

Full wwPDB X-ray Structure Validation Report (i)

Oct 3, 2023 – 02:15 AM EDT

PDB ID : 6PRD

Title : S. aureus dihydrofolate reductase co-crystallized with para-methoxyphenyl-di

methyoxydihydropthalazine inhibitor and NADP(H)

Authors: Bourne, C.R.; Thomas, L.M.

Deposited on : 2019-07-10

Resolution : 2.01 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*A user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : FAILED

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : FAILED

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.01 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 1568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dihydrofolate reductase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	162	Total 1335	C 862	N 225	O 245	S 3	0	3	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	A 159 LEU		-	expression tag	UNP P0A017
A	160 VAL		-	expression tag	UNP P0A017
A	A 161 PRO		-	expression tag	UNP P0A017
A	162	ARG	-	expression tag	UNP P0A017

• Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0


Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0


• Molecule 3 is $3-\{5-[(2,4-\text{diaminopyrimidin-5-yl})\text{methyl}]-2,3-\text{dimethoxyphenyl}\}-1-[(1S)-6,7-\text{dimethoxy-1-}(4-\text{methoxyphenyl})-3,4-\text{dihydrophthalazin-2}(1H)-yl]$ propan-1-one (three-letter code: OWG) (formula: $C_{33}H_{38}N_6O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	A	Aton	ns		ZeroOcc	AltConf
3	A	1	Total 45	C 33	N 6	O 6	0	0

• Molecule 4 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).





Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf
4	A	1	Total 48	C 21	_	O 17	P 3	0	0

• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	128	Total O 128 128	0	0

 $\operatorname{MolProbity}$ and EDS failed to run properly - this section is therefore empty.



3 Data and refinement statistics (i)

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants	79.06Å 79.06Å 105.51Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.79 - 2.01	Depositor
% Data completeness	96.7 (41.79-2.01)	Depositor
(in resolution range)	, , ,	•
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	11.96 (at 2.01Å)	Xtriage
Refinement program	PHENIX 1.8.4_1496	Depositor
R, R_{free}	0.190 , 0.226	Depositor
Wilson B-factor (\mathring{A}^2)	24.6	Xtriage
Anisotropy	0.375	Xtriage
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1568	wwPDB-VP
Average B, all atoms (\mathring{A}^2)	28.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.26% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $< L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

4 Model quality (i)

4.1 Standard geometry (i)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts (i)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains (i)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA (i)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Tuno	Chain	Res	Link	B	ond leng	gths	Bond angles		
MIOI	Type	e Chain Res Link		Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	401	-	5,5,5	0.35	0	5,5,5	0.74	0
3	OWG	A	403	-	47,49,49	2.19	14 (29%)	64,69,69	2.23	23 (35%)
2	GOL	A	402	-	5,5,5	0.33	0	5,5,5	0.29	0
4	NAP	A	404	-	45,52,52	1.10	3 (6%)	56,80,80	1.22	8 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	401	-	-	2/4/4/4	-
3	OWG	A	403	-	-	4/26/40/40	0/4/5/5
2	GOL	A	402	-	-	4/4/4/4	-
4	NAP	A	404	-	-	1/31/67/67	0/5/5/5

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(A)
3	A	403	OWG	C34-N35	6.17	1.46	1.33
3	A	403	OWG	C29-C28	4.63	1.47	1.38
3	A	403	OWG	C02-N01	4.46	1.45	1.34
3	A	403	OWG	C26-C27	-3.88	1.31	1.38
3	A	403	OWG	C29-C24	3.82	1.46	1.39
4	A	404	NAP	P2B-O2B	3.81	1.66	1.59
3	A	403	OWG	C26-C25	-3.79	1.33	1.39
3	A	403	OWG	C14-C13	3.29	1.58	1.51
3	A	403	OWG	C03-C02	-3.26	1.37	1.42
3	A	403	OWG	C15-C16	3.14	1.58	1.51
4	A	404	NAP	O4D-C1D	2.99	1.45	1.41
3	A	403	OWG	O47-C27	2.96	1.41	1.37
3	A	403	OWG	O08-C07	2.71	1.41	1.37
3	A	403	OWG	O45-C28	2.66	1.41	1.37
3	A	403	OWG	O79-C84	2.65	1.43	1.37
4	A	404	NAP	C5A-C4A	2.65	1.47	1.40
3	A	403	OWG	C24-C20	2.41	1.54	1.51



All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$Ideal(^{o})$
3	A	403	OWG	C46-O45-C28	-5.62	109.05	117.53
3	A	403	OWG	C14-C15-C16	-5.10	106.24	112.34
3	A	403	OWG	N33-C34-N36	-4.95	119.68	125.70
3	A	403	OWG	C28-C29-C24	-4.87	108.89	119.42
3	A	403	OWG	C29-C24-C25	4.28	124.18	119.39
3	A	403	OWG	C09-O08-C07	-4.19	111.20	117.53
3	A	403	OWG	C34-N36-C02	4.12	121.53	116.99
3	A	403	OWG	C19-C25-C24	-4.09	114.26	120.77
3	A	403	OWG	C48-O47-C27	-3.89	111.65	117.53
3	A	403	OWG	O45-C28-C29	-3.44	118.19	124.12
4	A	404	NAP	N3A-C2A-N1A	-3.41	123.36	128.68
3	A	403	OWG	C80-O79-C84	-2.99	111.03	117.51
3	A	403	OWG	N35-C34-N33	2.93	120.42	117.44
3	A	403	OWG	C69-C20-C24	-2.82	107.93	112.78
4	A	404	NAP	O3D-C3D-C2D	-2.61	103.37	111.82
3	A	403	OWG	C26-C27-C28	2.55	123.22	119.84
4	A	404	NAP	O2B-P2B-O1X	-2.54	99.60	109.39
3	A	403	OWG	O08-C07-C10	2.49	119.54	115.16
3	A	403	OWG	C20-N17-N18	2.46	122.00	118.93
3	A	403	OWG	O47-C27-C26	-2.45	119.90	124.12
3	A	403	OWG	O45-C28-C27	2.42	118.78	115.41
3	A	403	OWG	C29-C28-C27	2.41	123.03	119.84
4	A	404	NAP	C4A-C5A-N7A	-2.30	107.00	109.40
4	A	404	NAP	O2X-P2B-O1X	2.23	119.42	110.68
3	A	403	OWG	C32-C03-C02	2.23	118.76	115.72
4	A	404	NAP	O3X-P2B-O2X	2.16	115.89	107.64
4	A	404	NAP	C2A-N1A-C6A	2.12	122.39	118.75
3	A	403	OWG	C03-C32-N33	-2.08	120.35	123.82
4	A	404	NAP	C2D-C3D-C4D	2.04	106.61	102.64
3	A	403	OWG	C04-C03-C02	-2.03	119.36	122.17
3	A	403	OWG	C19-C25-C26	2.01	125.06	120.63

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	GOL	O1-C1-C2-C3
2	A	402	GOL	O1-C1-C2-C3
4	A	404	NAP	O4D-C1D-N1N-C6N
3	A	403	OWG	C27-C28-O45-C46
3	A	403	OWG	C29-C28-O45-C46

Continued on next page...



Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	A	402	GOL	O1-C1-C2-O2
2	A	402	GOL	C1-C2-C3-O3
2	A	401	GOL	O1-C1-C2-O2
3	A	403	OWG	C28-C27-O47-C48
3	A	403	OWG	C26-C27-O47-C48
2	A	402	GOL	O2-C2-C3-O3

There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



5 Fit of model and data (i)

5.1 Protein, DNA and RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates (i)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands (i)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers (i)

EDS failed to run properly - this section is therefore empty.

