



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 3, 2023 – 01:14 AM EDT

PDB ID : 6PSO
Title : Crystal structure of PsS1_19B C77S in complex with iota-neocarratetraose
Authors : Hettle, A.G.; Boraston, A.B.
Deposited on : 2019-07-13
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : **FAILED**
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : **FAILED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

There are no overall percentile quality scores available for this entry.

MolProbity and EDS failed to run properly - the sequence quality summary graphics cannot be shown.

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7555 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called exo-4S-kappa carrageenan S1 sulfatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	447	3545	2260	600	676	9	0	0	0
1	A	447	3544	2261	599	675	9	0	0	0

- Molecule 2 is an oligosaccharide called 3,6-anhydro-2-O-sulfo-alpha-D-galactopyranose-(1-3)-4-O-sulfo-beta-D-galactopyranose-(1-4)-3,6-anhydro-2-O-sulfo-alpha-D-galactopyranose-(1-3)-4-O-sulfo-beta-D-galactopyranose.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	O	S			
2	C	4	59	24	31	4	0	0	0
2	D	4	59	24	31	4	0	0	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
3	B	1	1	1	0	0
3	A	1	1	1	0	0

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Cl		
4	B	1	1	1	0	0
4	A	1	1	1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	140	Total 140	O 140	0	0
5	A	204	Total 204	O 204	0	0

MolProbity and EDS failed to run properly - this section is therefore empty.

3 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	65.26Å 94.00Å 170.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.01 – 2.00	Depositor
% Data completeness (in resolution range)	98.1 (29.01-2.00)	Depositor
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	10.73 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.8.0253	Depositor
R, R_{free}	0.197 , 0.231	Depositor
Wilson B-factor (Å ²)	24.3	Xtrriage
Anisotropy	0.058	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	7555	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

4 Model quality [i](#)

4.1 Standard geometry [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.2 Too-close contacts [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.2 Protein sidechains [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.3.3 RNA [i](#)

MolProbity failed to run properly - this section is therefore empty.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

8 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	G4S	C	1	2	16,16,16	1.26	2 (12%)	19,24,24	2.24	3 (15%)
2	DGS	C	2	2	15,15,16	1.28	1 (6%)	18,23,25	1.32	3 (16%)
2	G4S	C	3	2	15,15,16	1.36	2 (13%)	17,22,24	1.09	1 (5%)
2	DGS	C	4	2	15,15,16	1.48	2 (13%)	18,23,25	1.43	2 (11%)
2	G4S	D	1	2	16,16,16	1.28	2 (12%)	19,24,24	0.61	0
2	DGS	D	2	2	15,15,16	1.18	1 (6%)	18,23,25	1.72	3 (16%)
2	G4S	D	3	2	15,15,16	1.35	2 (13%)	17,22,24	1.30	2 (11%)
2	DGS	D	4	2	15,15,16	1.22	1 (6%)	18,23,25	1.11	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	G4S	C	1	2	-	1/7/27/27	0/1/1/1
2	DGS	C	2	2	-	0/5/27/30	0/3/2/2
2	G4S	C	3	2	-	1/7/24/27	0/1/1/1
2	DGS	C	4	2	-	0/5/27/30	0/3/2/2
2	G4S	D	1	2	-	1/7/27/27	0/1/1/1
2	DGS	D	2	2	-	0/5/27/30	0/3/2/2
2	G4S	D	3	2	-	1/7/24/27	0/1/1/1
2	DGS	D	4	2	-	0/5/27/30	0/3/2/2

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	1	G4S	O8-S	3.91	1.62	1.45
2	C	1	G4S	O7-S	3.78	1.61	1.45
2	D	3	G4S	O7-S	3.74	1.61	1.45
2	C	3	G4S	O8-S	3.74	1.61	1.45
2	C	4	DGS	O9-S	3.72	1.61	1.45
2	C	2	DGS	O2-C2	-3.59	1.41	1.47
2	D	2	DGS	O2-C2	-3.37	1.42	1.47
2	D	4	DGS	O2-C2	-3.09	1.42	1.47
2	C	4	DGS	O2-C2	-2.89	1.42	1.47
2	C	1	G4S	O4-C4	-2.24	1.41	1.46
2	D	3	G4S	O4-C4	-2.16	1.41	1.46
2	D	1	G4S	O4-C4	-2.07	1.42	1.46
2	C	3	G4S	O4-C4	-2.04	1.42	1.46

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1	G4S	O8-S-O7	-6.35	86.71	112.22
2	C	1	G4S	O9-S-O7	-5.70	88.68	108.49
2	D	2	DGS	O2-C2-C3	5.20	112.41	106.65
2	D	3	G4S	C1-C2-C3	4.02	114.61	109.67
2	C	3	G4S	C1-C2-C3	3.80	114.34	109.67
2	C	4	DGS	C2-O2-S	3.41	122.36	117.91
2	C	1	G4S	O9-S-O8	3.30	119.96	108.49
2	C	2	DGS	O5-C5-C6	-3.23	108.53	113.33
2	C	4	DGS	C2-C3-C4	-3.14	106.21	113.84
2	D	2	DGS	O5-C5-C6	-2.93	108.98	113.33
2	D	4	DGS	C2-C3-C4	-2.82	106.98	113.84
2	C	2	DGS	C2-O2-S	2.74	121.48	117.91
2	C	2	DGS	C2-C3-C4	-2.73	107.21	113.84
2	D	2	DGS	C2-O2-S	2.45	121.11	117.91
2	D	3	G4S	O3-C3-C2	-2.07	106.02	109.99

There are no chirality outliers.


All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	C	1	G4S	C4-O4-S-O8
2	C	3	G4S	O5-C5-C6-O6
2	D	3	G4S	O5-C5-C6-O6
2	D	1	G4S	C4-O4-S-O8

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





4.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

5 Fit of model and data [i](#)

5.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

5.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

5.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

5.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.