



## Full wwPDB EM Validation Report ⓘ

Mar 10, 2024 – 09:23 AM EDT

PDB ID : 6PSZ  
EMDB ID : EMD-20469  
Title : Poliovirus (Type 1 Mahoney), heat-catalysed 135S particle  
Authors : Hogle, J.M.; Filman, D.J.; Shah, P.N.M.  
Deposited on : 2019-07-14  
Resolution : 3.20 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
MapQ : 1.9.13  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	1	302	
2	2	272	
3	3	238	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10098 atoms, of which 5001 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VP1.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
1	1	193	3101	1013	1532	268	283	5	0	0

- Molecule 2 is a protein called VP2.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
2	2	222	3406	1107	1680	291	316	12	0	0

- Molecule 3 is a protein called VP3.

Mol	Chain	Residues	Atoms					AltConf	Trace	
			Total	C	H	N	O			S
3	3	231	3591	1150	1789	293	342	17	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: VP1



GLY LEU LEU GLN MET HIS MET LEU MET LEU SER MET ILE ASP ILE ASN THR VAL ARG F80 GLU THR VAL VAL GLY ALA ALA THR SER ARG ASP ALA LEU PRO PRO ASN THR D102 THR GLU ALA SER THR PRO HIS SER LYS GLU ILE PRO ALA LEU THR VAL VAL GLU THR THR ALA ASN PRO VAL VAL ASP THR

VAL GLN THR ARG HIS VAL VAL GLN H69 H70 H71 H72 S79 F80 F81 A82 R83 N94 P95 A96 S97 T98 M100 K101 K102 K103 D114 F124 T145 N146 M147 G148 H149 E168 D172 Y173 I183 I194 S213 LYS VAL PRO PRO LYS ASP GLN SER ALA ALA

LEU GLY ASP SER LEU TYR GLY ALA L234 L235 M235 H248 P263 K264 H265 I266 R272 G281 PRO GLY VAL ASP TYR LYS ASP GLY THR LEU THR PRO LEU SER THR LYS ASP LEU THR THR TYR


#### • Molecule 2: VP2



SER PRO ASN ILE GLU ALA CYS GLY TYR SER ASP ARG V113 L116 Q26 E27 A28 A29 R30 V33 R37 R43 ASP SER GLU ALA ASN PRO VAL ASP Q52 P53 T54 S55 P56 D57 V58 E74 L101 T107 Q119 G134 ASP SER ASN THR THR MET

HIS T143 G159 THR PHE THR PRO ASP ASN ASN GLN THR SER PRO PRO ALA ARG F174 D178 Y179 L180 M183 G184 T185 L186 L187 A190 R201 V209 V213 D219 S220 L234 M256 N261 R264 ASN ILE THR LEU PRO ARG LEU GLN

#### • Molecule 3: VP3



G1 T7 S10 D17 M18 D35 I36 P37 D56 L57 S58 A59 T60 K61 K62 N63 R69 D74 K75 P76 H77 D80 L87 A90 Y106 S123 K129 D140 K143 D157 I171 T174 I175 Y176 R177 Q178 T179 I180 D181 D182 S183 F184

T185 G188 I198 R206 E207 M208 N218 L225 R226 D227 I231 GLU GLN LYS ALA LEU ALA GLN

## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, I	Depositor
Number of particles used	67909	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	1.01	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	100.000	Depositor
Minimum map value	0.000	Depositor
Average map value	22.782	Depositor
Map value standard deviation	7.600	Depositor
Recommended contour level	50.0	Depositor
Map size (Å)	506.24, 506.24, 506.24	wwPDB
Map dimensions	448, 448, 448	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.13, 1.13, 1.13	Depositor

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	1	0.66	0/1618	0.61	0/2206
2	2	0.57	0/1773	0.65	0/2419
3	3	0.56	0/1849	0.65	0/2520
All	All	0.59	0/5240	0.64	0/7145

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	1569	1532	1525	13	0
2	2	1726	1680	1671	14	0
3	3	1802	1789	1784	17	0
All	All	5097	5001	4980	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:3:174:THR:OG1	3:3:176:TYR:O	2.02	0.78

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:2:201:ARG:NH1	3:3:123:SER:O	2.19	0.76
1:1:183:ILE:HD11	1:1:194:ILE:HG23	1.67	0.75
1:1:72:ARG:NH2	3:3:218:ASN:O	2.22	0.72
3:3:171:ILE:O	3:3:177:ARG:NH2	2.23	0.71
1:1:79:SER:O	1:1:83:ARG:NH1	2.26	0.69
2:2:107:THR:OG1	2:2:256:MET:SD	2.56	0.64
2:2:37:ARG:NH1	3:3:35:ASP:OD1	2.37	0.58
2:2:26:GLN:OE1	2:2:26:GLN:N	2.38	0.56
3:3:7:THR:O	3:3:10:SER:OG	2.22	0.54
3:3:198:ILE:HG21	3:3:208:MET:CE	2.38	0.53
1:1:263:PRO:O	1:1:266:ILE:HD11	2.08	0.53
2:2:178:ASP:OD1	2:2:179:TYR:N	2.42	0.53
2:2:219:ASP:OD1	2:2:220:SER:N	2.42	0.52
2:2:234:LEU:O	3:3:69:ARG:NE	2.39	0.51
2:2:183:ASN:O	2:2:185:THR:N	2.43	0.50
2:2:119:GLN:OE1	3:3:123:SER:N	2.45	0.50
3:3:106:TYR:O	3:3:225:LEU:HD12	2.12	0.49
1:1:124:PHE:O	1:1:272:ARG:N	2.44	0.49
2:2:213:VAL:HG22	3:3:37:PRO:HG2	1.93	0.49
1:1:147:ASN:O	1:1:149:HIS:N	2.44	0.49
3:3:179:THR:OG1	3:3:226:ARG:NH2	2.47	0.48
1:1:94:ASN:O	1:1:100:ASN:ND2	2.47	0.47
3:3:198:ILE:HG21	3:3:208:MET:HE3	1.98	0.45
3:3:60:THR:OG1	3:3:61:LYS:N	2.49	0.44
1:1:81:PHE:HB2	1:1:266:ILE:HD13	2.00	0.44
1:1:94:ASN:O	1:1:96:ALA:N	2.49	0.44
3:3:129:LYS:NZ	3:3:157:ASP:OD1	2.50	0.44
2:2:187:LEU:O	2:2:190:ALA:HB3	2.19	0.43
1:1:264:LYS:NZ	3:3:18:ASN:OD1	2.40	0.42
2:2:180:LEU:HD23	2:2:185:THR:O	2.20	0.42
1:1:172:ASP:OD1	1:1:173:TYR:N	2.51	0.42
1:1:101:LYS:NZ	1:1:102:ASP:OD2	2.34	0.41
3:3:87:LEU:HD13	3:3:188:GLY:HA3	2.03	0.41
2:2:101:LEU:N	2:2:261:ASN:O	2.44	0.41
2:2:33:VAL:HG22	2:2:209:VAL:HB	2.02	0.41
1:1:235:ASN:OD1	1:1:235:ASN:N	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	189/302 (63%)	163 (86%)	26 (14%)	0	100	100
2	2	214/272 (79%)	200 (94%)	14 (6%)	0	100	100
3	3	229/238 (96%)	211 (92%)	18 (8%)	0	100	100
All	All	632/812 (78%)	574 (91%)	58 (9%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	172/261 (66%)	172 (100%)	0	100	100
2	2	186/232 (80%)	186 (100%)	0	100	100
3	3	207/212 (98%)	206 (100%)	1 (0%)	88	95
All	All	565/705 (80%)	564 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	3	140	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:



Mol	Chain	Res	Type
1	1	153	GLN
1	1	248	HIS
2	2	196	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-20469. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



X




Y




Z


#### 6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

## 6.2 Central slices [i](#)


### 6.2.1 Primary map



X Index: 224




Y Index: 224




Z Index: 224


### 6.2.2 Raw map



X Index: 224



Y Index: 224




Z Index: 224


The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)


### 6.3.1 Primary map



X Index: 218




Y Index: 218




Z Index: 230


### 6.3.2 Raw map



X Index: 230



Y Index: 218




Z Index: 218


The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal standard-deviation projections (False-color) [i](#)


### 6.4.1 Primary map



X




Y




Z


### 6.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


## 6.5 Orthogonal surface views [i](#)

### 6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 50.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

### 6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


## 6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


### 6.6.1 emd\_20469\_msk\_1.map [i](#)



X



Y




Z

## 7 Map analysis [i](#)

This section contains the results of statistical analysis of the map.


### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.




## 7.2 Volume estimate [\(i\)](#)



The volume at the recommended contour level is  $1869 \text{ nm}^3$ ; this corresponds to an approximate mass of 1688 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum i




\*Reported resolution corresponds to spatial frequency of 0.312 Å<sup>-1</sup>

## 8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

### 8.1 FSC [i](#)



\*Reported resolution corresponds to spatial frequency of 0.312  $\text{\AA}^{-1}$

## 8.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.20	-	-
Author-provided FSC curve	3.22	3.54	3.29
Unmasked-calculated*	-	-	-


\*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-20469 and PDB model 6PSZ. Per-residue inclusion information can be found in section 3 on page 4.

### 9.1 Map-model overlays

#### 9.1.1 Map-model overlay [i](#)




#### 9.1.2 Map-model assembly overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 50.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

## 9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (50.0).









## 9.4 Atom inclusion [i](#)



At the recommended contour level, 71% of all backbone atoms, 71% of all non-hydrogen atoms, are inside the map.

## 9.5 Map-model fit summary

The table lists the average atom inclusion at the recommended contour level (50.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7130	 0.4360
1	 0.7320	 0.4420
2	 0.7100	 0.4360
3	 0.7040	 0.4320

