

wwPDB NMR Structure Validation Summary Report (i)

Sep 21, 2023 – 04:09 pm BST

PDB ID	:	8PSC
BMRB ID	:	34833
Title	:	Three-layered basket-type G-quadruplex from a G-rich sequence with five G-
		runs
Authors	:	Vianney, Y.M.; Schroeder, N.; Jana, J.; Weisz, K.
Deposited on	:	2023-07-13

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35.1

Clashscore

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 34%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metri	: Р	Percentile Ranks			
Clashscore			0		
	Worse		Better		
	Percentile relative to all structure	es			
	Percentile relative to all NMR st	tructures			
ЪЛани	Whole arch	ive NMR archiv	/e		
Metr	$\mathbf{c} (\# \text{Entries})$	s) (#Entries)			

158937

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

12864

Mol	Chain	Length	Quality of chain	
1		0.0		
	А	23	• 91%	•



2 Ensemble composition and analysis (i)

This entry contains 10 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 743 atoms, of which 256 are hydrogens and 0 are deuteriums.

• Molecule 1 is a DNA chain called DNA (5'-D(*AP*GP*(BG)P*GP*TP*AP*GP*GP*GP*GP*GP*CP*GP*GP*GP*GP*GP*GP*GP*GP*T)-3').

Mol	Chain	Residues		Atoms					Trace	
1	٨	02	Total	Br	С	Н	Ν	0	Р	0
	A	20	743	1	227	256	103	134	22	0



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: DNA (5'-D(*AP*GP*(BG)P*GP*TP*AP*GP*GP*GP*CP*GP*GP*CP*GP*GP*GP *GP*GP*CP*AP*GP*GP*GP*T)-3')

Chain A: •		91%	·
A1 62 63 65 65 69 69 69 610	<mark>611</mark> 612 615 615 617 617 621 723 723		

4.2 Residue scores for the representative (author defined) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.

• Molecule 1: DNA (5'-D(*AP*GP*(BG)P*GP*TP*AP*GP*GP*GP*CP*GP*GP*CP*GP*GP*GP *GP*GP*CP*AP*GP*GP*T)-3')

Chain A: .		78%	17%
A1 62 63 63 63 63 63 63 63 63	C10 C10 C12 C13 C13 C13 C13 C13 C13 C13 C13 C18 C18	A19 G21 G22 T23 T23	



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 10 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	structure calculation	18
Amber	refinement	18

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	159
Number of shifts mapped to atoms	159
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	34%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: BGM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	B	Sond lengths	Bond angles		
	Unam	RMSZ	$\#Z{>}5$	RMSZ	#Z>5	
1	А	$1.54{\pm}0.01$	$0{\pm}0/525~(~0.0{\pm}~0.0\%)$	2.42 ± 0.02	$39{\pm}2/813~(~4.8{\pm}~0.3\%)$	
All	All	1.54	0/5250 ($0.0%$)	2.42	388/8130~(~4.8%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	А	$0.0{\pm}0.0$	$2.9{\pm}1.0$
All	All	0	29

There are no bond-length outliers.

5 of 67 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	n Res Type Atoms Z Observed				Moo	dels		
	Unam	nes	Type	Atoms	Z	Observed(*)	Ideal(*)	Worst	Total
1	А	16	DG	O4'-C1'-N9	11.83	116.28	108.00	5	10
1	А	1	DA	N1-C6-N6	-10.07	112.56	118.60	4	10
1	А	19	DA	N1-C6-N6	-9.95	112.63	118.60	9	10
1	А	6	DA	N1-C6-N6	-9.49	112.91	118.60	6	10
1	А	21	DG	O4'-C1'-N9	8.67	114.07	108.00	7	9

There are no chirality outliers.

5 of 11 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	А	4	DG	Sidechain	10
				<u> </u>	1 /

Continued on next page...



001000	nucu jion	" pree	ous puge	• • •	
Mol	Chain	Res	Type	Group	Models (Total)
1	А	8	DG	Sidechain	4
1	А	10	DC	Sidechain	3
1	А	12	DG	Sidechain	2
1	А	14	DG	Sidechain	2

Continued from previous page...

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	4870	2560	2560	-

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is -.

There are no clashes.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

There are no protein molecules in this entry.

6.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds



that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mal	Turne	Chain	Dec	Tiple		Bond leng	gths
	туре	Unam	nes	Link	Counts	RMSZ	#Z>2
1	BGM	А	3	1	19,24,26	$1.08 {\pm} 0.01$	$1\pm0 (5\pm0\%)$

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mal	Turne	Chain	Dec	Tiple		Bond ang	gles
IVIOI	туре	Unam	nes	LIIIK	Counts	RMSZ	#Z>2
1	BGM	А	3	1	17,35,40	$1.01{\pm}0.04$	1±1 (4±3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	BGM	А	3	1	-	$0\pm 0,2,20,22$	$0\pm 0,3,3,3$

All unique bond outliers are listed below.

Mol	Chain	Ros	Type	Atoms	7	Observed(Å)	Ideal(Å)	Moo	
	Ullalli	Ites	Type	Atoms		Observed(A)	Ideal(A)	Worst	Total
1	А	3	BGM	C8-N7	2.93	1.30	1.34	2	10

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mal	Chain	Dog	Type	Atoms	7	Observed(°)	$Ideal(^{o})$	Moo	lels
	Ullalli	nes	туре	Atoms		Observed()	Ideal()	Worst	Total
1	А	3	BGM	C4'-O4'-C1'	2.15	104.26	109.45	5	7
1	А	3	BGM	C2'-C1'-N9	2.09	113.73	116.01	3	1



There are no chirality outliers. There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 34% for the well-defined parts and 34% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: assigned_chem_shift_list

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	159
Number of shifts mapped to atoms	159
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 34%, i.e. 152 atoms were assigned a chemical shift out of a possible 450. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathrm{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Sugar	91/264~(34%)	91/154~(59%)	0/110~(0%)	0/0 (%)
Base	61/186~(33%)	40/120~(33%)	21/30~(70%)	0/36~(0%)
Overall	152/450~(34%)	131/274~(48%)	21/140~(15%)	0/36~(0%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.



7.1.5 Random Coil Index (RCI) plots (i)

No random coil index (RCI) plot could be generated from the current chemical shift list. RCI is only applicable to proteins



8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	237
Intra-residue (i-j =0)	75
Sequential (i-j =1)	80
Medium range ($ i-j >1$ and $ i-j <5$)	8
Long range $(i-j \ge 5)$	26
Inter-chain	0
Hydrogen bond restraints	48
Disulfide bond restraints	0
Total dihedral-angle restraints	208
Number of unmapped restraints	17
Number of restraints per residue	19.3
Number of long range restraints per residue ¹	2.9

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	0.4	0.15
0.2-0.5 (Medium)	None	None
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	Max ($^{\circ}$)
1.0-10.0 (Small)	9.3	9.94
10.0-20.0 (Medium)	2.5	18.96
>20.0 (Large)	1.2	49.42



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destruction to the second	Count	$\%^1$	Vio	lated	3	Consis	$\begin{array}{c c c c c c c c c c c c c c c c c c c $	
Restraints type	Count	701	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	75	31.6	1	1.3	0.4	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	74	31.2	1	1.4	0.4	0	0.0	0.0
Sidechain-Sidechain	1	0.4	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	80	33.8	4	5.0	1.7	0	0.0	0.0
Backbone-Backbone	1	0.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	54	22.8	3	5.6	1.3	0	0.0	0.0
Sidechain-Sidechain	25	10.5	1	4.0	0.4	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	8	3.4	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	2	0.8	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	6	2.5	0	0.0	0.0	0	0.0	0.0
Long range $(i-j \ge 5)$	26	11.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	7	3.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	19	8.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	48	20.3	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	237	100.0	5	2.1	2.1	0	0.0	0.0
Backbone-Backbone	1	0.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	137	57.8	4	2.9	1.7	0	0.0	0.0
Sidechain-Sidechain	99	41.8	1	1.0	0.4	0	0.0	0.0

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models





9.1.1 Bar chart : Distribution of distance restraints and violations (i)

Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.


Model ID		Nur	nber o	f viola	ations	5	Mean (Å)	Max (Å)	SD^6 (Å)	Median (Å)
Model ID	IR^1	SQ^2	MR^3	LR ⁴	IC ⁵	Total	Mean (A)	Max (A)	SD (A)	Median (A)
1	0	0	0	0	0	0	0.0	0.0	0.0	0.0
2	0	0	0	0	0	0	0.0	0.0	0.0	0.0
3	0	1	0	0	0	1	0.13	0.13	0.0	0.13
4	0	0	0	0	0	0	0.0	0.0	0.0	0.0
5	0	1	0	0	0	1	0.11	0.11	0.0	0.11
6	0	1	0	0	0	1	0.15	0.15	0.0	0.15
7	0	0	0	0	0	0	0.0	0.0	0.0	0.0
8	0	0	0	0	0	0	0.0	0.0	0.0	0.0
9	0	0	0	0	0	0	0.0	0.0	0.0	0.0
10	1	1	0	0	0	2	0.11	0.11	0.0	0.11

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,



8PSC

⁵Inter-chain restraints, ⁶Standard deviation



9.2.1 Bar graph : Distance Violation statistics for each model (i)

The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 184(IR:74, SQ:76, MR:8, LR:26, IC:0) restraints are not violated in the ensemble.

Nu	mber	of vio	lated	aints	Fractio	n of the ensemble	
IR^1	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Count^6	%
1	4	0	0	0	5	1	10.0
0	0	0	0	0	0	2	20.0
0	0	0	0	0	0	3	30.0
0	0	0	0	0	0	4	40.0


Continued on next page...



00100	Continued from previous page												
Nu	\mathbf{mber}	of vio	lated	aints	Fraction of the ensemble								
IR^1	SQ^2	MR^3	LR^4	IC ⁵	Total	Count^6	%						
0	0	0	0	0	0	5	50.0						
0	0	0	0	0	0	6	60.0						
0	0	0	0	0	0	7	70.0						
0	0	0	0	0	0	8	80.0						
0	0	0	0	0	0	9	90.0						
0	0	0	0	0	0	10	100.0						

Continued from previous page...

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations



9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

No violations found



9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.

Data insufficient to plot histogram

9.5.2 Table : All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,140)	1:14:A:DG:H8	1:13:A:DC:H2"	6	0.15
(1,111)	1:12:A:DG:H8	1:11:A:DG:H3'	3	0.13
(1,121)	1:19:A:DA:H8	1:20:A:DG:H8	10	0.11
(1,28)	1:12:A:DG:H8	1:11:A:DG:H1'	5	0.11
(1,110)	1:11:A:DG:H8	1:11:A:DG:H3'	10	0.1




10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

Angle tripe	Count	$\%^1$	Vi	olated	3	Consis	tently	$^{\prime}$ Violated ⁴
Angle type	Count	70	Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
NU4	15	7.2	10	66.7	4.8	2	13.3	1.0
CHI	22	10.6	0	0.0	0.0	0	0.0	0.0
NU1	15	7.2	2	13.3	1.0	0	0.0	0.0
UNNAMED	111	53.4	0	0.0	0.0	0	0.0	0.0
NU2	15	7.2	3	20.0	1.4	0	0.0	0.0
NU3	15	7.2	8	53.3	3.8	1	6.7	0.5
NU0	15	7.2	11	73.3	5.3	2	13.3	1.0
Total	208	100.0	34	16.3	16.3	5	2.4	2.4

 1 percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models








Violated and consistently violated restraints are shown using different hatch patterns in their respective categories

10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID				Number of vi	olations	8			Mean (°)	Max (°)	SD $(^{\circ})$	Median (°)
Model ID	NU4	CHI	NU1	UNNAMED	NU2	NU3	NU0	Total	Mean ()	Max ()	SD ()	Median ()
1	4	0	1	0	1	3	5	14	9.63	31.13	9.24	7.12
2	4	0	0	0	1	4	6	15	7.03	15.85	4.66	5.59
3	4	0	0	0	0	1	4	9	4.34	9.2	2.59	3.32
4	6	0	1	0	2	3	8	20	11.06	37.0	9.84	7.89
5	4	0	1	0	0	2	4	11	7.63	16.46	4.88	6.34
6	3	0	0	0	0	1	3	7	8.5	15.73	3.43	7.25
7	7	0	1	0	1	4	5	18	10.6	41.14	11.33	5.77
8	5	0	0	0	0	2	4	11	5.15	11.74	2.84	4.92
9	5	0	1	Ö	1	2	5	14	11.52	35.38	11.17	7.56
10	3	0	1	0	1	2	4	11	17.13	49.42	15.5	9.14

10.2.1 Bar graph : Dihedral violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated




		Num	Fraction	n of the ensemble					
NU4	CHI	NU1	UNNAMED	NU2	NU3	NU0	Total	Count^1	%
1	0	0	0	0	2	2	5	1	10.0
2	0	0	0	2	3	2	9	2	20.0
4	0	2	0	1	2	3	12	3	30.0
0	0	0	0	0	0	0	0	4	40.0
0	0	0	0	0	0	1	1	5	50.0
0	0	0	0	0	0	0	0	6	60.0
0	0	0	0	0	0	0	0	7	70.0
1	0	0	0	0	0	1	2	8	80.0
0	0	0	0	0	0	0	0	9	90.0
2	0	0	0	0	1	2	5	10	100.0

restraints for a given fraction of ensemble.

¹ Number of models with violations

10.3.1	Bar graph :	Dihedral-angle	Violation	statistics	for the	ensemble	(\mathbf{i})
--------	-------------	----------------	-----------	------------	---------	----------	----------------




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram : Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models



in the ensemble



10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,78)	1:16:A:DG:C3'	1:16:A:DG:C4'	1:16:A:DG:O4'	1:16:A:DG:C1'	10	14.14	2.22	14.28
(1,74)	1:16:A:DG:C4'	1:16:A:DG:O4'	1:16:A:DG:C1'	1:16:A:DG:C2'	10	9.09	1.09	9.22
(1,77)	1:16:A:DG:C2'	1:16:A:DG:C3'	1:16:A:DG:C4'	1:16:A:DG:O4'	10	7.9	2.68	8.51
(1,93)	1:20:A:DG:C3'	1:20:A:DG:C4'	1:20:A:DG:O4'	1:20:A:DG:C1'	10	6.36	1.53	6.34
(1,89)	1:20:A:DG:C4'	1:20:A:DG:O4'	1:20:A:DG:C1'	1:20:A:DG:C2'	10	4.86	0.92	4.82
(1,43)	1:7:A:DG:C3'	1:7:A:DG:C4'	1:7:A:DG:O4'	1:7:A:DG:C1'	8	6.89	6.27	4.16
(1,39)	1:7:A:DG:C4'	1:7:A:DG:O4'	1:7:A:DG:C1'	1:7:A:DG:C2'	8	4.46	1.55	4.31
(1,59)	1:12:A:DG:C4'	1:12:A:DG:O4'	1:12:A:DG:C1'	1:12:A:DG:C2'	5	10.97	7.16	12.33
(1,56)	1:11:A:DG:C1'	1:11:A:DG:C2'	1:11:A:DG:C3'	1:11:A:DG:C4'	3	38.86	3.3	41.14
(1,55)	1:11:A:DG:O4'	1:11:A:DG:C1'	1:11:A:DG:C2'	1:11:A:DG:C3'	3	31.15	13.97	28.54


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)



10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



10.5.2 Table: All violated dihedral-angle restraints (i)

The following table provides the list of violations for the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation ($^{\circ}$)
(1,55)	1:11:A:DG:O4'	1:11:A:DG:C1'	1:11:A:DG:C2'	1:11:A:DG:C3'	10	49.42
(1,56)	1:11:A:DG:C1'	1:11:A:DG:C2'	1:11:A:DG:C3'	1:11:A:DG:C4'	10	41.24
(1,56)	1:11:A:DG:C1'	1:11:A:DG:C2'	1:11:A:DG:C3'	1:11:A:DG:C4'	7	41.14
(1,60)	1:12:A:DG:O4'	1:12:A:DG:C1'	1:12:A:DG:C2'	1:12:A:DG:C3'	4	37.0
(1,57)	1:11:A:DG:C2'	1:11:A:DG:C3'	1:11:A:DG:C4'	1:11:A:DG:O4'	9	35.38
(1,61)	1:12:A:DG:C1'	1:12:A:DG:C2'	1:12:A:DG:C3'	1:12:A:DG:C4'	4	34.71
(1,56)	1:11:A:DG:C1'	1:11:A:DG:C2'	1:11:A:DG:C3'	1:11:A:DG:C4'	9	34.2
(1,57)	1:11:A:DG:C2'	1:11:A:DG:C3'	1:11:A:DG:C4'	1:11:A:DG:O4'	7	34.19
(1,54)	1:11:A:DG:C4'	1:11:A:DG:O4'	1:11:A:DG:C1'	1:11:A:DG:C2'	10	33.04
(1,60)	1:12:A:DG:O4'	1:12:A:DG:C1'	1:12:A:DG:C2'	1:12:A:DG:C3'	1	31.13

