

Full wwPDB EM Validation Report (i)

Dec 20, 2023 – 04:34 pm GMT

PDB ID : 8PSX

EMDB ID : EMD-17864

Title : Tilapia Lake Virus polymerase in vRNA elongation state (transcriptase con-

formation)

Authors: Arragain, B.; Cusack, S.

Deposited on : 2023-07-13

Resolution : 2.96 Å(reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/EMValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev70

Mogul : 1.8.4, CSD as541be (2020)

MolProbity : 4.02b-467 buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

MapQ: 1.9.9

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	${ m EM\ structures} \ (\#{ m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion <40%). The numeric value is given above the bar.

Mol	Chain	Length		(Quality of chair	n		
1	A	419	21%		92%		7%	•
2	В	519	12%		87%		12%	
3	С	478	36%	76	%	12%	11%	-
4	S	40	28%	5%		68%		
4	V	40	10% 8% •		80%			-
5	Р	21	19%	57%	5%	38%		_



2 Entry composition (i)

There are 8 unique types of molecules in this entry. The entry contains 22416 atoms, of which 11016 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Polymerase acidic protein (PA-like).

Mol	Chain	Residues			Atom	.S			AltConf	Trace
1	A	417	Total 6652	C 2105	H 3314	N 597	O 617	S 19	0	0

• Molecule 2 is a protein called Putative PB1.

Mol	Chain	Residues			Atom	.S			AltConf	Trace
9	D	513	Total	С	Н	N	О	S	9	0
	Б	313	7958	2491	3987	688	762	30	2	U

• Molecule 3 is a protein called RNA-dependent RNA polymerase.

\mathbf{N}	Iol	Chain	Residues			Atom	.S			AltConf	Trace
,	3	С	426	Total 6676	C 2117	H 3339	N 596	O 610	S 14	0	0

There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	391	LYS	ARG	conflict	UNP A0A7G3S745
С	458	GLY	-	expression tag	UNP A0A7G3S745
С	459	SER	-	expression tag	UNP A0A7G3S745
С	460	GLY	-	expression tag	UNP A0A7G3S745
С	461	SER	-	expression tag	UNP A0A7G3S745
С	462	GLU	-	expression tag	UNP A0A7G3S745
С	463	ASN	-	expression tag	UNP A0A7G3S745
С	464	LEU	-	expression tag	UNP A0A7G3S745
С	465	TYR	-	expression tag	UNP A0A7G3S745
С	466	PHE	-	expression tag	UNP A0A7G3S745
С	467	GLN	-	expression tag	UNP A0A7G3S745
С	468	GLY	-	expression tag	UNP A0A7G3S745
С	469	HIS	-	expression tag	UNP A0A7G3S745
С	470	HIS	-	expression tag	UNP A0A7G3S745
С	471	HIS	-	expression tag	UNP A0A7G3S745



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
С	472	HIS	-	expression tag	UNP A0A7G3S745
С	473	HIS	-	expression tag	UNP A0A7G3S745
С	474	HIS	-	expression tag	UNP A0A7G3S745
С	475	HIS	-	expression tag	UNP A0A7G3S745
С	476	HIS	-	expression tag	UNP A0A7G3S745
С	477	HIS	-	expression tag	UNP A0A7G3S745
С	478	HIS	-	expression tag	UNP A0A7G3S745

• Molecule 4 is a RNA chain called 5' vRNA end - vRNA loop (40-mer).

Mol	Chain	Residues	Atoms					AltConf	Trace	
1	V	Q	Total	С	Н	N	О	Р	0	0
4	v	8	250	75	84	26	57	8	U	U
1	C	13	Total	С	Н	N	О	Р	0	0
4	S	10	422	126	140	53	90	13	U	U

• Molecule 5 is a RNA chain called Transcription-like product.


Mol	Chain	Residues			Atom	.S			AltConf	Trace
5	Р	13	Total 411	C 123	H 139	N 46	O 90	P 13	0	0

• Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	AltConf
6	A	1	Total Zn 1 1	0
6	С	2	Total Zn 2 2	0

• Molecule 7 is [(2 {R},3 {S},4 {R},5 {R})-5-(4-azanyl-2-oxidanylidene-pyrimidin-1-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methoxy- {N}-[oxidanyl(phosphonooxy)phosphoryl]phosphonami dic acid (three-letter code: A0I) (formula: $C_9H_{17}N_4O_{13}P_3$) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		A	Aton	ns			AltConf
7	D	1	Total	С	Н	N	О	Р	0
'	Б	1	42	9	13	4	13	3	0

 \bullet Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	AltConf
8	В	2	Total Mg 2 2	0




3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polymerase acidic protein (PA-like)










Chain P: 57% 5% 38%





4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	29892	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{Å}^2)$	40	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	0.081	Depositor
Minimum map value	-0.048	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.002	Depositor
Recommended contour level	0.02	Depositor
Map size (Å)	251.99998, 251.99998, 251.99998	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.84, 0.84, 0.84	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: A0I, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.26	0/3413	0.51	0/4611
2	В	0.25	0/4043	0.50	0/5472
3	С	0.25	0/3412	0.49	0/4608
4	S	0.26	0/316	0.73	0/491
4	V	0.21	0/184	0.70	0/283
5	Р	0.25	0/302	0.76	0/464
All	All	0.25	0/11670	0.52	0/15929

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3338	3314	3313	17	0
2	В	3971	3987	3984	38	0
3	С	3337	3339	3335	21	0
4	S	282	140	140	2	0
4	V	166	84	85	1	0
5	Р	272	139	140	1	0
6	A	1	0	0	0	0



Continued from previous page...

	Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
	6	С	2	0	0	0	0
ſ	7	В	29	13	0	0	0
	8	В	2	0	0	0	0
	All	All	11400	11016	10997	72	0

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is 3.

All (72) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
2 D = 4 CED 0 C	4 C 22 4 OP4	distance (Å)	overlap (Å)
2:B:74:SER:OG	4:S:29:A:OP1	1.88	0.90
2:B:335:THR:OG1	2:B:337:ASP:OD1	1.92	0.87
3:C:83:ARG:NH2	4:S:36:U:OP1	2.16	0.79
2:B:199:ALA:HB1	2:B:292:ILE:HG21	1.70	0.72
2:B:103:THR:OG1	2:B:125:ILE:O	2.06	0.70
3:C:204:LYS:NZ	3:C:228:VAL:O	2.27	0.68
3:C:195:VAL:O	3:C:202:ALA:N	2.28	0.66
2:B:36:GLU:O	2:B:39:THR:HG22	1.95	0.65
1:A:226:GLU:OE1	1:A:244:SER:OG	2.09	0.64
1:A:395:THR:O	1:A:399:SER:OG	2.16	0.62
2:B:418:ILE:HD11	2:B:428:CYS:HB3	1.79	0.62
2:B:401:THR:OG1	2:B:412:LEU:O	2.19	0.61
1:A:74:ASN:OD1	1:A:80:VAL:HG12	2.01	0.60
3:C:356:SER:O	3:C:359:ARG:NH1	2.34	0.59
2:B:427:ASP:OD1	2:B:427:ASP:O	2.21	0.59
2:B:506:ASN:HB2	2:B:509:ILE:HD12	1.83	0.59
2:B:39:THR:HG23	2:B:41:LYS:HG2	1.83	0.59
2:B:199:ALA:CB	2:B:292:ILE:HG21	2.32	0.58
3:C:334:THR:HG21	3:C:342:CYS:HB3	1.87	0.56
1:A:322:LEU:HD11	2:B:383:LEU:HD11	1.89	0.55
2:B:103:THR:HG21	2:B:111:LEU:HD13	1.89	0.54
1:A:101:ASP:N	1:A:101:ASP:OD1	2.40	0.53
3:C:361:ILE:HD12	3:C:363:ILE:HD11	1.92	0.52
2:B:189:THR:HG22	2:B:189:THR:O	2.10	0.52
3:C:90:GLU:OE1	3:C:90:GLU:N	2.42	0.52
2:B:465:LEU:HD23	2:B:465:LEU:O	2.09	0.52
1:A:45:TYR:OH	1:A:50:ASN:OD1	2.28	0.50
1:A:79:TYR:O	1:A:87:GLN:NE2	2.44	0.50
1:A:128:SER:O	2:B:44:ASN:ND2	2.38	0.50
2:B:438:ASP:OD1	3:C:129:ARG:NH1	2.42	0.50
		Continu	ed on nert nage



 $Continued\ from\ previous\ page...$

Continuea from previo		Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
2:B:213[B]:ASP:OD1	2:B:290:ASP:OD1	2.30	0.49
2:B:357:VAL:HG13	2:B:444:ASP:HB2	1.93	0.49
2:B:57:GLN:NE2	5:P:5:U:OP1	2.41	0.49
3:C:247:PRO:O	3:C:248:THR:HG23	2.13	0.48
2:B:323:SER:OG	2:B:326:SER:O	2.31	0.48
1:A:86:TYR:O	1:A:90:CYS:SG	2.71	0.48
2:B:103:THR:O	2:B:103:THR:HG23	2.12	0.48
2:B:506:ASN:CB	2:B:509:ILE:HD12	2.43	0.47
2:B:78:VAL:HG13	2:B:85:VAL:HG13	1.96	0.47
2:B:53:LYS:NZ	3:C:355:ARG:NH2	2.64	0.46
1:A:144:ASP:N	1:A:144:ASP:OD1	2.49	0.46
2:B:479:SER:OG	2:B:482:ASP:OD1	2.32	0.46
2:B:214:CYS:HB2	2:B:289:ASP:HB2	1.98	0.46
2:B:378:HIS:O	2:B:381:GLU:HG3	2.16	0.46
3:C:177:ILE:HG12	3:C:178:LYS:N	2.31	0.46
3:C:235:CYS:O	3:C:238:VAL:HG22	2.15	0.45
3:C:26:ASP:OD1	3:C:26:ASP:N	2.43	0.45
1:A:148:GLU:HG3	1:A:405:LEU:HD13	1.99	0.45
3:C:82:PHE:O	3:C:84:PRO:HD2	2.17	0.44
3:C:318:GLY:O	3:C:323:ARG:NH1	2.50	0.44
2:B:114:LYS:O	2:B:118:VAL:HG13	2.17	0.44
3:C:171:ASP:OD1	3:C:172:SER:N	2.51	0.44
4:V:9:U:OP1	4:V:10:U:H2'	2.18	0.43
1:A:43:ASP:OD2	1:A:62:ARG:NH1	2.47	0.43
2:B:167:LEU:HB2	2:B:242:LEU:HD13	2.01	0.43
2:B:103:THR:HG21	2:B:111:LEU:CD1	2.50	0.42
2:B:299:ALA:O	2:B:303:GLU:HG3	2.20	0.42
3:C:357:PRO:HB2	3:C:358:GLU:OE2	2.20	0.42
1:A:19:SER:O	1:A:19:SER:OG	2.34	0.42
3:C:176:VAL:HG12	3:C:185:MET:HG2	2.02	0.42
3:C:195:VAL:HG22	3:C:196:GLU:N	2.34	0.42
1:A:201:ARG:NH2	1:A:263:LYS:O	2.42	0.41
1:A:320:TYR:CZ	2:B:13:LEU:HD12	2.55	0.41
2:B:168:SER:HA	2:B:171:VAL:HG22	2.02	0.41
1:A:314:VAL:O	1:A:318:GLU:HG2	2.21	0.41
3:C:191:HIS:CE1	3:C:233:CYS:SG	3.14	0.41
2:B:292:ILE:O	2:B:292:ILE:HG23	2.20	0.41
2:B:289:ASP:O	2:B:289:ASP:CG	2.58	0.40
2:B:418:ILE:HD13	2:B:423:LEU:HB2	2.03	0.40
1:A:72:ILE:HB	1:A:80:VAL:HG13	2.04	0.40
2:B:285:LEU:HB3	2:B:292:ILE:HG22	2.04	0.40



Continued from previous page...

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
3:C:100:ASP:O	3:C:104:VAL:HG23	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	415/419 (99%)	403 (97%)	12 (3%)	0	100	100
2	В	511/519 (98%)	486 (95%)	25 (5%)	0	100	100
3	С	422/478 (88%)	401 (95%)	21 (5%)	0	100	100
All	All	1348/1416 (95%)	1290 (96%)	58 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	364/365 (100%)	354 (97%)	10 (3%)	44 74
2	В	445/447 (100%)	432 (97%)	13 (3%)	42 73
3	С	358/405 (88%)	331 (92%)	27 (8%)	13 40
All	All	1167/1217 (96%)	1117 (96%)	50 (4%)	33 62



All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	17	THR
1	A	31	THR THR
1	A	66	LYS
1	A	79	TYR
1	A	101	LYS TYR ASP
1	A A A A A B	144	ASP
1	A	176	GLU
1	A	380	ASP
1	A	399	SER
2	В	63	ARG
2	В	73	ARG
2	В	154	SER
2 2 2 2 2	В	159	SER LEU
2	В	180	SER SER
2	В	183	SER
2	В	255	ASP ASP LEU ASP
2	В	306	ASP
2	В	327	LEU
2	В	371	ASP
2 2 2 2 2 2	В	482	ASP
2	В	495	LYS
2	В	507	TYR HIS
3	B C C C C	22	HIS
3	С	39	ARG
3	С	45	ARG
3	С	67	SER
3	С	78	SER
3	С	81	SER
3	С	87	THR
3	C C C C C C C C	88	GLN
3	С	89	GLU
3	С	93	ASP
3	С	111	ASP ARG
3	С	120	SER
3	С	187	ASP
3	С	196	GLU
3	С	197	ASP
3	С	199	LYS
3	С	207	SER
3	С	237	LYS
3	С	268	ASP



Continued from previous page...

Mol	Chain	Res	Type
3	С	316	SER
3	С	346	LYS
3	С	358	GLU
3	С	377	PHE
3	С	388	SER
3	С	396	GLU
3	С	411	LYS
3	С	413	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	87	GLN
2	В	313	HIS

5.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
4	S	12/40 (30%)	0	0
4	V	7/40 (17%)	3 (42%)	0
5	Р	11/21 (52%)	0	0
All	All	30/101 (29%)	3 (10%)	0

All (3) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
4	V	5	A
4	V	7	С
4	V	10	U

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

Of 6 ligands modelled in this entry, 5 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Pog	Link	Bo	ond leng	ths	Bond angles		
MIOI	Туре	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	A0I	В	601	8	29,30,30	1.03	4 (13%)	42,47,47	1.08	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	A0I	В	601	8	-	3/19/38/38	0/2/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
7	В	601	A0I	P2-O5	2.73	1.50	1.46
7	В	601	A0I	P1-O11	2.70	1.50	1.46
7	В	601	A0I	P2-O6	-2.33	1.50	1.56
7	В	601	A0I	P1-O4	-2.32	1.50	1.56

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
7	В	601	A0I	O6-P2-O5	4.22	118.77	109.92
7	В	601	A0I	O4-P1-O11	4.01	118.32	109.92
7	В	601	A0I	O3-P1-O11	-2.34	105.22	114.24

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	В	601	A0I	P2-N4-P1-O3




Continued from previous page...

Mol	Chain	Res	Type	Atoms
7	В	601	A0I	P1-N4-P2-O5
7	В	601	A0I	P1-N4-P2-O7

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.




6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-17864. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections (i)

6.1.1 Primary map



6.1.2 Raw map




The images above show the map projected in three orthogonal directions.




6.2 Central slices (i)

Primary map 6.2.1








Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150

The images above show central slices of the map in three orthogonal directions.




6.3 Largest variance slices (i)

6.3.1 Primary map



6.3.2 Raw map




The images above show the largest variance slices of the map in three orthogonal directions.




6.4 Orthogonal standard-deviation projections (False-color) (i)

6.4.1 Primary map



6.4.2 Raw map



The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.




6.5 Orthogonal surface views (i)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.02. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.




6.6 Mask visualisation (i)


This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure

6.6.1 emd_17864_msk_1.map (i)








7 Map analysis (i)

This section contains the results of statistical analysis of the map.


7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)




The volume at the recommended contour level is $30~\mathrm{nm^3}$; this corresponds to an approximate mass of $27~\mathrm{kDa}$.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)




^{*}Reported resolution corresponds to spatial frequency of 0.338 $\rm \mathring{A}^{-1}$



8 Fourier-Shell correlation (i)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC (i)



^{*}Reported resolution corresponds to spatial frequency of 0.338 $\rm \mathring{A}^{-1}$



8.2 Resolution estimates (i)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)			
rtesolution estimate (A)	0.143	0.5	Half-bit	
Reported by author	2.96	-	-	
Author-provided FSC curve	2.93	3.44	2.96	
Unmasked-calculated*	3.66	7.16	3.78	


^{*}Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.66 differs from the reported value 2.96 by more than 10 %



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-17864 and PDB model 8PSX. Per-residue inclusion information can be found in section 3 on page 6.


9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.02 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.




9.2 Q-score mapped to coordinate model (i)



The images above show the model with each residue coloured according its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.


9.3 Atom inclusion mapped to coordinate model (i)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.02).



9.4 Atom inclusion (i)



At the recommended contour level, 60% of all backbone atoms, 61% of all non-hydrogen atoms, are inside the map.



9.5 Map-model fit summary (i)

The table lists the average atom inclusion at the recommended contour level (0.02) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.6090	0.5950
A	0.6290	0.5980
В	0.7020	0.6150
С	0.4950	0.5730
Р	0.5880	0.5530
S	0.7230	0.5750
V	0.6990	0.6060



