



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 12:11 pm GMT

PDB ID : 8PWA
Title : Crystal structure of the human METTL3-METTL14 in complex with a bisubstrate analogue (BA4)
Authors : Bedi, R.K.; Etheve-Quellejeu, M.; Caffisch, A.
Deposited on : 2023-07-19
Resolution : 2.10 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	228	
2	B	290	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 3742 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called N6-adenosine-methyltransferase catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	202	1604	1029	279	286	10	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	353	MET	-	initiating methionine	UNP Q86U44


- Molecule 2 is a protein called N6-adenosine-methyltransferase non-catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	238	1876	1205	317	341	13	0	1	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	106	GLY	-	expression tag	UNP A4IFD8

- Molecule 3 is (2 {S})-4-[[[2 {R},3 {S},4 {R},5 {R}]-5-(6-aminopurin-9-yl)-3,4-bis(oxidanyl)oxolan-2-yl]methyl-[3-[[9-[(2 {R},5 {R}]-5-(hydroxymethyl)-3,4-bis(oxidanyl)oxolan-2-yl]-7 {H}-purin-6-yl]amino]propyl]amino]-2-azanyl-butanoic acid (three-letter code: I0C) (formula:) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
			Total	C	N			O
3	A	1	39	22	12	5	0	0

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
4	B	1	4	2	2	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	84	Total	O	0	0
			84	84		
6	B	134	Total	O	0	0
			134	134		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: N6-adenosine-methyltransferase catalytic subunit

Chain A: 



- Molecule 2: N6-adenosine-methyltransferase non-catalytic subunit

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	63.89Å 63.89Å 225.18Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.54 – 2.10 44.54 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.7 (44.54-2.10) 99.7 (44.54-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.15 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.191 , 0.225 0.190 , 0.225	Depositor DCC
R_{free} test set	1606 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	38.9	Xtrriage
Anisotropy	0.084	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 50.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.030 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3742	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: I0C, MG, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/1649	0.64	0/2246
2	B	0.42	0/1923	0.63	1/2614 (0.0%)
All	All	0.42	0/3572	0.64	1/4860 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	395	LEU	CB-CG-CD1	-5.89	100.98	111.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1604	0	1543	9	0
2	B	1876	0	1774	21	0
3	A	39	0	0	1	0
4	B	4	0	3	0	0
5	B	1	0	0	0	0
6	A	84	0	0	2	0
6	B	134	0	0	8	0
All	All	3742	0	3320	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:701:I0C:CBS	3:A:701:I0C:OBC	1.63	1.17
1:A:375:CYS:SG	6:A:877:HOH:O	2.35	0.84
2:B:276:ASP:OD1	6:B:501:HOH:O	1.97	0.81
2:B:153:ILE:N	6:B:504:HOH:O	2.20	0.75
2:B:168:MET:HE3	2:B:369:TYR:HA	1.71	0.72
2:B:250:LYS:NZ	6:B:506:HOH:O	2.25	0.69
2:B:229:SER:OG	6:B:502:HOH:O	2.15	0.64
1:A:391:VAL:HB	1:A:530:LYS:HG2	1.81	0.62
2:B:174:ILE:O	6:B:503:HOH:O	2.17	0.59
2:B:185:LYS:NZ	6:B:509:HOH:O	2.39	0.56
1:A:373:TRP:HB2	1:A:551:LEU:HD13	1.87	0.54
1:A:456:ILE:HG12	2:B:285:LYS:HE2	1.91	0.53
2:B:178:ASP:O	2:B:181:GLU:HG3	2.08	0.53
2:B:326:LYS:NZ	6:B:510:HOH:O	2.41	0.52
2:B:260[B]:CYS:SG	2:B:287:HIS:CE1	3.06	0.49
1:A:480:LYS:HE2	2:B:260[B]:CYS:SG	2.55	0.47
2:B:260[A]:CYS:SG	2:B:285:LYS:HD2	2.55	0.47
1:A:490:ASN:HB2	6:A:875:HOH:O	2.15	0.46
2:B:173:ASP:HB3	6:B:557:HOH:O	2.15	0.45
1:A:453:ASP:OD2	2:B:280:VAL:N	2.46	0.44
2:B:120:CYS:O	2:B:124:VAL:HG23	2.18	0.44
1:A:478:HIS:HE1	2:B:257:GLU:OE1	2.01	0.43
2:B:177:PHE:CE2	2:B:179:ILE:HA	2.54	0.42
2:B:395:LEU:HA	2:B:395:LEU:HD13	1.72	0.42
1:A:456:ILE:HD13	1:A:456:ILE:HA	1.91	0.41
2:B:230:PHE:CE1	2:B:339:LEU:HD22	2.56	0.41
2:B:341:ARG:HD3	2:B:385:LEU:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	199/228 (87%)	193 (97%)	4 (2%)	2 (1%)	15	11
2	B	229/290 (79%)	226 (99%)	3 (1%)	0	100	100
All	All	428/518 (83%)	419 (98%)	7 (2%)	2 (0%)	29	26

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	404	LEU
1	A	403	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/200 (84%)	169 (100%)	0	100	100
2	B	196/258 (76%)	194 (99%)	2 (1%)	76	82
All	All	365/458 (80%)	363 (100%)	2 (0%)	86	92

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	136	ASP
2	B	277	PRO

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	478	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	I0C	A	701	-	36,43,53	4.58	8 (22%)	35,61,77	2.13	10 (28%)
4	ACT	B	401	-	3,3,3	1.48	1 (33%)	3,3,3	1.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	I0C	A	701	-	-	3/20/40/62	0/5/5/6

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	701	I0C	CBO-CBS	-18.08	1.26	1.53
3	A	701	I0C	OBC-CBS	16.36	1.63	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	701	I0C	CBM-CBQ	-6.99	1.35	1.53
3	A	701	I0C	CBF-NBA	5.86	1.45	1.34
3	A	701	I0C	CBO-CBM	5.14	1.67	1.53
3	A	701	I0C	C6-N6	2.74	1.44	1.34
3	A	701	I0C	OBC-CBQ	2.63	1.50	1.45
4	B	401	ACT	CH3-C	2.11	1.57	1.49
3	A	701	I0C	CBJ-NAX	-2.05	1.34	1.37

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	701	I0C	CAK-NAV-CBF	5.16	121.02	116.59
3	A	701	I0C	N3-C2-N1	-4.87	121.07	128.68
3	A	701	I0C	NAX-CAK-NAV	-4.38	121.83	128.68
3	A	701	I0C	OBC-CBS-CBO	-3.92	101.19	106.93
3	A	701	I0C	CAN-CAR-NBT	-3.09	106.04	113.84
3	A	701	I0C	CAK-NAX-CBJ	3.02	120.52	113.45
3	A	701	I0C	CBJ-CBH-NAZ	-2.98	106.30	109.40
3	A	701	I0C	C4-C5-N7	-2.66	106.62	109.40
3	A	701	I0C	CAP-NBA-CBF	-2.31	119.01	122.89
3	A	701	I0C	OBC-CBQ-CAT	-2.00	105.58	108.90

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	701	I0C	CAR-CAN-CAP-NBA
3	A	701	I0C	C-CA-CB-CAS
3	A	701	I0C	CAP-CAN-CAR-NBT


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	701	I0C	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.

Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	202/228 (88%)	0.12	14 (6%) 16 21	28, 41, 69, 96	0
2	B	238/290 (82%)	0.11	9 (3%) 40 46	26, 40, 72, 90	0
All	All	440/518 (84%)	0.11	23 (5%) 27 32	26, 40, 72, 96	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	402	MET	7.4
1	A	401	HIS	6.5
1	A	574	ILE	6.3
1	A	573	ILE	5.4
2	B	274	THR	4.9
2	B	309	VAL	4.8
1	A	570	PRO	4.4
2	B	155	LEU	4.4
1	A	406	TYR	4.3
1	A	404	LEU	4.0
2	B	395	LEU	3.9
1	A	463	LEU	3.9
2	B	394	ARG	3.8
2	B	392	ILE	3.2
1	A	575	SER	3.2
1	A	405	PRO	3.1
1	A	572	GLY	3.1
1	A	571	ASP	2.9
1	A	403	GLU	2.9
2	B	136	ASP	2.4
2	B	159	LEU	2.2
1	A	369	PHE	2.2
2	B	393	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	I0C	A	701	39/48	0.91	0.13	33,42,109,127	0
4	ACT	B	401	4/4	0.98	0.10	30,32,35,37	0
5	MG	B	402	1/1	0.98	0.12	48,48,48,48	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.