



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 12, 2022 – 02:10 pm GMT

PDB ID : 7PXQ
Title : GH115 alpha-1,2-glucuronidase D303A
Authors : Wilkens, C.; Morth, J.P.; Polikarpov, I.
Deposited on : 2021-10-08
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.24
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0267
CCP4 : 7.1.010 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.24


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	841	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6982 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called xylan alpha-1,2-glucuronidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	821	6708	4262	1122	1300	24	0	0	0

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	3	Total	Ca	0	0
			3	3		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	271	Total	O	0	0
			271	271		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: xylan alpha-1,2-glucuronidase



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	148.15Å 148.15Å 274.46Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	50.47 – 2.30 65.18 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.47-2.30) 100.0 (65.18-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.190 , 0.219 0.185 , 0.211	Depositor DCC
R_{free} test set	4045 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	66.9	Xtriage
Anisotropy	0.194	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6982	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.42	0/6879	0.60	0/9341

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6708	0	6417	44	0
2	A	3	0	0	0	0
3	A	271	0	0	6	0
All	All	6982	0	6417	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:423:ILE:HG23	1:A:458:LEU:HD22	1.69	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:773:VAL:O	1:A:777:THR:HG23	1.87	0.74
1:A:423:ILE:HD12	1:A:423:ILE:H	1.55	0.71
1:A:667:VAL:HB	1:A:817:VAL:HG13	1.77	0.67
1:A:659:VAL:HG12	1:A:715:ARG:HB3	1.77	0.66
1:A:268:ASP:OD1	1:A:270:THR:OG1	2.12	0.66
1:A:103:VAL:HG13	1:A:115:ILE:HD13	1.80	0.63
1:A:30:GLU:OE2	3:A:1001:HOH:O	2.16	0.61
1:A:423:ILE:HG22	1:A:474:LEU:HD22	1.87	0.57
1:A:666:VAL:HG13	1:A:668:VAL:HG23	1.89	0.55
1:A:423:ILE:H	1:A:423:ILE:CD1	2.18	0.54
1:A:423:ILE:HG12	1:A:454:TYR:HE2	1.73	0.53
1:A:423:ILE:HD12	1:A:423:ILE:N	2.25	0.52
1:A:60:LEU:O	1:A:62:ASP:N	2.42	0.51
1:A:392:GLU:OE1	1:A:392:GLU:N	2.41	0.50
1:A:583:ARG:HH11	1:A:583:ARG:HB3	1.76	0.50
1:A:837:ILE:O	1:A:838:GLU:HB2	2.10	0.50
1:A:719:PRO:HB2	1:A:721:GLU:HG2	1.94	0.49
1:A:177:ASP:HA	1:A:220:TRP:CZ2	2.48	0.48
1:A:287:VAL:HG12	1:A:290:THR:OG1	2.14	0.48
1:A:587:PRO:HG3	1:A:654:ALA:HB3	1.96	0.48
1:A:422:PRO:HG2	1:A:425:ASN:ND2	2.29	0.47
1:A:262:ASP:OD1	1:A:262:ASP:N	2.45	0.47
1:A:555:LYS:HA	1:A:559:LEU:HB3	1.95	0.47
1:A:681:GLU:O	1:A:682:GLU:HB2	2.14	0.47
1:A:25:ASN:HD22	1:A:56:PRO:HD2	1.80	0.46
1:A:46:ASP:OD1	1:A:49:ARG:NH2	2.48	0.46
1:A:788:THR:O	3:A:1002:HOH:O	2.20	0.46
1:A:17:LEU:O	1:A:24:THR:HG23	2.17	0.45
1:A:835:LYS:HB2	1:A:835:LYS:HE2	1.65	0.44
1:A:716:MET:SD	1:A:817:VAL:HG11	2.58	0.44
1:A:423:ILE:CG1	1:A:454:TYR:HE2	2.28	0.44
1:A:732:LYS:NZ	3:A:1004:HOH:O	2.32	0.43
1:A:733:SER:HA	1:A:783:VAL:HG23	2.00	0.43
1:A:754:LYS:HE3	1:A:754:LYS:HB2	1.91	0.43
1:A:53:THR:HG22	3:A:1042:HOH:O	2.19	0.43
1:A:219:MET:HG3	3:A:1102:HOH:O	2.19	0.42
1:A:171:ARG:HG2	1:A:463:ALA:HB1	2.01	0.42
1:A:583:ARG:HH11	1:A:583:ARG:CB	2.33	0.41
1:A:666:VAL:CG1	1:A:668:VAL:HG23	2.51	0.41
1:A:741:LYS:HE2	1:A:742:GLY:O	2.21	0.41
1:A:16:THR:O	1:A:24:THR:HG21	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:ARG:HD2	3:A:1222:HOH:O	2.21	0.41
1:A:85:LYS:O	1:A:86:ILE:HD13	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	817/841 (97%)	784 (96%)	32 (4%)	1 (0%)	51 64

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	61	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	721/736 (98%)	689 (96%)	32 (4%)	28 39

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	24	THR
1	A	30	GLU
1	A	53	THR
1	A	103	VAL
1	A	107	LEU
1	A	219	MET
1	A	264	TYR
1	A	272	ASN
1	A	286	ARG
1	A	287	VAL
1	A	289	ASP
1	A	325	ARG
1	A	336	GLU
1	A	345	SER
1	A	467	THR
1	A	468	ARG
1	A	475	LEU
1	A	501	TYR
1	A	525	TRP
1	A	532	TYR
1	A	627	LYS
1	A	666	VAL
1	A	687	VAL
1	A	741	LYS
1	A	744	GLU
1	A	753	SER
1	A	773	VAL
1	A	777	THR
1	A	817	VAL
1	A	835	LYS
1	A	836	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	821/841 (97%)	0.79	78 (9%) 8 11	50, 79, 121, 151	0

All (78) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	264	TYR	6.2
1	A	269	TYR	6.0
1	A	791	LEU	5.7
1	A	103	VAL	5.4
1	A	266	VAL	5.2
1	A	325	ARG	5.1
1	A	327	LEU	5.1
1	A	751	GLY	4.9
1	A	86	ILE	4.7
1	A	792	HIS	4.7
1	A	274	GLU	4.6
1	A	284	ILE	4.5
1	A	330	ILE	4.5
1	A	257	TRP	4.4
1	A	275	VAL	4.0
1	A	259	ARG	3.9
1	A	73	GLU	3.8
1	A	260	ASN	3.7
1	A	261	ARG	3.7
1	A	336	GLU	3.6
1	A	797	GLY	3.6
1	A	267	TRP	3.6
1	A	280	PHE	3.5
1	A	262	ASP	3.5
1	A	753	SER	3.4
1	A	89	ASP	3.4
1	A	263	LYS	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	161	TYR	3.3
1	A	332	LYS	3.2
1	A	279	PHE	3.1
1	A	283	GLY	3.1
1	A	799	TYR	3.1
1	A	276	ILE	3.0
1	A	18	LEU	3.0
1	A	250	MET	2.9
1	A	345	SER	2.9
1	A	84	ARG	2.9
1	A	278	GLN	2.9
1	A	533	ILE	2.8
1	A	268	ASP	2.8
1	A	592	TRP	2.7
1	A	9	GLU	2.6
1	A	525	TRP	2.6
1	A	333	ASN	2.6
1	A	326	LEU	2.5
1	A	219	MET	2.5
1	A	798	PHE	2.5
1	A	252	ARG	2.5
1	A	521	GLU	2.5
1	A	334	GLN	2.5
1	A	340	ASP	2.5
1	A	87	LYS	2.4
1	A	528	VAL	2.4
1	A	725	VAL	2.4
1	A	573	TYR	2.3
1	A	540	LEU	2.3
1	A	790	ASP	2.3
1	A	322	ASP	2.3
1	A	7	TYR	2.3
1	A	690	TYR	2.3
1	A	562	VAL	2.3
1	A	281	ARG	2.2
1	A	518	TYR	2.2
1	A	577	MET	2.2
1	A	341	VAL	2.2
1	A	806	LEU	2.2
1	A	158	ARG	2.2
1	A	99	LEU	2.2
1	A	520	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	532	TYR	2.2
1	A	55	ALA	2.2
1	A	77	ILE	2.2
1	A	517	THR	2.1
1	A	537	ALA	2.1
1	A	194	TYR	2.1
1	A	338	ILE	2.0
1	A	581	LEU	2.0
1	A	6	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)


There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	CA	A	902	1/1	0.92	0.05	105,105,105,105	0
2	CA	A	903	1/1	0.93	0.06	81,81,81,81	0
2	CA	A	901	1/1	1.00	0.15	63,63,63,63	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around CA A 903:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.