

wwPDB EM Validation Summary Report (i)

Feb 21, 2022 - 05:17 am GMT

PDB ID 7Q0A : EMDB ID EMD-13742 : Title : SARS-CoV-2 Spike ectodomain with Fab FI3A Authors Duyvesteyn, H.M.E.; Ren, J.; Stuart, D.I. : Deposited on 2021-10-14 : 4.80 Å(reported) Resolution : Based on initial model 7ND5 ·

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	0.0.0 dev97
Mogul	:	1.8.4, CSD as541be (2020)
MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.26

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 4.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	EM structures
INTEGI IC	$(\# {\rm Entries})$	$(\# { m Entries})$
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for $\geq=3, 2, 1$ and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq=5\%$ The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion < 40%). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	Quality of chain							
1	А	1316	67%	8%	24%						
1	В	1316	65%	10%	25%						
1	С	1316	68%	8%	25%						
2	Y	116	61% 87%		13%						
3	Z	107	71%		12%						
4	D	2	50%	50%							
4	Е	2	100%								
4	F	2	50%	50%							

Continued on next page...



Contr	nued fron	<i>i</i> previous	page
Mol	Chain	Length	Quality of chain
	â	_	
4	G	2	100%
4	т	0	
4	1	2	100%
1	т	2	
- 4	1	<u>∠</u>	100%
4	K	2	E00/
-	11		8, 0C
4	М	2	100%
4	Ν	2	100%
4	0	2	100%
	Ъ		
4	P	2	100%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 25125 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

•	Molecule 1	is a	protein	called	Spike	glycoprot	ein,Spike	ectodom	ain,Spike	protein	S2'.
•	monecule 1	no a	protein	canca	opine	SIJCOPIO	,opine	cciouom	am,opine	proton	04.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	А	995	Total 7628	C 4883	N 1266	0 1444	${ m S}\ 35$	0	0
1	В	984	Total 7530	C 4818	N 1249	O 1428	S 35	0	0
1	С	991	Total 7582	C 4853	N 1264	0 1431	S 34	0	0

Chain	Residue	Modelled	Actual	Comment	Reference
А	682	GLY	ARG	conflict	UNP P0DTC2
А	683	SER	ARG	conflict	UNP P0DTC2
А	685	SER	ARG	conflict	UNP P0DTC2
А	986	PRO	LYS	conflict	UNP P0DTC2
А	987	PRO	VAL	conflict	UNP P0DTC2
В	682	GLY	ARG	conflict	UNP P0DTC2
В	683	SER	ARG	conflict	UNP P0DTC2
В	685	SER	ARG	conflict	UNP P0DTC2
В	986	PRO	LYS	conflict	UNP P0DTC2
В	987	PRO	VAL	conflict	UNP P0DTC2
С	682	GLY	ARG	conflict	UNP P0DTC2
С	683	SER	ARG	conflict	UNP P0DTC2
С	685	SER	ARG	conflict	UNP P0DTC2
С	986	PRO	LYS	conflict	UNP P0DTC2
С	987	PRO	VAL	conflict	UNP P0DTC2

There are 15 discrepancies between the modelled and reference sequences:

• Molecule 2 is a protein called FI3A fab heavy chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	Y	116	Total 884	C 552	N 155	0 171	${ m S}{ m 6}$	0	0

• Molecule 3 is a protein called FI3A fab Light chain.



Mol	Chain	Residues	Atoms					AltConf	Trace
3	Z	107	Total 815	C 511	N 134	0 167	${ m S} { m 3}$	0	0

• Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	A	Aton	ns		AltConf	Trace	
4	р	n	Total	С	Ν	0	0	0	
4	D	2	28	16	2	10	0	0	
4	Б	ე	Total	С	Ν	0	0	0	
4	E	2	28	16	2	10	0	0	
4	Б	ე	Total	С	Ν	0	0	0	
4	Г	2	28	16	2	10	0	0	
4	С	n	Total	С	Ν	0	0	0	
4	G	2	28	16	2	10	0	0	
4	Т	ე	Total	С	Ν	0	0	0	
4	1	2	28	16	2	10	0	0	
4	т	ე	Total	С	Ν	0	0	0	
4	J	2	28	16	2	10	0	0	
4	K	9	Total	С	Ν	0	0	0	
4	Γ	2	28	16	2	10	0	0	
4	М	9	Total	С	Ν	0	0	0	
4	111	2	28	16	2	10	0	0	
4	N	n	Total	С	Ν	0	0	0	
4	11	2	28	16	2	10	0	0	
4	0	9	Total	С	Ν	0	0	0	
4	U	2	28	16	2	10	U	U	
4	Р	D 9		С	Ν	0	0	0	
±	1	2	28	16	2	10	U	0	

• Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	1	Aton	ns		AltConf	
F	٨	1	Total	С	Ν	0	0	
0	A	1	112	64	8	40	0	
۲.	Δ	1	Total	С	Ν	0	0	
0	A	L	112	64	8	40	0	
5	Λ	1	Total	С	Ν	0	0	
0	Л	T	112	64	8	40	0	
5	Δ	1	Total	С	Ν	0	0	
0	Π	T	112	64	8	40	0	
5	Δ	1	Total	С	Ν	0	0	
0	11	I	112	64	8	40	0	
5	Δ	1	Total	С	Ν	0	0	
0	11	I	112	64	8	40	0	
5	Δ	1	Total	С	Ν	Ο	0	
		T	112	64	8	40	0	
5	А	1	Total	С	Ν	0	0	
		1	112	64	8	40	0	
5	В	1	Total	С	Ν	Ο	0	
		±	154	88	11	55	Ŭ	
5	В	1	Total	С	Ν	0	0	
		1	154	88	11	55	0	
5	В	1	Total	С	Ν	Ο	0	
		1	154	88	11	55	0	
5	В	1	Total	С	Ν	Ο	0	
		1	154	88	11	55	0	
5	В	1	Total	С	Ν	Ο	0	
	-	-	154	88	11	55		
5	В	1	Total	С	Ν	0	0	
		_ *	154	88	11	55		

Continued on next page...



Mol	Chain	Residues		Aton	ns		AltConf	
E	D	1	Total	С	Ν	0	0	
5	D	1	154	88	11	55	0	
F	D	1	Total	С	Ν	0	0	
5	D	1	154	88	11	55	0	
5	В	1	Total	С	Ν	0	0	
5	D	1	154	88	11	55	0	
5	В	1	Total	С	Ν	0	0	
0	D	1	154	88	11	55	0	
5	В	1	Total	С	Ν	0	0	
5	D	1	154	88	11	55	0	
5	С	1	Total	С	Ν	0	0	
5	U	1	112	64	8	40	0	
5	С	1	Total	С	Ν	0	0	
0	0	I	112	64	8	40	0	
5	С	1	Total	С	Ν	0	0	
		1	112	64	8	40	0	
5	С	1	Total	С	Ν	0	0	
0	0	I	112	64	8	40	0	
5	С	1	Total	С	Ν	0	0	
0	0	1	112	64	8	40	0	
5	С	1	Total	С	Ν	0	0	
0	0	1	112	64	8	40	0	
5	С	1	Total	С	Ν	0	0	
		1	112	64	8	40		
5	С	1	Total	С	Ν	0	0	
0		1	112	64	8	40		

Continued from previous page...



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Spike glycoprotein, Spike ectodomain, Spike protein S2'

 • Molecule 1: Spike glycoprotein,
Spike ectodomain, Spike protein $\mathrm{S2'}$

Chain B:

65%

10%

25%





LYS TYR PHE LYS ASN HIS

• Molecule 1: Spike glycoprotein, Spike ectodomain, Spike protein S2'







opyranose

Chain E:

100%



NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:	50%	50%	
NAG1 NAG2			
• Molecule 4 opyranose	: 2-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain G:	10	00%	•
NAG1 NAG2			
• Molecule 4 opyranose	: 2-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain I:	100)%	
NAG1 NAG2			
• Molecule 4 opyranose	: 2-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain J:	10	0%	
NAG1 NAG2			
• Molecule 4 opyranose	: 2-acetamido-2-deoxy-beta-I	D-glucopyranose-(1-4)-2-acetamid	o-2-deoxy-beta-D-gluc
Chain K:	50%	50%	•
-			

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain M:

100%

NAG1 NAG2



• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain N:

100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain O:

100%

NAG1 NAG2

• Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain P:

100%

NAG1 NAG2



4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, C1	Depositor
Number of particles used	9095	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	47.7	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	1.092	Depositor
Minimum map value	-0.517	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.022	Depositor
Recommended contour level	0.0261	Depositor
Map size (Å)	459.19998, 459.19998, 459.19998	wwPDB
Map dimensions	280, 280, 280	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.64, 1.64, 1.64	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
IVI01	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.26	0/7800	0.48	0/10634
1	В	0.26	0/7698	0.48	0/10494
1	С	0.26	0/7754	0.47	0/10577
2	Y	0.27	0/902	0.52	0/1221
3	Ζ	0.25	0/832	0.49	0/1131
All	All	0.26	0/24986	0.48	0/34057

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	7628	0	7354	66	0
1	В	7530	0	7213	84	0
1	С	7582	0	7280	62	0
2	Y	884	0	859	9	0
3	Ζ	815	0	789	8	0
4	D	28	0	25	2	0
4	Е	28	0	25	0	0
4	F	28	0	25	1	0

Continued on next page...



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	G	28	0	25	0	0
4	Ι	28	0	25	0	0
4	J	28	0	25	0	0
4	Κ	28	0	25	1	0
4	М	28	0	25	0	0
4	N	28	0	25	0	0
4	0	28	0	25	0	0
4	Р	28	0	25	0	0
5	А	112	0	104	0	0
5	В	154	0	143	1	0
5	С	112	0	104	1	0
All	All	25125	0	24121	209	0

Continued from previous page...

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 209 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1129:VAL:HB	1:B:1132:ILE:HD11	1.61	0.82
1:B:1126:CYS:HB2	1:B:1132:ILE:HD13	1.59	0.82
1:B:1040:VAL:HG21	1:C:1035:GLY:HA3	1.67	0.76
1:B:388:ASN:HB3	1:B:527:PRO:HD2	1.69	0.74
1:A:736:VAL:HG11	1:A:1004:LEU:HD21	1.72	0.71

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	nalysed Favoured Allowed		Outliers	Percentiles
1	А	977/1316~(74%)	946 (97%)	31 (3%)	0	100 100

Continued on next page...



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	В	962/1316~(73%)	915~(95%)	46~(5%)	1 (0%)	51	85
1	С	975/1316~(74%)	939~(96%)	36~(4%)	0	100	100
2	Y	114/116~(98%)	110 (96%)	3~(3%)	1 (1%)	17	56
3	Ζ	105/107~(98%)	100~(95%)	5 (5%)	0	100	100
All	All	3133/4171 (75%)	3010 (96%)	121 (4%)	2~(0%)	54	85

Continued from previous page...

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	Y	102	PRO
1	В	332	ILE

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	834/1139~(73%)	829~(99%)	5 (1%)	86	92
1	В	819/1139~(72%)	815 (100%)	4 (0%)	88	93
1	С	822/1139~(72%)	819 (100%)	3~(0%)	91	94
2	Y	95/95~(100%)	94~(99%)	1 (1%)	73	85
3	Ζ	92/92~(100%)	92 (100%)	0	100	100
All	All	2662/3604~(74%)	2649 (100%)	13 (0%)	89	93

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	643	PHE
1	В	1017	GLU
2	Y	98	ASP
1	С	517	LEU
1	С	616	ASN



Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

22 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	L Type Chain Reg Link		Bo	Bond lengths			Bond angles			
IVIOI	Type	Ullalli	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
4	NAG	D	1	1,4	14,14,15	0.27	0	$17,\!19,\!21$	0.47	0
4	NAG	D	2	4	14,14,15	0.26	0	17,19,21	0.43	0
4	NAG	Е	1	1,4	14,14,15	0.23	0	17,19,21	0.40	0
4	NAG	Е	2	4	14,14,15	0.24	0	17,19,21	0.44	0
4	NAG	F	1	1,4	14,14,15	0.20	0	17,19,21	0.42	0
4	NAG	F	2	4	14,14,15	0.25	0	17,19,21	0.41	0
4	NAG	G	1	1,4	14,14,15	0.22	0	17,19,21	0.41	0
4	NAG	G	2	4	14,14,15	0.23	0	17,19,21	0.42	0
4	NAG	Ι	1	1,4	14,14,15	0.20	0	17,19,21	0.43	0
4	NAG	Ι	2	4	14,14,15	0.24	0	17,19,21	0.41	0
4	NAG	J	1	1,4	14,14,15	0.20	0	17,19,21	0.42	0
4	NAG	J	2	4	14,14,15	0.24	0	$17,\!19,\!21$	0.41	0
4	NAG	K	1	1,4	14,14,15	0.18	0	17,19,21	0.42	0
4	NAG	K	2	4	14,14,15	0.23	0	$17,\!19,\!21$	0.41	0
4	NAG	М	1	1,4	14,14,15	0.24	0	$17,\!19,\!21$	0.44	0
4	NAG	М	2	4	14,14,15	0.21	0	17,19,21	0.40	0
4	NAG	N	1	1,4	14,14,15	0.26	0	$17,\!19,\!21$	0.47	0
4	NAG	N	2	4	14,14,15	0.30	0	17,19,21	0.40	0
4	NAG	0	1	1,4	14,14,15	0.20	0	17,19,21	0.44	0



Mal	Mal Type Chain Beg	Dec	Tink	Bo	ond leng	$_{\rm ths}$	Bond angles			
WIOI	туре	Unam	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NAG	0	2	4	14,14,15	0.27	0	17,19,21	0.41	0
4	NAG	Р	1	1,4	14,14,15	0.19	0	17,19,21	0.47	0
4	NAG	Р	2	4	14,14,15	0.27	0	17,19,21	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	D	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	D	2	4	_	1/6/23/26	0/1/1/1
4	NAG	Е	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	Е	2	4	-	1/6/23/26	0/1/1/1
4	NAG	F	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	F	2	4	-	0/6/23/26	0/1/1/1
4	NAG	G	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	G	2	4	-	2/6/23/26	0/1/1/1
4	NAG	Ι	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	Ι	2	4	-	2/6/23/26	0/1/1/1
4	NAG	J	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	J	2	4	-	2/6/23/26	0/1/1/1
4	NAG	K	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	K	2	4	-	0/6/23/26	0/1/1/1
4	NAG	М	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	М	2	4	-	1/6/23/26	0/1/1/1
4	NAG	Ν	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	Ν	2	4	-	2/6/23/26	0/1/1/1
4	NAG	0	1	1,4	-	2/6/23/26	0/1/1/1
4	NAG	0	2	4	-	0/6/23/26	0/1/1/1
4	NAG	Р	1	1,4	-	0/6/23/26	0/1/1/1
4	NAG	Р	2	4	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 27 torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
4	Ι	1	NAG	O5-C5-C6-O6
4	Ν	2	NAG	O5-C5-C6-O6
4	G	1	NAG	O5-C5-C6-O6
4	G	2	NAG	C4-C5-C6-O6
4	Ι	1	NAG	C4-C5-C6-O6

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Κ	1	NAG	1	0
4	F	1	NAG	1	0
4	D	1	NAG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.













































5.6 Ligand geometry (i)

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Type	when Chain Reg Lin		Tink	Link Bond lengths			Bond angles		
WIOI	туре	Ullalli	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
5	NAG	С	1407	1	14,14,15	0.38	0	17,19,21	0.51	0
5	NAG	В	1408	1	14,14,15	0.22	0	17,19,21	0.41	0
5	NAG	В	1409	1	14,14,15	0.23	0	17,19,21	0.38	0



Mal	Turne	Chain	Dec	Timle	Bo	ond leng	ths	В	ond ang	gles
IVIOI	Type	Chain	nes		Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	В	1407	1	14,14,15	0.21	0	17,19,21	0.47	0
5	NAG	С	1403	1	14,14,15	0.22	0	17,19,21	0.50	0
5	NAG	В	1403	1	14,14,15	0.20	0	17,19,21	0.37	0
5	NAG	А	1402	1	14,14,15	0.34	0	17,19,21	0.44	0
5	NAG	В	1406	1	14,14,15	0.20	0	17,19,21	0.36	0
5	NAG	С	1401	1	14,14,15	0.22	0	17,19,21	0.40	0
5	NAG	А	1401	1	14,14,15	0.25	0	17,19,21	0.35	0
5	NAG	А	1407	1	14,14,15	0.22	0	17,19,21	0.41	0
5	NAG	А	1403	1	14,14,15	0.35	0	17,19,21	0.38	0
5	NAG	С	1408	1	14,14,15	0.25	0	17,19,21	0.44	0
5	NAG	В	1401	1	14,14,15	0.27	0	17,19,21	0.46	0
5	NAG	В	1405	1	14,14,15	0.31	0	17,19,21	0.36	0
5	NAG	А	1408	1	14,14,15	0.26	0	17,19,21	0.57	0
5	NAG	А	1405	1	14,14,15	0.29	0	17,19,21	0.54	0
5	NAG	В	1410	1	14,14,15	0.33	0	17,19,21	0.47	0
5	NAG	С	1404	1	14,14,15	0.21	0	17,19,21	0.43	0
5	NAG	С	1402	1	14,14,15	0.23	0	17,19,21	0.36	0
5	NAG	В	1404	1	14,14,15	0.27	0	17,19,21	0.41	0
5	NAG	С	1406	1	14,14,15	0.23	0	17,19,21	0.40	0
5	NAG	А	1404	1	14,14,15	0.20	0	17,19,21	0.42	0
5	NAG	С	1405	1	14,14,15	0.21	0	17,19,21	0.41	0
5	NAG	А	1406	1	14,14,15	0.21	0	17,19,21	0.43	0
5	NAG	В	1402	1	14,14,15	0.21	0	17,19,21	0.46	0
5	NAG	В	1411	1	14,14,15	0.22	0	17,19,21	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	С	1407	1	-	2/6/23/26	0/1/1/1
5	NAG	В	1408	1	-	0/6/23/26	0/1/1/1
5	NAG	В	1409	1	-	0/6/23/26	0/1/1/1
5	NAG	В	1407	1	-	2/6/23/26	0/1/1/1
5	NAG	С	1403	1	-	2/6/23/26	0/1/1/1
5	NAG	В	1403	1	-	2/6/23/26	0/1/1/1
5	NAG	А	1402	1	-	2/6/23/26	0/1/1/1
5	NAG	В	1406	1	-	1/6/23/26	0/1/1/1
5	NAG	С	1401	1	-	2/6/23/26	0/1/1/1
5	NAG	А	1401	1	-	1/6/23/26	0/1/1/1

Continued on next page...



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	А	1407	1	-	2/6/23/26	0/1/1/1
5	NAG	А	1403	1	-	2/6/23/26	0/1/1/1
5	NAG	С	1408	1	-	0/6/23/26	0/1/1/1
5	NAG	В	1401	1	-	2/6/23/26	0/1/1/1
5	NAG	В	1405	1	-	2/6/23/26	0/1/1/1
5	NAG	А	1408	1	-	1/6/23/26	0/1/1/1
5	NAG	А	1405	1	-	2/6/23/26	0/1/1/1
5	NAG	В	1410	1	-	2/6/23/26	0/1/1/1
5	NAG	С	1404	1	-	2/6/23/26	0/1/1/1
5	NAG	С	1402	1	-	2/6/23/26	0/1/1/1
5	NAG	В	1404	1	-	1/6/23/26	0/1/1/1
5	NAG	С	1406	1	-	2/6/23/26	0/1/1/1
5	NAG	А	1404	1	-	2/6/23/26	0/1/1/1
5	NAG	С	1405	1	-	2/6/23/26	0/1/1/1
5	NAG	А	1406	1	-	2/6/23/26	0/1/1/1
5	NAG	В	1402	1	-	1/6/23/26	0/1/1/1
5	NAG	В	1411	1	-	2/6/23/26	0/1/1/1

Continued from previous page...

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 43 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	А	1407	NAG	O5-C5-C6-O6
5	С	1406	NAG	C4-C5-C6-O6
5	В	1407	NAG	O5-C5-C6-O6
5	А	1404	NAG	O5-C5-C6-O6
5	В	1411	NAG	O5-C5-C6-O6

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	С	1401	NAG	1	0
5	В	1405	NAG	1	0



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Map visualisation (i)

This section contains visualisations of the EMDB entry EMD-13742. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections (i)

6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices (i)

6.2.1 Primary map



X Index: 140

Y Index: 140



Z Index: 140

The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices (i)

6.3.1 Primary map



X Index: 142

Y Index: 137

Z Index: 130

The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views (i)

6.4.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0261. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.



6.5 Mask visualisation (i)

This section was not generated. No masks/segmentation were deposited.



7 Map analysis (i)

This section contains the results of statistical analysis of the map.

7.1 Map-value distribution (i)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.



7.2 Volume estimate (i)



The volume at the recommended contour level is 1066 $\rm nm^3;$ this corresponds to an approximate mass of 963 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.



7.3 Rotationally averaged power spectrum (i)



*Reported resolution corresponds to spatial frequency of 0.208 $\mathrm{\AA^{-1}}$



8 Fourier-Shell correlation (i)

This section was not generated. No FSC curve or half-maps provided.



9 Map-model fit (i)

This section contains information regarding the fit between EMDB map EMD-13742 and PDB model 7Q0A. Per-residue inclusion information can be found in section 3 on page 8.

9.1 Map-model overlay (i)



The images above show the 3D surface view of the map at the recommended contour level 0.0261 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.



9.2 Atom inclusion (i)



At the recommended contour level, 95% of all backbone atoms, 94% of all non-hydrogen atoms, are inside the map.

