

# Full wwPDB X-ray Structure Validation Report (i)

### Jan 13, 2024 - 02:51 pm GMT

; Moroz, O.; Blagova, E.

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY \, DIFFRACTION$ 

The reported resolution of this entry is 1.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Motric	Whole archive	Similar resolution
	$(\# {\rm Entries})$	$(\# { m Entries},  { m resolution}  { m range}({ m \AA}))$
$R_{free}$	130704	4690 (1.50-1.46)
Clashscore	141614	4955 (1.50-1.46)
Ramachandran outliers	138981	4846 (1.50-1.46)
Sidechain outliers	138945	4844 (1.50-1.46)
RSRZ outliers	127900	4614 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	А	785	94%
1	В	785	94%
2	С	9	100%
3	D	2	100%
3	F	2	100%



Mol	Chain	Length	Quality of chain
	_		
4	E	10	100%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	EDO	В	925	-	-	Х	-



# 2 Entry composition (i)

There are 10 unique types of molecules in this entry. The entry contains 14338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Exo-1,4-beta-xylosidase xlnD.

Mol	Chain	Residues		A	$\operatorname{toms}$		ZeroOcc	AltConf	Trace	
1	В	769	Total 5945	C 3752	N 982	O 1195	S 16	0	8	0
1	А	770	Total 5987	C 3775	N 993	O 1203	S 16	0	11	0

• Molecule 2 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyran ose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deox y-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	С	9	Total 105	$\begin{array}{c} \mathrm{C} \\ 58 \end{array}$	N 2	O 45	0	0	0

• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	D	2	Total 28	C 16	N 2	O 10	0	0	0
3	F	2	Total 28	C 16	N 2	O 10	0	0	0

• Molecule 4 is an oligosaccharide called alpha-D-mannopyranose-(1-2)-alpha-D-mannopyran



ose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-gluco



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
4	Е	10	Total         C         N         O           116         64         2         50	0	0	0

• Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula:  $C_8H_{15}NO_6$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	$\begin{array}{cccc} \text{Total} & \text{C} & \text{N} & \text{O} \\ 14 & 8 & 1 & 5 \end{array}$	0	0
5	В	1	Total         C         N         O           14         8         1         5	0	0
5	В	1	Total         C         N         O           14         8         1         5	0	0
5	В	1	Total         C         N         O           14         8         1         5	0	0
5	В	1	Total         C         N         O           14         8         1         5	0	0
5	В	1	Total         C         N         O           14         8         1         5	0	0

Continued on next page...

WIDE

Mol	Chain	Residues	A	ton	ns		ZeroOcc	AltConf	
5	Р	1	Total	С	Ν	0	0	0	
5	D	L	14	8	1	5	0	0	
5	В	1	Total	С	Ν	Ο	0	0	
0			T	14	8	1	5	0	0
5	В	1	Total	С	Ν	Ο	0	0	
0	D	I	14	8	1	5	0	0	
5	Δ	1	Total	С	Ν	Ο	0	0	
5	11	I	14	8	1	5	0	0	
5	Δ	1	Total	С	Ν	Ο	0	0	
0	11	I	14	8	1	5	0	0	
5	Δ	1	Total	С	Ν	Ο	0	0	
0		±	14	8	1	5	0	0	
5	А	1	Total	С	Ν	Ο	0	0	
		T	14	8	1	5	Ŭ	· · · · · · · · · · · · · · · · · · ·	
5	А	1	Total	С	Ν	Ο	0	0	
		Ŧ	14	8	1	5	Ŭ	· · · · · · · · · · · · · · · · · · ·	
5	А	1	Total	С	Ν	Ο	0	0	
		±	14	8	1	5	Ŭ	0	
5	А	1	Total	С	Ν	Ο	0	0	
		*	14	8	1	5		V	
5	А	1	Total	С	Ν	Ο	0	0	
			14	8	1	5			

• Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
6	В	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 4  2  2 \end{array}$	0	0
6	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 4 & 2 & 2 \end{array}$	0	0

• Molecule 7 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf	
7	В	1	Total 4	C 2	0 2	0	0



• Molecule 8 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
8	В	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0
8	А	1	$\begin{array}{ccc} \text{Total} & \text{C} & \text{O} \\ 6 & 3 & 3 \end{array}$	0	0

• Molecule 9 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	В	3	Total K 3 3	0	0
9	А	1	Total K 1 1	0	0

• Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	В	873	Total O 873 873	0	0
10	А	948	Total O 948 948	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Exo-1,4-beta-xylosidase xlnD

 $\label{eq:mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-$ 

Chain C:

100%

#### NAG1 NAG2 BMA3 MAN5 MAN5 MAN5 MAN6 MAN8 MAN8

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:

100%



• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:

100%

#### NAG1 NAG2

 $\label{eq:mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]alpha-D-mannopyranose-(1-6)-[alpha-D-mannopyranose-(1-2)-alpha-D-mannopyranose-(1-3)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glu copyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glu copyranose$ 

Chain E:

100%

#### NAG1 NAG2 BMA3 BMA3 MAN4 MAN5 MAN5 MAN7 MAN8 MAN8 MAN9 MAN10



# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	70.47Å 90.58Å 244.37Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Bosolution (Å)	33.24 - 1.48	Depositor
Resolution (A)	33.22 - 1.48	EDS
% Data completeness	99.9 (33.24-1.48)	Depositor
(in resolution range)	100.0 (33.22 - 1.48)	EDS
$R_{merge}$	(Not available)	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.79 (at 1.48 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0238	Depositor
D D.	0.159 , $0.193$	Depositor
II, II, <i>free</i>	0.162 , $0.196$	DCC
$R_{free}$ test set	13022 reflections $(5.01\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	18.8	Xtriage
Anisotropy	0.176	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.34 , $41.8$	EDS
L-test for $twinning^2$	$ < L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	14338	wwPDB-VP
Average B, all atoms $(Å^2)$	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 11.20% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PCA, ACT, EDO, NAG, MAN, BMA, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bo	nd lengths	Bond angles		
1VIOI	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5	
1	А	0.79	1/6127~(0.0%)	0.88	1/8373~(0.0%)	
1	В	0.78	0/6086	0.90	5/8320~(0.1%)	
All	All	0.79	1/12213~(0.0%)	0.89	6/16693~(0.0%)	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
1	А	596	GLU	CD-OE2	5.79	1.32	1.25

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	В	114	ARG	NE-CZ-NH1	7.56	124.08	120.30
1	В	387	ARG	CG-CD-NE	-6.27	98.63	111.80
1	В	716	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	В	169	TYR	CB-CG-CD2	-5.39	117.77	121.00
1	В	169	TYR	CB-CG-CD1	5.25	124.15	121.00
1	А	368	ARG	NE-CZ-NH2	-5.23	117.68	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



6Q7I	
0.0011	

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	5987	0	5631	21	0
1	В	5945	0	5589	19	0
2	С	105	0	88	0	0
3	D	28	0	25	0	0
3	F	28	0	25	0	0
4	Е	116	0	97	0	0
5	А	112	0	104	0	0
5	В	126	0	117	0	0
6	А	28	0	41	8	0
6	В	16	0	24	10	0
7	В	4	0	3	1	0
8	А	6	0	8	0	0
8	В	12	0	16	3	0
9	А	1	0	0	0	0
9	В	3	0	0	0	0
10	А	948	0	0	13	0
10	В	873	0	0	7	0
All	All	14338	0	11768	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
6:A:921:EDO:H11	10:A:1522:HOH:O	1.79	0.83
1:B:208:GLY:HA2	6:B:925:EDO:H21	1.62	0.79
6:A:923:EDO:H22	10:A:1782:HOH:O	1.80	0.79
6:B:924:EDO:HO1	6:B:925:EDO:HO1	1.20	0.78
6:B:924:EDO:O1	6:B:925:EDO:O1	1.92	0.76
6:A:923:EDO:C2	10:A:1782:HOH:O	2.38	0.70
1:B:265:LYS:NZ	10:B:1001:HOH:O	2.23	0.69
1:B:716:ARG:HB3	6:B:923:EDO:H11	1.75	0.69
1:A:515:ASN:ND2	6:A:923:EDO:O2	2.27	0.67
1:A:285[A]:LYS:HD2	10:A:1756:HOH:O	1.98	0.63
1:A:233:ASN:ND2	1:A:514:GLU:OE1	2.31	0.63
1:B:734:VAL:CG1	1:B:748:LEU:HD11	2.29	0.62
1:B:145:HIS:NE2	6:B:924:EDO:H11	2.14	0.62
1:B:38:LEU:H	7:B:922:ACT:H2	1.65	0.61
1:B:307:ASP:OD2	8:B:926:GOL:H32	2.03	0.59
1:A:777[B]:LEU:C	1:A:777[B]:LEU:HD23	2.23	0.58
1:B:382:GLU:HG3	10:B:1456:HOH:O	2.05	0.57



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
8:B:927:GOL:H2	10:B:1106:HOH:O	2.04	0.56
6:A:924:EDO:H12	10:A:1374:HOH:O	2.05	0.55
6:B:921:EDO:H11	10:B:1413:HOH:O	2.06	0.55
6:A:924:EDO:C2	10:A:1374:HOH:O	2.56	0.53
6:B:923:EDO:H22	10:B:1394:HOH:O	2.07	0.53
6:A:921:EDO:C1	10:A:1522:HOH:O	2.50	0.51
1:A:768:GLU:OE2	10:A:1002:HOH:O	2.19	0.51
1:A:468:HIS:CG	1:A:492:GLN:HE22	2.28	0.50
1:B:149:THR:HG22	6:B:925:EDO:H22	1.95	0.48
1:A:432:ILE:HA	1:A:470:ALA:O	2.13	0.48
1:B:417:ASP:O	1:B:418:GLU:CB	2.61	0.48
1:A:528:LYS:NZ	10:A:1006:HOH:O	2.40	0.48
6:B:925:EDO:H11	10:B:1231:HOH:O	2.15	0.46
1:A:468:HIS:CG	1:A:492:GLN:NE2	2.84	0.46
6:B:921:EDO:H12	10:B:1029:HOH:O	2.16	0.46
1:A:313:ASN:HA	1:A:316:ASN:O	2.16	0.46
1:B:702:GLY:HA2	1:B:749:TYR:CD1	2.52	0.45
1:A:128:SER:HB2	10:A:1203:HOH:O	2.17	0.45
1:B:432:ILE:HA	1:B:470:ALA:O	2.16	0.44
1:B:270:MET:HA	1:B:305:SER:O	2.18	0.44
1:B:446:TYR:OH	8:B:926:GOL:H2	2.17	0.44
6:A:924:EDO:H22	10:A:1374:HOH:O	2.17	0.44
1:B:313:ASN:HA	1:B:316:ASN:O	2.18	0.43
1:B:678[A]:THR:HA	1:B:728:LEU:O	2.17	0.43
1:A:316:ASN:HA	1:A:317:PRO:HA	1.79	0.43
1:A:724[A]:ASP:OD2	10:A:1003:HOH:O	2.21	0.43
1:A:504:ASN:HA	1:A:507:GLU:O	2.19	0.43
1:A:248:GLU:HA	1:A:251:GLU:HG2	2.01	0.42
1:A:700:THR:OG1	10:A:1001:HOH:O	1.72	0.41
1:A:539:LEU:HD21	1:A:578:ILE:HD11	2.02	0.41
1:B:370:TYR:O	1:B:374:VAL:HG23	2.19	0.41
1:A:270:MET:HA	1:A:305:SER:O	2.19	0.41
1:B:268:SER:HA	1:B:303:TYR:O	2.21	0.41
1:A:542:GLY:O	1:A:564:GLY:HA2	2.21	0.41
1:B:504:ASN:HA	1:B:507:GLU:O	2.21	0.40
1:A:178:HIS:CG	1:A:179:PRO:HD2	2.57	0.40
1:A:777[B]:LEU:C	1:A:777[B]:LEU:CD2	2.89	0.40

There are no symmetry-related clashes.



### 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured Allowed		Outliers	Percentiles		
1	А	780/785~(99%)	753 (96%)	26 (3%)	1 (0%)	51	25	
1	В	775/785~(99%)	750 (97%)	24 (3%)	1 (0%)	51	25	
All	All	1555/1570~(99%)	1503 (97%)	50 (3%)	2(0%)	51	25	

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type	
1	А	231	SER	
1	В	231	SER	

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	641/647~(99%)	639~(100%)	2(0%)	92 84
1	В	636/647~(98%)	634 (100%)	2(0%)	92 84
All	All	1277/1294~(99%)	1273 (100%)	4 (0%)	92 84

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	В	176	PHE
1	В	233	ASN
1	А	176	PHE
1	А	233	ASN



Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such side chains are listed below:

Mol	Chain	Res	Type
1	А	515	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Tyj	Turne	Chain	Res	Link	Bond lengths			Bond angles		
	туре	Ullaill			Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z >2
1	PCA	А	19	1	7,8,9	0.61	0	9,10,12	1.01	1 (11%)
1	PCA	В	19	1	7,8,9	0.52	0	9,10,12	0.77	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PCA	А	19	1	-	0/0/11/13	0/1/1/1
1	PCA	В	19	1	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
1	А	19	PCA	O-C-CA	-2.10	119.29	124.78

There are no chirality outliers.



There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.5 Carbohydrates (i)

23 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Bos	Link	Bo	ond leng	ths	B	ond ang	les
WIOI	Type	Ullalli	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	С	1	2,1	14,14,15	0.91	1 (7%)	17,19,21	1.68	3 (17%)
2	NAG	С	2	2	14,14,15	0.56	0	17,19,21	1.24	1 (5%)
2	BMA	С	3	2	11,11,12	1.56	1 (9%)	15,15,17	1.31	2 (13%)
2	MAN	С	4	2	11,11,12	0.67	0	15,15,17	1.24	2 (13%)
2	MAN	С	5	2	11,11,12	0.88	0	15,15,17	1.52	3 (20%)
2	MAN	С	6	2	11,11,12	0.65	0	15,15,17	1.39	2 (13%)
2	MAN	С	7	2	11,11,12	0.71	0	15,15,17	1.36	1 (6%)
2	MAN	С	8	2	11,11,12	1.09	1 (9%)	15,15,17	0.96	0
2	MAN	С	9	2	11,11,12	0.90	1 (9%)	15,15,17	1.85	4 (26%)
3	NAG	D	1	3,1	14,14,15	0.69	0	17,19,21	1.25	1 (5%)
3	NAG	D	2	3	14,14,15	0.76	0	17,19,21	1.26	3 (17%)
4	NAG	Е	1	4,1	14,14,15	1.47	2 (14%)	17,19,21	2.28	4 (23%)
4	MAN	Е	10	4	11,11,12	1.01	0	15,15,17	1.48	4 (26%)
4	NAG	Е	2	4	14,14,15	0.92	1 (7%)	17,19,21	1.58	4 (23%)
4	BMA	Е	3	4	11,11,12	0.65	0	15,15,17	1.71	2 (13%)
4	MAN	Е	4	4	11,11,12	0.80	0	15,15,17	1.53	4 (26%)
4	MAN	Е	5	4	11,11,12	0.88	1 (9%)	15,15,17	3.07	6 (40%)
4	MAN	E	6	4	11,11,12	1.32	2 (18%)	15,15,17	2.75	5 (33%)
4	MAN	Е	7	4	11,11,12	0.48	0	15,15,17	1.00	1 (6%)
4	MAN	Е	8	4	11,11,12	0.48	0	15,15,17	1.06	1 (6%)
4	MAN	Е	9	4	11,11,12	0.63	0	15,15,17	1.67	4 (26%)



Mol Type	Turne	Chain	Dec	Tinle	Bo	ond leng	$_{\rm ths}$	Bond angles		
	Ullalli	nes		Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2	
3	NAG	F	1	3,1	14,14,15	0.76	0	$17,\!19,\!21$	1.39	3 (17%)
3	NAG	F	2	3	14,14,15	0.91	0	17,19,21	1.78	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	С	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	С	2	2	_	0/6/23/26	0/1/1/1
2	BMA	С	3	2	_	0/2/19/22	0/1/1/1
2	MAN	С	4	2	-	0/2/19/22	0/1/1/1
2	MAN	С	5	2	-	2/2/19/22	0/1/1/1
2	MAN	С	6	2	_	0/2/19/22	0/1/1/1
2	MAN	С	7	2	-	0/2/19/22	0/1/1/1
2	MAN	С	8	2	-	0/2/19/22	0/1/1/1
2	MAN	С	9	2	-	0/2/19/22	0/1/1/1
3	NAG	D	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	2/6/23/26	0/1/1/1
4	NAG	Е	1	4,1	-	2/6/23/26	0/1/1/1
4	MAN	Е	10	4	-	2/2/19/22	0/1/1/1
4	NAG	Е	2	4	-	0/6/23/26	0/1/1/1
4	BMA	Е	3	4	-	0/2/19/22	0/1/1/1
4	MAN	Е	4	4	-	0/2/19/22	0/1/1/1
4	MAN	Е	5	4	-	0/2/19/22	0/1/1/1
4	MAN	Е	6	4	-	1/2/19/22	0/1/1/1
4	MAN	Е	7	4	-	0/2/19/22	0/1/1/1
4	MAN	Е	8	4	-	0/2/19/22	0/1/1/1
4	MAN	E	9	4	-	2/2/19/22	0/1/1/1
3	NAG	F	1	3,1	_	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	0/6/23/26	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
2	С	3	BMA	O5-C1	-4.11	1.37	1.43
4	Ε	1	NAG	C1-C2	3.71	1.57	1.52
4	Ε	6	MAN	O5-C1	-2.94	1.39	1.43
4	Ε	1	NAG	C8-C7	2.73	1.56	1.50



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\operatorname{Ideal}(\operatorname{\AA})$
4	Е	2	NAG	C1-C2	2.30	1.55	1.52
2	С	9	MAN	O2-C2	2.28	1.48	1.43
2	С	1	NAG	C1-C2	2.24	1.55	1.52
2	С	8	MAN	O5-C1	-2.07	1.40	1.43
4	Е	5	MAN	C2-C3	-2.04	1.49	1.52
4	Е	6	MAN	C2-C3	2.01	1.55	1.52

All (65) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	Е	5	MAN	O2-C2-C3	-8.48	93.16	110.14
4	Е	1	NAG	O5-C1-C2	-7.50	99.45	111.29
4	Е	6	MAN	C1-O5-C5	6.33	120.77	112.19
4	Е	6	MAN	O5-C5-C6	-6.08	97.68	107.20
2	С	1	NAG	O5-C1-C2	-4.95	103.48	111.29
4	Е	5	MAN	O2-C2-C1	4.56	118.47	109.15
4	Е	9	MAN	O5-C5-C6	4.21	113.80	107.20
4	Е	3	BMA	C1-O5-C5	-3.95	106.84	112.19
2	С	7	MAN	C6-C5-C4	-3.87	103.94	113.00
4	Е	3	BMA	C2-C3-C4	-3.86	104.22	110.89
3	F	2	NAG	C1-O5-C5	3.78	117.32	112.19
4	Е	5	MAN	O4-C4-C3	-3.73	101.72	110.35
2	С	5	MAN	C6-C5-C4	-3.68	104.39	113.00
4	Е	5	MAN	C6-C5-C4	-3.67	104.41	113.00
2	С	9	MAN	O2-C2-C1	3.57	116.46	109.15
2	С	6	MAN	O5-C1-C2	-3.51	105.35	110.77
2	С	9	MAN	C1-C2-C3	-3.24	105.68	109.67
3	F	2	NAG	O3-C3-C4	-3.11	103.15	110.35
2	С	2	NAG	C2-N2-C7	3.11	127.33	122.90
4	Е	2	NAG	O5-C5-C6	3.09	112.05	107.20
4	Е	1	NAG	O7-C7-C8	-2.98	116.52	122.06
2	С	1	NAG	C4-C3-C2	-2.98	106.66	111.02
4	Е	4	MAN	O2-C2-C3	-2.97	104.18	110.14
4	Е	2	NAG	O6-C6-C5	-2.92	101.28	111.29
4	Е	6	MAN	O3-C3-C2	2.85	115.44	109.99
4	Е	10	MAN	O3-C3-C2	-2.84	104.55	109.99
3	F	2	NAG	O5-C1-C2	-2.83	106.82	111.29
3	F	1	NAG	C3-C4-C5	-2.80	105.25	110.24
3	F	2	NAG	O4-C4-C5	2.72	116.06	109.30
2	С	4	MAN	O5-C5-C6	-2.65	103.04	107.20
4	Е	6	MAN	$C\overline{3}-C4-C5$	-2.63	105.55	110.24
4	Е	2	NAG	O4-C4-C5	-2.59	102.87	109.30



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
4	Е	10	MAN	O4-C4-C5	-2.58	102.89	109.30
2	С	3	BMA	C1-C2-C3	-2.56	106.52	109.67
2	С	9	MAN	O2-C2-C3	2.55	115.25	110.14
4	Е	7	MAN	C1-O5-C5	-2.54	108.75	112.19
4	Е	2	NAG	C4-C3-C2	2.53	114.72	111.02
3	F	1	NAG	O5-C1-C2	-2.46	107.40	111.29
2	С	6	MAN	C1-C2-C3	-2.44	106.67	109.67
3	D	1	NAG	O6-C6-C5	-2.44	102.93	111.29
3	D	2	NAG	C2-N2-C7	2.43	126.37	122.90
2	С	4	MAN	O3-C3-C2	2.41	114.61	109.99
2	С	9	MAN	C1-O5-C5	2.40	115.45	112.19
4	Ε	4	MAN	O3-C3-C4	-2.40	104.81	110.35
2	С	3	BMA	O5-C5-C6	2.40	110.96	107.20
4	Ε	4	MAN	O6-C6-C5	-2.39	103.11	111.29
3	D	2	NAG	O7-C7-N2	2.35	126.28	121.95
3	F	2	NAG	C6-C5-C4	-2.35	107.50	113.00
4	Ε	4	MAN	O2-C2-C1	2.34	113.94	109.15
3	F	1	NAG	C2-N2-C7	2.28	126.15	122.90
4	Ε	9	MAN	O3-C3-C4	-2.23	105.20	110.35
4	Ε	1	NAG	O7-C7-N2	2.22	126.03	121.95
4	Ε	1	NAG	O5-C5-C4	-2.21	105.45	110.83
2	С	1	NAG	O5-C5-C4	-2.18	105.53	110.83
4	Ε	9	MAN	O3-C3-C2	2.17	114.14	109.99
4	Ε	6	MAN	C1-C2-C3	-2.16	107.02	109.67
4	Ε	10	MAN	O3-C3-C4	2.14	115.30	110.35
4	Ε	5	MAN	O3-C3-C4	-2.14	105.41	110.35
2	С	5	MAN	O3-C3-C2	-2.13	105.91	109.99
3	D	2	NAG	C8-C7-N2	-2.12	112.50	116.10
4	Ε	10	MAN	O2-C2-C1	2.12	113.49	109.15
4	Ε	8	MAN	O5-C5-C6	-2.11	103.90	107.20
2	С	5	MAN	$O4-C4-C\overline{3}$	2.09	115.18	110.35
4	E	9	MAN	O4-C4-C3	-2.03	105.65	110.35
4	Ε	5	MAN	O5-C5-C6	2.03	110.38	107.20

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	2	NAG	C4-C5-C6-O6
3	D	2	NAG	O5-C5-C6-O6
4	Е	9	MAN	O5-C5-C6-O6
4	Е	6	MAN	O5-C5-C6-O6



Mol	Chain	Res	Type	Atoms
2	С	5	MAN	O5-C5-C6-O6
4	Е	10	MAN	C4-C5-C6-O6
4	Е	10	MAN	O5-C5-C6-O6
4	Е	9	MAN	C4-C5-C6-O6
2	С	1	NAG	C3-C2-N2-C7
4	Е	1	NAG	C3-C2-N2-C7
4	Е	1	NAG	C1-C2-N2-C7
2	С	5	MAN	C4-C5-C6-O6
2	С	1	NAG	C1-C2-N2-C7

Continued from previous page...

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.















## 5.6 Ligand geometry (i)

Of 36 ligands modelled in this entry, 4 are monoatomic - leaving 32 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mal	Turne	Chain	Dec	Timle	Bond lengths		B	Bond angles		
	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
6	EDO	А	923	-	3,3,3	0.52	0	2,2,2	0.13	0
6	EDO	А	926	6	3,3,3	0.26	0	2,2,2	0.47	0
5	NAG	А	901	1	14,14,15	0.88	0	17,19,21	1.52	4 (23%)
6	EDO	В	925	-	3,3,3	0.47	0	2,2,2	1.21	0
5	NAG	А	917	1	14,14,15	0.72	0	17,19,21	1.54	4 (23%)
7	ACT	В	922	-	3,3,3	0.77	0	3,3,3	1.33	0
5	NAG	В	901	1	14,14,15	1.19	2 (14%)	17,19,21	2.18	3 (17%)
8	GOL	В	926	-	5,5,5	0.18	0	$5,\!5,\!5$	1.10	0
5	NAG	А	920	1	14,14,15	0.97	1 (7%)	17,19,21	1.84	6 (35%)
8	GOL	В	927	-	5,5,5	0.11	0	$5,\!5,\!5$	0.36	0
5	NAG	В	917	1	14,14,15	0.61	0	17,19,21	2.29	8 (47%)
5	NAG	А	919	1	14,14,15	0.51	0	17,19,21	0.81	0
6	EDO	В	921	-	3,3,3	0.32	0	2,2,2	0.36	0
8	GOL	А	928	-	5,5,5	0.08	0	$5,\!5,\!5$	0.49	0
6	EDO	В	924	-	3,3,3	0.33	0	2,2,2	0.19	0
6	EDO	A	925	-	3,3,3	0.32	0	2,2,2	0.38	0
5	NAG	В	918	1	14,14,15	0.52	0	17,19,21	1.24	3 (17%)
5	NAG	В	913	1	14,14,15	0.68	0	17,19,21	0.86	0
5	NAG	В	914	1	14,14,15	0.94	1 (7%)	17,19,21	1.66	4 (23%)
5	NAG	В	920	1	14,14,15	1.34	2 (14%)	17,19,21	1.85	2 (11%)
6	EDO	А	924	-	3,3,3	0.57	0	2,2,2	0.62	0
5	NAG	В	912	1	14,14,15	0.75	0	17,19,21	0.95	0
5	NAG	В	919	1	14,14,15	0.77	1 (7%)	17,19,21	1.10	2 (11%)
5	NAG	А	914	1	14,14,15	1.14	1 (7%)	17,19,21	2.14	4 (23%)
5	NAG	А	918	1	14,14,15	1.01	0	17,19,21	1.61	3 (17%)
6	EDO	А	921	-	3,3,3	0.15	0	2,2,2	0.21	0
5	NAG	В	911	1	14,14,15	0.60	0	17,19,21	0.97	0
5	NAG	А	915	1	14,14,15	0.51	0	17,19,21	1.22	1 (5%)
5	NAG	А	916	1	14,14,15	0.87	1 (7%)	17,19,21	1.08	1 (5%)
6	EDO	В	923	-	3,3,3	0.66	0	2,2,2	0.63	0
6	EDO	А	922	-	3,3,3	0.11	0	2,2,2	0.70	0
6	EDO	A	927	6	3,3,3	0.27	0	2,2,2	0.58	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	А	923	-	-	1/1/1/1	-

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	А	926	6	-	1/1/1/1	-
5	NAG	А	901	1	-	0/6/23/26	0/1/1/1
6	EDO	В	925	-	-	0/1/1/1	_
5	NAG	А	917	1	-	0/6/23/26	0/1/1/1
5	NAG	В	901	1	-	1/6/23/26	0/1/1/1
8	GOL	В	926	-	-	3/4/4/4	-
5	NAG	А	920	1	-	0/6/23/26	0/1/1/1
8	GOL	В	927	-	-	0/4/4/4	-
5	NAG	В	917	1	-	2/6/23/26	0/1/1/1
5	NAG	А	919	1	-	0/6/23/26	0/1/1/1
6	EDO	В	921	-	-	1/1/1/1	-
8	GOL	А	928	-	-	0/4/4/4	-
6	EDO	В	924	-	-	0/1/1/1	-
6	EDO	А	925	-	-	1/1/1/1	-
5	NAG	В	918	1	-	0/6/23/26	0/1/1/1
5	NAG	В	913	1	-	0/6/23/26	0/1/1/1
5	NAG	В	914	1	-	0/6/23/26	0/1/1/1
5	NAG	В	920	1	-	0/6/23/26	0/1/1/1
6	EDO	А	924	-	-	0/1/1/1	-
5	NAG	В	912	1	-	0/6/23/26	0/1/1/1
5	NAG	В	919	1	-	0/6/23/26	0/1/1/1
5	NAG	А	914	1	-	0/6/23/26	0/1/1/1
5	NAG	А	918	1	-	0/6/23/26	0/1/1/1
6	EDO	А	921	-	-	0/1/1/1	-
5	NAG	В	911	1	-	0/6/23/26	0/1/1/1
5	NAG	А	915	1	-	0/6/23/26	0/1/1/1
5	NAG	A	916	1	-	0/6/23/26	0/1/1/1
6	EDO	В	923	-	-	1/1/1/1	-
6	EDO	А	922	-	-	1/1/1/1	-
6	EDO	А	927	6	-	0/1/1/1	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
5	В	920	NAG	C1-C2	3.55	1.57	1.52
5	В	901	NAG	C8-C7	2.86	1.56	1.50
5	В	920	NAG	O5-C1	2.51	1.47	1.43
5	А	914	NAG	O7-C7	2.49	1.28	1.23
5	В	914	NAG	O5-C5	2.41	1.48	1.43
5	А	920	NAG	O6-C6	2.36	1.52	1.42
5	В	901	NAG	O5-C5	2.30	1.48	1.43
5	В	919	NAG	C1-C2	2.28	1.55	1.52



Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	$\mathrm{Ideal}(\mathrm{\AA})$
5	А	916	NAG	C8-C7	2.15	1.55	1.50

All (45) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	В	901	NAG	C2-N2-C7	6.88	132.71	122.90
5	А	914	NAG	C8-C7-N2	-6.30	105.44	116.10
5	В	917	NAG	C1-O5-C5	4.88	118.81	112.19
5	В	920	NAG	C1-O5-C5	4.85	118.77	112.19
5	А	920	NAG	C1-O5-C5	4.55	118.36	112.19
5	В	920	NAG	O5-C1-C2	-4.54	104.11	111.29
5	А	918	NAG	O5-C5-C4	-3.83	101.50	110.83
5	В	917	NAG	C1-C2-N2	-3.79	104.02	110.49
5	В	901	NAG	C8-C7-N2	-3.60	110.00	116.10
5	В	914	NAG	C1-O5-C5	3.42	116.83	112.19
5	В	917	NAG	O7-C7-C8	3.25	128.09	122.06
5	В	917	NAG	C8-C7-N2	-3.13	110.79	116.10
5	В	901	NAG	O5-C1-C2	-2.94	106.64	111.29
5	А	914	NAG	O7-C7-N2	2.91	127.30	121.95
5	В	914	NAG	O5-C5-C4	-2.89	103.80	110.83
5	А	901	NAG	O5-C5-C6	2.86	111.68	107.20
5	А	914	NAG	O7-C7-C8	2.79	127.24	122.06
5	В	918	NAG	O5-C5-C4	-2.77	104.08	110.83
5	А	918	NAG	C1-O5-C5	2.74	115.91	112.19
5	А	917	NAG	C8-C7-N2	-2.71	111.51	116.10
5	А	915	NAG	O5-C5-C4	-2.70	104.26	110.83
5	В	914	NAG	O4-C4-C3	-2.64	104.24	110.35
5	В	917	NAG	C2-N2-C7	-2.52	119.32	122.90
5	А	920	NAG	O5-C1-C2	-2.51	107.33	111.29
5	В	914	NAG	O3-C3-C2	-2.43	104.43	109.47
5	А	917	NAG	C1-O5-C5	2.42	115.47	112.19
5	А	916	NAG	O4-C4-C3	-2.41	104.78	110.35
5	В	917	NAG	O5-C5-C6	-2.40	103.44	107.20
5	В	918	NAG	O6-C6-C5	-2.39	103.09	111.29
5	А	901	NAG	O4-C4-C5	2.38	115.21	109.30
5	А	920	NAG	O3-C3-C4	-2.35	104.92	110.35
5	А	920	NAG	O5-C5-C6	2.29	110.79	107.20
5	В	918	NAG	C1-O5-C5	-2.27	109.11	112.19
5	В	919	NAG	O5-C5-C6	2.24	110.72	107.20
5	А	917	NAG	O7-C7-C8	2.22	126.19	122.06
5	А	901	NAG	O3-C3-C4	-2.22	105.22	110.35
5	А	920	NAG	O6-C6-C5	2.20	118.84	111.29



6Q/1
------

Mol	Chain	Res	Type	Atoms	Ζ	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
5	А	917	NAG	O5-C5-C6	2.19	110.64	107.20
5	А	901	NAG	C2-N2-C7	-2.19	119.79	122.90
5	В	917	NAG	O6-C6-C5	-2.18	103.80	111.29
5	В	917	NAG	O5-C1-C2	-2.17	107.85	111.29
5	А	914	NAG	C2-N2-C7	2.08	125.87	122.90
5	В	919	NAG	O4-C4-C3	-2.05	105.62	110.35
5	А	918	NAG	O5-C5-C6	2.02	110.38	107.20
5	А	920	NAG	C1-C2-N2	2.02	113.94	110.49

Continued from previous page...

There are no chirality outliers.

Mol	Chain	$\operatorname{Res}$	Type	Atoms
8	В	926	GOL	C1-C2-C3-O3
5	В	917	NAG	C4-C5-C6-O6
8	В	926	GOL	O1-C1-C2-O2
8	В	926	GOL	O2-C2-C3-O3
5	В	917	NAG	O5-C5-C6-O6
6	В	921	EDO	O1-C1-C2-O2
6	В	923	EDO	O1-C1-C2-O2
5	В	901	NAG	C1-C2-N2-C7
6	А	925	EDO	O1-C1-C2-O2
6	А	922	EDO	O1-C1-C2-O2
6	А	926	EDO	O1-C1-C2-O2
6	А	923	EDO	O1-C1-C2-O2

All (12) torsion outliers are listed below:

There are no ring outliers.

10 monomers are involved in 22 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	А	923	EDO	3	0
6	В	925	EDO	5	0
7	В	922	ACT	1	0
8	В	926	GOL	2	0
8	В	927	GOL	1	0
6	В	921	EDO	2	0
6	В	924	EDO	3	0
6	А	924	EDO	3	0
6	А	921	EDO	2	0
6	В	923	EDO	2	0



# 5.7 Other polymers (i)

There are no such residues in this entry.

# 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	769/785~(97%)	-0.38	4 (0%) 91 93	14, 19, 31, 45	19 (2%)
1	В	768/785~(97%)	-0.28	5 (0%) 87 90	14, 21, 36, 48	14 (1%)
All	All	1537/1570~(97%)	-0.33	9 (0%) 89 91	14, 20, 33, 48	33 (2%)

All (9) RSRZ outliers are listed below:

Mol	Chain	$\mathbf{Res}$	Type	RSRZ
1	А	788	VAL	4.8
1	А	787	PHE	3.2
1	А	57	LEU	3.1
1	В	425	ASP	2.7
1	А	24	TYR	2.2
1	В	533	GLY	2.2
1	В	493	ALA	2.1
1	В	423	SER	2.1
1	В	787	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
1	PCA	А	19	8/9	0.95	0.07	17,19,20,20	0
1	PCA	В	19	8/9	0.98	0.04	17,18,20,20	0



### 6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	MAN	Е	10	11/12	0.62	0.19	$58,\!63,\!68,\!70$	0
3	NAG	F	2	14/15	0.78	0.24	47,63,73,74	0
3	NAG	D	2	14/15	0.84	0.28	40,45,55,66	0
4	MAN	Е	6	11/12	0.85	0.20	42,47,51,64	0
2	MAN	С	6	11/12	0.85	0.18	$39,\!47,\!51,\!55$	0
4	MAN	Е	5	11/12	0.86	0.22	34,41,46,50	0
2	MAN	С	9	11/12	0.89	0.11	29,39,45,49	0
2	MAN	С	5	11/12	0.89	0.15	34,40,42,46	0
4	MAN	Е	9	11/12	0.90	0.12	37,45,56,57	0
3	NAG	F	1	14/15	0.91	0.10	25,29,41,45	0
3	NAG	D	1	14/15	0.91	0.12	25,27,37,41	0
2	MAN	С	7	11/12	0.93	0.08	27,31,40,42	0
4	MAN	Е	8	11/12	0.94	0.07	19,23,28,31	0
2	MAN	С	8	11/12	0.94	0.11	21,25,30,33	0
4	BMA	Е	3	11/12	0.94	0.09	22,25,30,30	0
4	NAG	Е	1	14/15	0.95	0.07	17,22,29,29	0
2	MAN	С	4	11/12	0.95	0.10	26,28,29,31	0
4	MAN	Е	7	11/12	0.95	0.10	$26,\!29,\!33,\!35$	0
4	NAG	Е	2	14/15	0.96	0.07	22,23,29,31	0
2	BMA	С	3	11/12	0.96	0.09	23,26,30,31	0
4	MAN	Е	4	11/12	0.96	0.09	24,26,28,32	0
2	NAG	С	2	14/15	0.97	0.06	22,23,26,32	0
2	NAG	С	1	14/15	0.97	0.04	18,21,26,29	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





![](_page_31_Picture_4.jpeg)

![](_page_32_Figure_3.jpeg)

![](_page_32_Picture_4.jpeg)

![](_page_33_Figure_3.jpeg)

![](_page_33_Picture_4.jpeg)

![](_page_34_Figure_3.jpeg)

### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathrm{\AA}^2)$	Q<0.9
6	EDO	В	921	4/4	0.75	0.14	$45,\!57,\!58,\!59$	0
5	NAG	В	920	14/15	0.76	0.24	$26,\!41,\!44,\!49$	0
6	EDO	А	925	4/4	0.82	0.15	34,47,49,56	0
6	EDO	В	923	4/4	0.84	0.13	33,37,39,41	0
8	GOL	В	927	6/6	0.84	0.13	32,42,50,53	0

![](_page_34_Picture_8.jpeg)

6Q7I
------

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(A^2)$	Q<0.9
6	EDO	А	926	4/4	0.85	0.14	37,39,39,42	4
5	NAG	В	917	14/15	0.86	0.13	32,39,48,53	0
6	EDO	А	927	4/4	0.86	0.12	21,23,26,26	4
5	NAG	В	919	14/15	0.86	0.14	36,39,48,50	0
6	EDO	А	924	4/4	0.87	0.15	36,37,41,45	0
6	EDO	В	925	4/4	0.88	0.21	30,31,34,38	0
5	NAG	В	914	14/15	0.88	0.11	22,28,30,40	0
5	NAG	В	918	14/15	0.89	0.14	34,41,52,53	0
6	EDO	А	922	4/4	0.90	0.16	32,36,37,52	0
5	NAG	А	919	14/15	0.90	0.13	40,48,52,52	0
5	NAG	А	916	14/15	0.90	0.11	30,34,42,43	0
6	EDO	В	924	4/4	0.91	0.24	30,38,39,41	0
5	NAG	А	918	14/15	0.91	0.09	21,28,31,37	0
5	NAG	В	901	14/15	0.92	0.10	21,24,38,40	0
5	NAG	А	920	14/15	0.92	0.10	27,31,38,39	0
8	GOL	В	926	6/6	0.92	0.10	20,22,23,29	0
5	NAG	A	915	14/15	0.92	0.13	$29,\!36,\!45,\!47$	0
5	NAG	В	913	14/15	0.93	0.12	$29,\!39,\!48,\!52$	0
6	EDO	А	923	4/4	0.93	0.10	26,28,31,32	0
7	ACT	В	922	4/4	0.94	0.11	16,20,22,29	0
5	NAG	А	914	14/15	0.94	0.07	20,26,34,37	0
5	NAG	В	911	14/15	0.94	0.10	$25,\!30,\!34,\!36$	0
8	GOL	A	928	6/6	0.94	0.07	19,22,24,24	0
5	NAG	В	912	14/15	0.96	0.09	20,22,27,27	0
5	NAG	А	901	14/15	0.96	0.06	20,22,25,28	0
9	K	В	929	1/1	0.96	0.06	39,39,39,39	0
6	EDO	A	921	4/4	0.97	0.05	27,30,30,31	0
5	NAG	A	917	14/15	0.97	0.07	22,26,28,30	0
9	К	В	930	1/1	0.97	0.13	37,37,37,37	0
9	К	В	928	1/1	0.99	0.03	32,32,32,32	0
9	K	A	929	1/1	0.99	0.04	28,28,28,28	0

# 6.5 Other polymers (i)

There are no such residues in this entry.

![](_page_35_Picture_7.jpeg)