



# wwPDB X-ray Structure Validation Summary Report

Aug 31, 2020 – 07:16 AM BST


PDB ID : 6Q8O  
Title : Respiratory complex I from *Thermus thermophilus* with bound Piericidin A  
Authors : Gutierrez-Fernandez, J.; Minhas, G.S.; Sazanov, L.A.  
Deposited on : 2018-12-15  
Resolution : 3.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : **FAILED**  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1353 (3.70-3.50)
Ramachandran outliers	138981	1307 (3.70-3.50)
Sidechain outliers	138945	1307 (3.70-3.50)

























The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Note EDS failed to run properly.

Mol	Chain	Length	Quality of chain
1	1	438	53% 45% .
1	B	438	51% 48% .
2	2	181	60% 38% ..
2	C	181	61% 35% ..
3	3	783	57% 39% ..
3	D	783	56% 39% ..
4	4	409	48% 45% . 6%
4	E	409	41% 52% . 6%

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	5	207	 61% 32% 5%
5	F	207	 47% 45% 5%
6	6	181	 43% 46% 8%
6	G	181	 41% 48% 8%
7	9	182	 65% 34% ..
7	O	182	 53% 46% .
8	7	129	 59% 39% ..
8	I	129	 63% 35% ..
9	W	131	 71% 26% .
9	X	131	 66% 30% ..
10	A	119	 55% 43% ..
10	P	119	 52% 45% ..
11	J	176	 51% 39% . 9%
11	R	176	 60% 29% . 9%
12	K	95	 64% 34% .
12	S	95	 65% 34% .
13	L	606	 64% 35%
13	T	606	 62% 36% .
14	M	469	 59% 39% .
14	U	469	 58% 41%
15	N	427	 66% 34%
15	V	427	 63% 36% .
16	H	365	 52% 42% . .
16	Q	365	 51% 43% . .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
17	SF4	9	201	-	-	X	-
17	SF4	9	202	-	-	X	-
17	SF4	D	801	-	-	X	-
17	SF4	D	803	-	-	X	-
17	SF4	G	201	-	-	X	-
17	SF4	O	202	-	-	X	-

## 2 Entry composition [i](#)

There are 20 unique types of molecules in this entry. The entry contains 74146 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH-quinone oxidoreductase subunit 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	1	437	Total 3417	C 2180	N 595	O 624	S 18	0	0	0
1	B	437	Total 3417	C 2180	N 595	O 624	S 18	0	0	0

- Molecule 2 is a protein called NADH-quinone oxidoreductase subunit 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	2	178	Total 1406	C 895	N 238	O 265	S 8	0	0	0
2	C	178	Total 1406	C 895	N 238	O 265	S 8	0	0	0

- Molecule 3 is a protein called NADH-quinone oxidoreductase subunit 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	3	756	Total 5895	C 3754	N 1057	O 1053	S 31	0	0	0
3	D	756	Total 5895	C 3754	N 1057	O 1053	S 31	0	0	0

- Molecule 4 is a protein called NADH-quinone oxidoreductase subunit 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	4	384	Total 3067	C 1975	N 522	O 559	S 11	0	0	0
4	E	384	Total 3067	C 1975	N 522	O 559	S 11	0	0	0

- Molecule 5 is a protein called NADH-quinone oxidoreductase subunit 5.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	5	196	Total	C	N	O	S	0	0	0
			1607	1043	273	288	3			
5	F	196	Total	C	N	O	S	0	0	0
			1607	1043	273	288	3			

- Molecule 6 is a protein called NADH-quinone oxidoreductase subunit 6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	6	166	Total	C	N	O	S	0	0	0
			1289	815	235	226	13			
6	G	166	Total	C	N	O	S	0	0	0
			1289	815	235	226	13			

- Molecule 7 is a protein called NADH-quinone oxidoreductase subunit 9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	9	180	Total	C	N	O	S	0	0	0
			1388	890	232	255	11			
7	O	180	Total	C	N	O	S	0	0	0
			1388	890	232	255	11			

- Molecule 8 is a protein called NADH-quinone oxidoreductase subunit 15.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	7	127	Total	C	N	O	S	0	0	0
			1031	664	183	181	3			
8	I	127	Total	C	N	O	S	0	0	0
			1031	664	183	181	3			

- Molecule 9 is a protein called NADH-quinone oxidoreductase subunit 16.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
9	W	127	Total	C	N	O	S	0	0	0
			967	623	165	175	4			
9	X	127	Total	C	N	O	S	0	0	0
			967	623	165	175	4			

- Molecule 10 is a protein called NADH-quinone oxidoreductase subunit 7.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
10	A	117	Total	C	N	O	S	0	0	0
			910	624	138	144	4			

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
10	P	117	910	624	138	144	4	0	0	0

- Molecule 11 is a protein called NADH-quinone oxidoreductase subunit 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
11	J	160	1183	806	183	191	3	0	0	0
11	R	160	1183	806	183	191	3	0	0	0

- Molecule 12 is a protein called NADH-quinone oxidoreductase subunit 11.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
12	K	95	703	456	118	126	3	0	0	0
12	S	95	703	456	118	126	3	0	0	0

- Molecule 13 is a protein called NADH-quinone oxidoreductase subunit 12.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
13	L	605	4604	3089	740	756	19	0	0	0
13	T	605	4604	3089	740	756	19	0	0	0

- Molecule 14 is a protein called NADH-quinone oxidoreductase subunit 13.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
14	M	467	3489	2363	546	572	8	0	0	0
14	U	467	3489	2363	546	572	8	0	0	0


- Molecule 15 is a protein called NADH-quinone oxidoreductase subunit 14.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
15	N	427	3154	2125	505	518	6	0	0	0
15	V	427	3154	2125	505	518	6	0	0	0

- Molecule 16 is a protein called NADH-quinone oxidoreductase subunit 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
16	H	353	Total 2838	C 1943	N 431	O 457	S 7	0	0	0
16	Q	353	Total 2838	C 1943	N 431	O 457	S 7	0	0	0

- Molecule 17 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	Fe	S		
17	1	1	Total 8	Fe 4	S 4	0	0
17	3	1	Total 8	Fe 4	S 4	0	0
17	3	1	Total 8	Fe 4	S 4	0	0
17	3	1	Total 8	Fe 4	S 4	0	0
17	6	1	Total 8	Fe 4	S 4	0	0
17	9	1	Total 8	Fe 4	S 4	0	0
17	9	1	Total 8	Fe 4	S 4	0	0
17	B	1	Total 8	Fe 4	S 4	0	0
17	D	1	Total 8	Fe 4	S 4	0	0


*Continued on next page...*



Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
17	D	1	Total	Fe	S	0	0
			8	4	4		
17	D	1	Total	Fe	S	0	0
			8	4	4		
17	G	1	Total	Fe	S	0	0
			8	4	4		
17	O	1	Total	Fe	S	0	0
			8	4	4		
17	O	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 18 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C<sub>17</sub>H<sub>21</sub>N<sub>4</sub>O<sub>9</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
18	1	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
18	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 19 is FE2/S2 (INORGANIC) CLUSTER (three-letter code: FES) (formula: Fe<sub>2</sub>S<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
19	2	1	Total Fe S 4 2 2	0	0
19	3	1	Total Fe S 4 2 2	0	0
19	C	1	Total Fe S 4 2 2	0	0
19	D	1	Total Fe S 4 2 2	0	0

- Molecule 20 is Piericidin A (three-letter code: HQH) (formula:  $C_{25}H_{37}NO_4$ ) (labeled as "Ligand of Interest" by author).




<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>				<b>ZeroOcc</b>	<b>AltConf</b>
20	4	1	Total	C	N	O	0	0
			30	25	1	4		
20	E	1	Total	C	N	O	0	0
			30	25	1	4		

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS failed to run properly.

- Molecule 1: NADH-quinone oxidoreductase subunit 1




- Molecule 1: NADH-quinone oxidoreductase subunit 1







• Molecule 2: NADH-quinone oxidoreductase subunit 2




• Molecule 2: NADH-quinone oxidoreductase subunit 2




• Molecule 3: NADH-quinone oxidoreductase subunit 3





● Molecule 3: NADH-kinone oxidoreductase subunit 3


Chain D: 56% 39%




● Molecule 4: NADH-kinone oxidoreductase subunit 4

Chain 4: 48% 45% 6%







• Molecule 4: NADH-quinone oxidoreductase subunit 4





• Molecule 5: NADH-quinone oxidoreductase subunit 5






- Molecule 5: NADH-quinone oxidoreductase subunit 5

Chain F: 47% 45% 5%



- Molecule 6: NADH-quinone oxidoreductase subunit 6

Chain 6: 43% 46% 8%




- Molecule 6: NADH-quinone oxidoreductase subunit 6

Chain G: 41% 48% 8%



- Molecule 7: NADH-quinone oxidoreductase subunit 9

Chain 9: 65% 34% 2%








- Molecule 7: NADH-quinone oxidoreductase subunit 9




- Molecule 8: NADH-quinone oxidoreductase subunit 15



- Molecule 8: NADH-quinone oxidoreductase subunit 15




- Molecule 9: NADH-quinone oxidoreductase subunit 16




- Molecule 9: NADH-quinone oxidoreductase subunit 16







• Molecule 10: NADH-quinone oxidoreductase subunit 7




• Molecule 10: NADH-quinone oxidoreductase subunit 7



• Molecule 11: NADH-quinone oxidoreductase subunit 10



• Molecule 11: NADH-quinone oxidoreductase subunit 10




• Molecule 12: NADH-quinone oxidoreductase subunit 11








● Molecule 14: NADH-quinone oxidoreductase subunit 13



● Molecule 14: NADH-quinone oxidoreductase subunit 13








● Molecule 16: NADH-quinone oxidoreductase subunit 8

Chain Q: 51% 43%



## 4 Data and refinement statistics

EDS failed to run properly - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	95.14Å 336.59Å 262.79Å 90.00° 100.41° 90.00°	Depositor
Resolution (Å)	49.16 – 3.60	Depositor
% Data completeness (in resolution range)	79.8 (49.16-3.60)	Depositor
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.73 (at 3.57Å)	Xtrriage
Refinement program	PHENIX (1.13rc1_2961: ???)	Depositor
R, $R_{free}$	0.218 , 0.235	Depositor
Wilson B-factor (Å <sup>2</sup> )	107.9	Xtrriage
Anisotropy	0.002	Xtrriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.36$ , $\langle L^2 \rangle = 0.18$	Xtrriage
Estimated twinning fraction	0.347 for h,-k,-h-l	Xtrriage
Reported twinning fraction	0.490 for -H,-K,H+L	Depositor
Outliers	0 of 148790 reflections	Xtrriage
Total number of atoms	74146	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	111.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HQH, FMN, SF4, FES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	1	0.27	0/3506	0.47	0/4745
1	B	0.27	0/3506	0.47	0/4745
2	2	0.27	0/1439	0.46	0/1953
2	C	0.26	0/1439	0.48	0/1953
3	3	0.28	0/6035	0.50	0/8185
3	D	0.28	0/6035	0.49	0/8185
4	4	0.28	0/3150	0.48	0/4284
4	E	0.27	0/3150	0.47	0/4284
5	5	0.27	0/1656	0.48	0/2246
5	F	0.28	0/1656	0.48	0/2246
6	6	0.30	0/1319	0.52	0/1786
6	G	0.28	0/1319	0.51	0/1786
7	9	0.32	0/1423	0.51	0/1933
7	O	0.30	0/1423	0.52	0/1933
8	7	0.27	0/1059	0.47	0/1429
8	I	0.26	0/1059	0.47	0/1429
9	W	0.26	0/985	0.47	0/1335
9	X	0.26	0/985	0.46	0/1335
10	A	0.29	0/940	0.49	0/1280
10	P	0.28	0/940	0.48	0/1280
11	J	0.26	0/1206	0.45	0/1649
11	R	0.26	0/1206	0.45	0/1649
12	K	0.28	0/710	0.44	0/962
12	S	0.25	0/710	0.44	0/962
13	L	0.26	0/4741	0.44	0/6460
13	T	0.26	0/4741	0.45	0/6460
14	M	0.27	0/3591	0.47	0/4896
14	U	0.26	0/3591	0.45	0/4896
15	N	0.27	0/3238	0.43	0/4434
15	V	0.27	0/3238	0.44	0/4434
16	H	0.30	0/2935	0.52	0/4014
16	Q	0.29	0/2935	0.51	0/4014



Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
All	All	0.28	0/75866	0.47	0/103182

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
7	9	0	1
7	O	0	1
16	H	0	2
16	Q	0	2
All	All	0	6

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
7	9	21	PRO	Peptide
16	H	217	THR	Peptide
16	H	266	GLY	Peptide
7	O	21	PRO	Peptide
16	Q	217	THR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	1	3417	0	3388	150	0
1	B	3417	0	3388	181	0
2	2	1406	0	1373	68	0
2	C	1406	0	1373	67	0
3	3	5895	0	5930	231	0
3	D	5895	0	5931	253	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	4	3067	0	3049	164	0
4	E	3067	0	3049	217	0
5	5	1607	0	1574	65	0
5	F	1607	0	1574	93	0
6	6	1289	0	1298	80	0
6	G	1289	0	1299	106	0
7	9	1388	0	1383	61	0
7	O	1388	0	1383	82	0
8	7	1031	0	1029	57	0
8	I	1031	0	1029	40	0
9	W	967	0	1010	24	0
9	X	967	0	1010	32	0
10	A	910	0	939	59	0
10	P	910	0	939	58	0
11	J	1183	0	1286	72	0
11	R	1183	0	1286	54	0
12	K	703	0	747	36	0
12	S	703	0	747	29	0
13	L	4604	0	4734	173	0
13	T	4604	0	4734	167	0
14	M	3489	0	3606	138	0
14	U	3489	0	3606	146	0
15	N	3154	0	3343	112	0
15	V	3154	0	3343	121	0
16	H	2838	0	2903	171	0
16	Q	2838	0	2903	179	0
17	1	8	0	0	0	0
17	3	24	0	0	1	0
17	6	8	0	0	0	0
17	9	16	0	0	6	0
17	B	8	0	0	1	0
17	D	24	0	0	5	0
17	G	8	0	0	4	0
17	O	16	0	0	3	0
18	1	31	0	19	4	0
18	B	31	0	19	7	0
19	2	4	0	0	1	0
19	3	4	0	0	0	0
19	C	4	0	0	0	0
19	D	4	0	0	0	0
20	4	30	0	0	3	0
20	E	30	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	74146	0	75224	2994	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 20.

The worst 5 of 2994 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
13:L:355:LEU:HD12	13:L:359:LEU:HG	1.35	1.09
3:3:611:ARG:HA	3:3:624:LEU:O	1.62	0.98
16:Q:43:GLN:HE21	16:Q:45:ARG:NH2	1.61	0.97
16:H:146:LYS:HA	16:H:149:LEU:HB2	1.47	0.94
3:D:651:ARG:NH1	3:D:652:PRO:O	2.03	0.91

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	1	435/438 (99%)	410 (94%)	25 (6%)	0	100	100
1	B	435/438 (99%)	405 (93%)	30 (7%)	0	100	100
2	2	176/181 (97%)	167 (95%)	9 (5%)	0	100	100
2	C	176/181 (97%)	169 (96%)	7 (4%)	0	100	100
3	3	750/783 (96%)	702 (94%)	48 (6%)	0	100	100
3	D	750/783 (96%)	699 (93%)	51 (7%)	0	100	100
4	4	382/409 (93%)	358 (94%)	24 (6%)	0	100	100
4	E	382/409 (93%)	357 (94%)	25 (6%)	0	100	100
5	5	194/207 (94%)	187 (96%)	7 (4%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
5	F	194/207 (94%)	186 (96%)	8 (4%)	0	100	100
6	6	164/181 (91%)	153 (93%)	11 (7%)	0	100	100
6	G	164/181 (91%)	149 (91%)	15 (9%)	0	100	100
7	9	178/182 (98%)	167 (94%)	11 (6%)	0	100	100
7	O	178/182 (98%)	168 (94%)	10 (6%)	0	100	100
8	7	125/129 (97%)	117 (94%)	8 (6%)	0	100	100
8	I	125/129 (97%)	114 (91%)	11 (9%)	0	100	100
9	W	125/131 (95%)	121 (97%)	4 (3%)	0	100	100
9	X	125/131 (95%)	120 (96%)	5 (4%)	0	100	100
10	A	115/119 (97%)	103 (90%)	12 (10%)	0	100	100
10	P	115/119 (97%)	103 (90%)	12 (10%)	0	100	100
11	J	158/176 (90%)	143 (90%)	15 (10%)	0	100	100
11	R	158/176 (90%)	144 (91%)	14 (9%)	0	100	100
12	K	93/95 (98%)	88 (95%)	5 (5%)	0	100	100
12	S	93/95 (98%)	90 (97%)	3 (3%)	0	100	100
13	L	603/606 (100%)	568 (94%)	34 (6%)	1 (0%)	47	79
13	T	603/606 (100%)	572 (95%)	30 (5%)	1 (0%)	47	79
14	M	465/469 (99%)	435 (94%)	30 (6%)	0	100	100
14	U	465/469 (99%)	434 (93%)	31 (7%)	0	100	100
15	N	425/427 (100%)	400 (94%)	25 (6%)	0	100	100
15	V	425/427 (100%)	403 (95%)	22 (5%)	0	100	100
16	H	351/365 (96%)	297 (85%)	48 (14%)	6 (2%)	9	45
16	Q	351/365 (96%)	298 (85%)	49 (14%)	4 (1%)	14	53
All	All	9478/9796 (97%)	8827 (93%)	639 (7%)	12 (0%)	51	83

5 of 12 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
16	H	51	VAL
16	Q	44	VAL
16	Q	51	VAL
16	H	44	VAL
16	H	50	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	1	355/356 (100%)	343 (97%)	12 (3%)	37	69
1	B	355/356 (100%)	344 (97%)	11 (3%)	40	71
2	2	150/152 (99%)	146 (97%)	4 (3%)	44	73
2	C	150/152 (99%)	144 (96%)	6 (4%)	31	65
3	3	609/628 (97%)	599 (98%)	10 (2%)	62	83
3	D	609/628 (97%)	598 (98%)	11 (2%)	59	81
4	4	332/355 (94%)	324 (98%)	8 (2%)	49	75
4	E	332/355 (94%)	325 (98%)	7 (2%)	53	78
5	5	167/175 (95%)	164 (98%)	3 (2%)	59	81
5	F	167/175 (95%)	162 (97%)	5 (3%)	41	71
6	6	135/149 (91%)	125 (93%)	10 (7%)	13	46
6	G	135/149 (91%)	127 (94%)	8 (6%)	19	55
7	9	148/150 (99%)	146 (99%)	2 (1%)	67	85
7	O	148/150 (99%)	146 (99%)	2 (1%)	67	85
8	7	104/106 (98%)	103 (99%)	1 (1%)	76	88
8	I	104/106 (98%)	103 (99%)	1 (1%)	76	88
9	W	99/101 (98%)	98 (99%)	1 (1%)	76	88
9	X	99/101 (98%)	97 (98%)	2 (2%)	55	79
10	A	90/92 (98%)	88 (98%)	2 (2%)	52	77
10	P	90/92 (98%)	88 (98%)	2 (2%)	52	77
11	J	118/130 (91%)	115 (98%)	3 (2%)	47	75
11	R	118/130 (91%)	114 (97%)	4 (3%)	37	69
12	K	71/71 (100%)	69 (97%)	2 (3%)	43	72
12	S	71/71 (100%)	69 (97%)	2 (3%)	43	72
13	L	453/454 (100%)	445 (98%)	8 (2%)	59	81
13	T	453/454 (100%)	443 (98%)	10 (2%)	52	77

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
14	M	332/332 (100%)	327 (98%)	5 (2%)	65	84
14	U	332/332 (100%)	324 (98%)	8 (2%)	49	75
15	N	302/302 (100%)	297 (98%)	5 (2%)	60	82
15	V	302/302 (100%)	295 (98%)	7 (2%)	50	76
16	H	293/300 (98%)	284 (97%)	9 (3%)	40	71
16	Q	293/300 (98%)	285 (97%)	8 (3%)	44	73
All	All	7516/7706 (98%)	7337 (98%)	179 (2%)	49	75

5 of 179 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
16	H	196	PHE
2	C	116	LEU
15	V	50	PHE
16	H	307	ARG
1	B	353	CYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 57 such sidechains are listed below:

Mol	Chain	Res	Type
14	M	255	GLN
2	C	71	GLN
14	U	221	ASN
15	N	245	ASN
16	H	112	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

22 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
17	SF4	9	202	7	0,12,12	0.00	-	-		
17	SF4	3	802	3	0,12,12	0.00	-	-		
17	SF4	B	501	1	0,12,12	0.00	-	-		
18	FMN	1	502	-	31,33,33	1.45	4 (12%)	40,50,50	1.58	5 (12%)
20	HQH	E	501	-	30,30,30	1.38	3 (10%)	36,40,40	1.91	7 (19%)
17	SF4	D	802	3	0,12,12	0.00	-	-		
17	SF4	3	803	3	0,12,12	0.00	-	-		
19	FES	D	804	3	0,4,4	0.00	-	-		
17	SF4	O	202	7	0,12,12	0.00	-	-		
19	FES	C	201	2	0,4,4	0.00	-	-		
17	SF4	9	201	7	0,12,12	0.00	-	-		
20	HQH	4	501	-	30,30,30	1.39	3 (10%)	36,40,40	1.73	9 (25%)
17	SF4	D	801	3	0,12,12	0.00	-	-		
17	SF4	3	801	3	0,12,12	0.00	-	-		
19	FES	3	804	3	0,4,4	0.00	-	-		
17	SF4	1	501	1	0,12,12	0.00	-	-		
18	FMN	B	502	-	31,33,33	1.42	4 (12%)	40,50,50	1.63	6 (15%)
17	SF4	G	201	6	0,12,12	0.00	-	-		
17	SF4	D	803	3	0,12,12	0.00	-	-		
17	SF4	O	201	7	0,12,12	0.00	-	-		
17	SF4	6	201	6	0,12,12	0.00	-	-		
19	FES	2	201	2	0,4,4	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	SF4	9	202	7	-	-	0/6/5/5
17	SF4	3	802	3	-	-	0/6/5/5

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
17	SF4	B	501	1	-	-	0/6/5/5
18	FMN	1	502	-	-	8/18/18/18	0/3/3/3
20	HQH	E	501	-	-	13/29/29/29	0/1/1/1
17	SF4	D	802	3	-	-	0/6/5/5
17	SF4	3	803	3	-	-	0/6/5/5
19	FES	D	804	3	-	-	0/1/1/1
18	FMN	B	502	-	-	8/18/18/18	0/3/3/3
19	FES	C	201	2	-	-	0/1/1/1
17	SF4	9	201	7	-	-	0/6/5/5
20	HQH	4	501	-	-	12/29/29/29	0/1/1/1
17	SF4	D	801	3	-	-	0/6/5/5
17	SF4	3	801	3	-	-	0/6/5/5
19	FES	3	804	3	-	-	0/1/1/1
17	SF4	1	501	1	-	-	0/6/5/5
17	SF4	O	202	7	-	-	0/6/5/5
17	SF4	G	201	6	-	-	0/6/5/5
17	SF4	D	803	3	-	-	0/6/5/5
17	SF4	O	201	7	-	-	0/6/5/5
17	SF4	6	201	6	-	-	0/6/5/5
19	FES	2	201	2	-	-	0/1/1/1

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
20	4	501	HQH	C24-N5	5.65	1.40	1.32
20	E	501	HQH	C24-N5	5.54	1.40	1.32
18	1	502	FMN	C10-N1	4.20	1.38	1.33
18	B	502	FMN	C10-N1	4.06	1.38	1.33
18	1	502	FMN	C4A-N5	3.47	1.38	1.33

The worst 5 of 27 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
20	E	501	HQH	C29-O2-C24	-7.71	109.58	117.21
20	4	501	HQH	C29-O2-C24	-5.99	111.28	117.21
18	1	502	FMN	C4-N3-C2	5.51	119.79	115.14
18	B	502	FMN	C4-N3-C2	5.39	119.69	115.14
18	B	502	FMN	C1'-N10-C10	4.79	122.70	118.41

There are no chirality outliers.

5 of 41 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
18	1	502	FMN	N10-C1'-C2'-O2'
18	1	502	FMN	N10-C1'-C2'-C3'
18	1	502	FMN	C1'-C2'-C3'-O3'
18	1	502	FMN	C1'-C2'-C3'-C4'
18	1	502	FMN	O4'-C4'-C5'-O5'


There are no ring outliers.


14 monomers are involved in 36 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
17	9	202	SF4	3	0
17	B	501	SF4	1	0
18	1	502	FMN	4	0
20	E	501	HQH	1	0
17	O	202	SF4	2	0
17	9	201	SF4	3	0
20	4	501	HQH	3	0
17	D	801	SF4	2	0
17	3	801	SF4	1	0
18	B	502	FMN	7	0
17	G	201	SF4	4	0
17	D	803	SF4	3	0
17	O	201	SF4	1	0
19	2	201	FES	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS failed to run properly - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS failed to run properly - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS failed to run properly - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS failed to run properly - this section is therefore empty.