



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 21, 2024 – 04:36 PM EST

PDB ID : 4QF0  
Title : Structure of a 16 nm protein cage designed by fusing symmetric oligomeric domains, quadruple mutant, P21212 form  
Authors : Lai, Y.-T.; Yeates, T.O.  
Deposited on : 2014-05-19  
Resolution : 6.49 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 6.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1000 (9.00-3.90)
Clashscore	141614	1064 (9.00-3.90)
Ramachandran outliers	138981	1012 (9.00-3.88)
Sidechain outliers	138945	1010 (9.00-3.84)
RSRZ outliers	127900	1002 (9.00-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	456	
1	B	456	
1	C	456	
1	D	456	
1	E	456	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	456	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into three segments: a red segment on the left labeled '46%', a green segment in the middle labeled '88%', and a yellow segment on the right labeled '9%'. A small grey dot is visible at the far right end of the bar.</p>

## 2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 20256 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-haem bromoperoxidase BPO-A2, Matrix protein 1 chimera.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	441	3376	2151	569	648	8	0	0	0
1	B	441	3376	2151	569	648	8	0	0	0
1	C	441	3376	2151	569	648	8	0	0	0
1	D	441	3376	2151	569	648	8	0	0	0
1	E	441	3376	2151	569	648	8	0	0	0
1	F	441	3376	2151	569	648	8	0	0	0

There are 114 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	THR	GLN	engineered mutation	UNP P29715
A	51	ALA	TYR	engineered mutation	UNP P29715
A	118	ALA	LYS	engineered mutation	UNP P29715
A	278	ALA	-	linker	UNP P03485
A	279	GLN	-	linker	UNP P03485
A	280	GLU	-	linker	UNP P03485
A	281	ALA	-	linker	UNP P03485
A	282	GLN	-	linker	UNP P03485
A	283	LYS	-	linker	UNP P03485
A	284	GLN	-	linker	UNP P03485
A	285	LYS	-	linker	UNP P03485
A	448	LEU	-	expression tag	UNP P03485
A	449	GLU	-	expression tag	UNP P03485
A	450	HIS	-	expression tag	UNP P03485
A	451	HIS	-	expression tag	UNP P03485
A	452	HIS	-	expression tag	UNP P03485
A	453	HIS	-	expression tag	UNP P03485

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	454	HIS	-	expression tag	UNP P03485
A	455	HIS	-	expression tag	UNP P03485
B	24	THR	GLN	engineered mutation	UNP P29715
B	51	ALA	TYR	engineered mutation	UNP P29715
B	118	ALA	LYS	engineered mutation	UNP P29715
B	278	ALA	-	linker	UNP P03485
B	279	GLN	-	linker	UNP P03485
B	280	GLU	-	linker	UNP P03485
B	281	ALA	-	linker	UNP P03485
B	282	GLN	-	linker	UNP P03485
B	283	LYS	-	linker	UNP P03485
B	284	GLN	-	linker	UNP P03485
B	285	LYS	-	linker	UNP P03485
B	448	LEU	-	expression tag	UNP P03485
B	449	GLU	-	expression tag	UNP P03485
B	450	HIS	-	expression tag	UNP P03485
B	451	HIS	-	expression tag	UNP P03485
B	452	HIS	-	expression tag	UNP P03485
B	453	HIS	-	expression tag	UNP P03485
B	454	HIS	-	expression tag	UNP P03485
B	455	HIS	-	expression tag	UNP P03485
C	24	THR	GLN	engineered mutation	UNP P29715
C	51	ALA	TYR	engineered mutation	UNP P29715
C	118	ALA	LYS	engineered mutation	UNP P29715
C	278	ALA	-	linker	UNP P03485
C	279	GLN	-	linker	UNP P03485
C	280	GLU	-	linker	UNP P03485
C	281	ALA	-	linker	UNP P03485
C	282	GLN	-	linker	UNP P03485
C	283	LYS	-	linker	UNP P03485
C	284	GLN	-	linker	UNP P03485
C	285	LYS	-	linker	UNP P03485
C	448	LEU	-	expression tag	UNP P03485
C	449	GLU	-	expression tag	UNP P03485
C	450	HIS	-	expression tag	UNP P03485
C	451	HIS	-	expression tag	UNP P03485
C	452	HIS	-	expression tag	UNP P03485
C	453	HIS	-	expression tag	UNP P03485
C	454	HIS	-	expression tag	UNP P03485
C	455	HIS	-	expression tag	UNP P03485
D	24	THR	GLN	engineered mutation	UNP P29715
D	51	ALA	TYR	engineered mutation	UNP P29715

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	118	ALA	LYS	engineered mutation	UNP P29715
D	278	ALA	-	linker	UNP P03485
D	279	GLN	-	linker	UNP P03485
D	280	GLU	-	linker	UNP P03485
D	281	ALA	-	linker	UNP P03485
D	282	GLN	-	linker	UNP P03485
D	283	LYS	-	linker	UNP P03485
D	284	GLN	-	linker	UNP P03485
D	285	LYS	-	linker	UNP P03485
D	448	LEU	-	expression tag	UNP P03485
D	449	GLU	-	expression tag	UNP P03485
D	450	HIS	-	expression tag	UNP P03485
D	451	HIS	-	expression tag	UNP P03485
D	452	HIS	-	expression tag	UNP P03485
D	453	HIS	-	expression tag	UNP P03485
D	454	HIS	-	expression tag	UNP P03485
D	455	HIS	-	expression tag	UNP P03485
E	24	THR	GLN	engineered mutation	UNP P29715
E	51	ALA	TYR	engineered mutation	UNP P29715
E	118	ALA	LYS	engineered mutation	UNP P29715
E	278	ALA	-	linker	UNP P03485
E	279	GLN	-	linker	UNP P03485
E	280	GLU	-	linker	UNP P03485
E	281	ALA	-	linker	UNP P03485
E	282	GLN	-	linker	UNP P03485
E	283	LYS	-	linker	UNP P03485
E	284	GLN	-	linker	UNP P03485
E	285	LYS	-	linker	UNP P03485
E	448	LEU	-	expression tag	UNP P03485
E	449	GLU	-	expression tag	UNP P03485
E	450	HIS	-	expression tag	UNP P03485
E	451	HIS	-	expression tag	UNP P03485
E	452	HIS	-	expression tag	UNP P03485
E	453	HIS	-	expression tag	UNP P03485
E	454	HIS	-	expression tag	UNP P03485
E	455	HIS	-	expression tag	UNP P03485
F	24	THR	GLN	engineered mutation	UNP P29715
F	51	ALA	TYR	engineered mutation	UNP P29715
F	118	ALA	LYS	engineered mutation	UNP P29715
F	278	ALA	-	linker	UNP P03485
F	279	GLN	-	linker	UNP P03485
F	280	GLU	-	linker	UNP P03485

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
F	281	ALA	-	linker	UNP P03485
F	282	GLN	-	linker	UNP P03485
F	283	LYS	-	linker	UNP P03485
F	284	GLN	-	linker	UNP P03485
F	285	LYS	-	linker	UNP P03485
F	448	LEU	-	expression tag	UNP P03485
F	449	GLU	-	expression tag	UNP P03485
F	450	HIS	-	expression tag	UNP P03485
F	451	HIS	-	expression tag	UNP P03485
F	452	HIS	-	expression tag	UNP P03485
F	453	HIS	-	expression tag	UNP P03485
F	454	HIS	-	expression tag	UNP P03485
F	455	HIS	-	expression tag	UNP P03485

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Non-haem bromoperoxidase BPO-A2, Matrix protein 1 chimera





- Molecule 1: Non-haem bromoperoxidase BPO-A2, Matrix protein 1 chimera










- Molecule 1: Non-haem bromoperoxidase BPO-A2, Matrix protein 1 chimera





- Molecule 1: Non-haem bromoperoxidase BPO-A2, Matrix protein 1 chimera








• Molecule 1: Non-haem bromoperoxidase BPO-A2, Matrix protein 1 chimera



• Molecule 1: Non-haem bromoperoxidase BPO-A2, Matrix protein 1 chimera





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	175.50Å 147.73Å 167.63Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.71 – 6.49 93.71 – 6.49	Depositor EDS
% Data completeness (in resolution range)	99.6 (93.71-6.49) 99.6 (93.71-6.49)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.07 (at 6.72Å)	Xtrriage
Refinement program	REFMAC 5.8.0071	Depositor
R, $R_{free}$	0.271 , 0.324 0.273 , 0.331	Depositor DCC
$R_{free}$ test set	451 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	398.2	Xtrriage
Anisotropy	0.081	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.39 , 410.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.45$ , $\langle L^2 \rangle = 0.28$	Xtrriage
Estimated twinning fraction	0.003 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.86	EDS
Total number of atoms	20256	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	313.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.41	0/3452	0.55	0/4702
1	B	0.40	0/3452	0.56	0/4702
1	C	0.41	0/3452	0.56	0/4702
1	D	0.40	0/3452	0.56	0/4702
1	E	0.40	0/3452	0.56	0/4702
1	F	0.41	0/3452	0.56	0/4702
All	All	0.40	0/20712	0.56	0/28212

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3376	0	3286	16	0
1	B	3376	0	3286	16	0
1	C	3376	0	3286	15	0
1	D	3376	0	3286	16	0
1	E	3376	0	3286	18	0
1	F	3376	0	3286	18	0
All	All	20256	0	19716	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 92 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:30:HIS:CE1	1:F:34:LEU:O	2.28	0.85
1:B:30:HIS:CE1	1:B:34:LEU:O	2.32	0.82
1:A:30:HIS:CE1	1:A:34:LEU:O	2.33	0.81
1:C:30:HIS:CE1	1:C:34:LEU:O	2.33	0.81
1:E:30:HIS:CE1	1:E:34:LEU:O	2.38	0.77

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	439/456 (96%)	413 (94%)	26 (6%)	0	100	100
1	B	439/456 (96%)	413 (94%)	26 (6%)	0	100	100
1	C	439/456 (96%)	414 (94%)	25 (6%)	0	100	100
1	D	439/456 (96%)	413 (94%)	26 (6%)	0	100	100
1	E	439/456 (96%)	414 (94%)	25 (6%)	0	100	100
1	F	439/456 (96%)	414 (94%)	25 (6%)	0	100	100
All	All	2634/2736 (96%)	2481 (94%)	153 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	348/369 (94%)	336 (97%)	12 (3%)	37	60
1	B	348/369 (94%)	337 (97%)	11 (3%)	39	61
1	C	348/369 (94%)	337 (97%)	11 (3%)	39	61
1	D	348/369 (94%)	337 (97%)	11 (3%)	39	61
1	E	348/369 (94%)	337 (97%)	11 (3%)	39	61
1	F	348/369 (94%)	337 (97%)	11 (3%)	39	61
All	All	2088/2214 (94%)	2021 (97%)	67 (3%)	39	61

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	136	PRO
1	F	217	ASP
1	F	338	LEU
1	C	136	PRO
1	C	91	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	188	ASN
1	E	188	ASN
1	F	188	ASN
1	B	188	ASN
1	A	188	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	441/456 (96%)	2.63	244 (55%) 0 0	121, 325, 340, 340	0
1	B	441/456 (96%)	2.21	215 (48%) 0 1	122, 314, 340, 340	0
1	C	441/456 (96%)	1.87	169 (38%) 0 1	122, 313, 340, 340	0
1	D	441/456 (96%)	2.96	258 (58%) 0 0	122, 321, 340, 340	0
1	E	441/456 (96%)	2.71	251 (56%) 0 0	122, 320, 340, 340	0
1	F	441/456 (96%)	2.18	208 (47%) 0 1	121, 317, 340, 340	0
All	All	2646/2736 (96%)	2.43	1345 (50%) 0 1	121, 318, 340, 340	0

The worst 5 of 1345 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	358	GLN	14.3
1	A	359	ARG	14.1
1	D	33	PRO	14.0
1	A	66	GLN	13.5
1	B	359	ARG	13.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.