



Full wwPDB NMR Structure Validation Report ⓘ

May 31, 2020 – 11:53 am BST

PDB ID : 6QH2
Title : Solution NMR ensemble for a chimeric KH-S1 domain construct of exosomal polynucleotide phosphrylase at 298K compiled using the CoMAND method
Authors : ElGamacy, M.; Truffault, V.; Zhu, H.; Coles, M.
Deposited on : 2019-01-14

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 70%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	171	

2 Ensemble composition and analysis

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:15-A:76 (62)	0.56	18
2	A:79-A:153 (75)	0.52	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 4 single-model clusters were found.

Cluster number	Models
1	3, 6, 8, 9, 10, 12, 13, 16
2	1, 11, 19, 20
3	2, 14
4	5, 18
Single-model clusters	4; 7; 15; 17

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 2238 atoms, of which 1144 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Polyribonucleotide nucleotidyltransferase.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	142	2238	683	1144	197	213	1	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP B6I1N9
A	2	ALA	-	expression tag	UNP B6I1N9
A	3	MET	-	expression tag	UNP B6I1N9
A	4	GLY	-	expression tag	UNP B6I1N9

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Polyribonucleotide nucleotidyltransferase



4.2.2 Score per residue for model 2

- Molecule 1: Polyribonucleotide nucleotidyltransferase




PRO
ALA
ALA
GLY
GLN
GLY
GLU

4.2.3 Score per residue for model 3

- Molecule 1: Polyribonucleotide nucleotidyltransferase

Chain A: 68% 9% 17%



GLN
PRO
ALA
ALA
GLY
GLY
PRO
GLY
GLU
ALA
ALA
PRO
SER
GLU
ALA
ALA
GLY
GLU
ALA
GLY
GLN
GLY
GLU

4.2.4 Score per residue for model 4

- Molecule 1: Polyribonucleotide nucleotidyltransferase

Chain A: 68% 9% 17%



GLN
PRO
ALA
ALA
GLY
GLY
PRO
GLY
GLU
ALA
ALA
PRO
SER
GLU
ALA
ALA
GLY
GLU
ALA
GLY
GLN
GLY
GLU

4.2.5 Score per residue for model 5

- Molecule 1: Polyribonucleotide nucleotidyltransferase

Chain A: 70% 8% 17%




ALA
ALA
GLU
GLN
GLY
GLY
GLU

4.2.6 Score per residue for model 6

- Molecule 1: Polyribonucleotide nucleotidyltransferase


Chain A: 70% 8% 17%



4.2.7 Score per residue for model 7 (medoid)

- Molecule 1: Polyribonucleotide nucleotidyltransferase


Chain A: 72% 6% 17%



4.2.8 Score per residue for model 8

- Molecule 1: Polyribonucleotide nucleotidyltransferase


Chain A: 71% 8% 17%



4.2.9 Score per residue for model 9


- Molecule 1: Polyribonucleotide nucleotidyltransferase

Chain A: 66% 11% 17%




4.2.10 Score per residue for model 10

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.11 Score per residue for model 11

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.12 Score per residue for model 12

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.13 Score per residue for model 13

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.14 Score per residue for model 14

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.15 Score per residue for model 15

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.16 Score per residue for model 16

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.17 Score per residue for model 17

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.18 Score per residue for model 18

- Molecule 1: Polyribonucleotide nucleotidyltransferase




4.2.19 Score per residue for model 19

- Molecule 1: Polyribonucleotide nucleotidyltransferase



4.2.20 Score per residue for model 20

- Molecule 1: Polyribonucleotide nucleotidyltransferase



5 Refinement protocol and experimental data overview

The models were refined using the following method: *R-factor based frame picking*.

Of the 10000 calculated structures, 20 were deposited, based on the following criterion: *back calculated data agree with experimental NOESY spectrum*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Shine	structure calculation	
Rosetta	structure calculation	3.6
NAMD	refinement	2.12
CoMAND	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 6 of this report.

Chemical shift file(s)	input_cs.cif
Number of chemical shift lists	1
Total number of shifts	1440
Number of shifts mapped to atoms	1440
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	70%

No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	1051	1106	1106	0±1
All	All	21020	22120	22120	10

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:36:ILE:HD11	1:A:49:ILE:HG22	0.58	1.75	13	1
1:A:43:THR:HG21	1:A:68:ALA:HA	0.47	1.85	16	1
1:A:47:ILE:HG23	1:A:57:ILE:HD13	0.46	1.88	10	1
1:A:30:GLY:HA3	1:A:35:VAL:H	0.46	1.69	16	1
1:A:122:VAL:HG12	1:A:126:LEU:HD12	0.45	1.89	18	1
1:A:136:VAL:HG22	1:A:147:LEU:HD23	0.45	1.89	16	1
1:A:59:ALA:CB	1:A:65:ALA:HB2	0.43	2.43	4	1
1:A:29:ILE:O	1:A:29:ILE:HG22	0.43	2.13	10	2
1:A:97:ALA:O	1:A:109:VAL:N	0.41	2.53	14	1

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/171 (80%)	133±1 (97±1%)	3±1 (2±1%)	1±1 (1±1%)	26	73
All	All	2740/3420 (80%)	2658 (97%)	63 (2%)	19 (1%)	26	73

All 10 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	51	ASP	5
1	A	102	GLY	3
1	A	91	ARG	3
1	A	59	ALA	2
1	A	111	ILE	1
1	A	105	LYS	1
1	A	145	ILE	1
1	A	139	VAL	1
1	A	30	GLY	1
1	A	52	ASP	1

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	113/134 (84%)	110±1 (97±1%)	3±1 (3±1%)	46 90
All	All	2260/2680 (84%)	2197 (97%)	63 (3%)	46 90

All 30 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	134	VAL	7
1	A	140	ASP	6
1	A	123	THR	5
1	A	132	VAL	4
1	A	146	ARG	3
1	A	90	THR	3
1	A	45	THR	3
1	A	56	LYS	3
1	A	151	GLU	2
1	A	142	GLN	2
1	A	118	ARG	2
1	A	86	THR	2
1	A	144	ARG	2
1	A	133	PRO	2
1	A	91	ARG	2
1	A	141	ARG	1
1	A	67	HIS	1
1	A	26	LYS	1
1	A	36	ILE	1
1	A	17	THR	1
1	A	153	THR	1
1	A	121	LYS	1
1	A	128	MET	1
1	A	113	GLN	1
1	A	108	LEU	1
1	A	61	ASP	1
1	A	94	ASP	1
1	A	18	ILE	1
1	A	127	GLN	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	51	ASP	1

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

There are no ligands in this entry.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 70% for the well-defined parts and 69% for the entire structure.

6.1 Chemical shift list 1

File name: input_cs.cif

Chemical shift list name: *kh-s1_bmr_bBA780W.str*

6.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1440
Number of shifts mapped to atoms	1440
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

6.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
¹³ C _α	143	-0.40 \pm 0.12	None needed (< 0.5 ppm)
¹³ C _β	128	0.07 \pm 0.15	None needed (< 0.5 ppm)
¹³ C'	126	-0.28 \pm 0.10	None needed (< 0.5 ppm)
¹⁵ N	131	0.35 \pm 0.39	None needed (< 0.5 ppm)

6.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 70%, i.e. 1169 atoms were assigned a chemical shift out of a possible 1669. 20 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	644/681 (95%)	260/272 (96%)	258/274 (94%)	126/135 (93%)
Sidechain	517/933 (55%)	215/536 (40%)	300/352 (85%)	2/45 (4%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	8/55 (15%)	4/30 (13%)	4/22 (18%)	0/3 (0%)
Overall	1169/1669 (70%)	479/838 (57%)	562/648 (87%)	128/183 (70%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 69%, i.e. 1194 atoms were assigned a chemical shift out of a possible 1738. 20 out of 20 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	660/706 (93%)	266/282 (94%)	265/284 (93%)	129/140 (92%)
Sidechain	526/977 (54%)	216/562 (38%)	308/366 (84%)	2/49 (4%)
Aromatic	8/55 (15%)	4/30 (13%)	4/22 (18%)	0/3 (0%)
Overall	1194/1738 (69%)	486/874 (56%)	577/672 (86%)	131/192 (68%)

6.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

6.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

