



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 2, 2023 – 03:43 PM EDT

PDB ID : 3QI2
Title : A Galpha P-loop mutation prevents transition to the activated state: G42R bound to RGS14 GoLoco
Authors : Bosch, D.E.; Willard, F.S.; Kimple, A.J.; Miley, M.J.; Siderovski, D.P.
Deposited on : 2011-01-26
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 76%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 21%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 3%; margin-top: 5px;">76% 21% .</p>
1	B	328	<div style="display: flex; align-items: center;"> <div style="width: 8%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 74%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 22%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 0%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 3%; margin-top: 5px;">74% 22% .</p>
2	C	36	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 67%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 25%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 8%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 3%; margin-top: 5px;">67% 25% 8%</p>
2	D	36	<div style="display: flex; align-items: center;"> <div style="width: 25%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 50%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 33%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: grey; margin-right: 5px;"></div> </div> <p style="margin-left: 3%; margin-top: 5px;">50% 33% 17%</p>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	A	2	-	-	-	X

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 5714 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Guanine nucleotide-binding protein G(i) subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	318	2560	1625	434	486	15	0	0	0
1	B	314	2530	1609	426	480	15	0	0	0

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	SER	-	expression tag	UNP P63096
A	28	ASN	-	expression tag	UNP P63096
A	29	ALA	-	expression tag	UNP P63096
A	30	GLY	-	expression tag	UNP P63096
A	42	ARG	GLY	engineered mutation	UNP P63096
B	27	SER	-	expression tag	UNP P63096
B	28	ASN	-	expression tag	UNP P63096
B	29	ALA	-	expression tag	UNP P63096
B	30	GLY	-	expression tag	UNP P63096
B	42	ARG	GLY	engineered mutation	UNP P63096

- Molecule 2 is a protein called Regulator of G-protein signaling 14.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	C	33	269	169	48	52	0	0	0
2	D	30	246	155	44	47	0	0	0

- Molecule 3 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: C₁₀H₁₅N₅O₁₁P₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
			Total	C	N	O			P
3	A	1	28	10	5	11	2	0	0
3	B	1	28	10	5	11	2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O S		
4	A	1	5	4 1	0	0
4	B	1	5	4 1	0	0

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	C O	0	0
			6	3 3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	19	Total	O	0	0
			19	19		
6	B	13	Total	O	0	0
			13	13		
6	C	2	Total	O	0	0
			2	2		
6	D	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Guanine nucleotide-binding protein G(i) subunit alpha-1




- Molecule 1: Guanine nucleotide-binding protein G(i) subunit alpha-1



- Molecule 2: Regulator of G-protein signaling 14



- Molecule 2: Regulator of G-protein signaling 14



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	70.00Å 131.02Å 203.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.70 – 2.80 29.71 – 2.80	Depositor EDS
% Data completeness (in resolution range)	78.4 (29.70-2.80) 78.5 (29.71-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.55 (at 2.80Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.6_289)	Depositor
R, R_{free}	0.196 , 0.273 0.194 , 0.274	Depositor DCC
R_{free} test set	952 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	41.9	Xtrriage
Anisotropy	0.412	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 58.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5714	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.79% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, GDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.22	0/2606	0.36	0/3510
1	B	0.22	0/2575	0.35	0/3469
2	C	0.20	0/270	0.35	0/361
2	D	0.21	0/246	0.33	0/328
All	All	0.22	0/5697	0.35	0/7668

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2560	0	2537	43	0
1	B	2530	0	2507	44	0
2	C	269	0	273	8	0
2	D	246	0	254	9	0
3	A	28	0	12	1	0
3	B	28	0	12	0	0
4	A	5	0	0	0	0
4	B	5	0	0	0	0
5	A	6	0	8	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	19	0	0	0	0
6	B	13	0	0	0	0
6	C	2	0	0	0	0
6	D	3	0	0	0	0
All	All	5714	0	5603	94	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (94) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:211:TRP:HB3	1:B:215:PHE:HE1	1.51	0.76
1:A:112:GLY:HA2	2:C:524:LEU:HD13	1.69	0.75
1:A:149:ASN:H	2:C:515:GLN:HE22	1.38	0.72
1:B:252:SER:HB3	2:D:500:LEU:HD22	1.72	0.71
1:B:181:THR:HG21	1:B:202:GLY:HA3	1.73	0.69
2:D:526:LEU:HD12	2:D:527:PRO:HD2	1.76	0.67
1:B:38:LEU:HD23	1:B:222:ILE:HB	1.76	0.66
1:B:42:ARG:HG2	1:B:43:GLU:HG2	1.78	0.66
1:B:211:TRP:HB3	1:B:215:PHE:CE1	2.31	0.66
2:C:526:LEU:HD12	2:C:527:PRO:HD2	1.80	0.64
1:A:38:LEU:HD23	1:A:222:ILE:HB	1.80	0.62
1:B:235:ALA:HB3	1:B:238:GLU:HG2	1.81	0.62
1:A:61:TYR:H	1:A:171:GLN:HE21	1.48	0.61
1:A:183:GLY:HA2	1:A:204:GLN:HG2	1.82	0.61
1:B:54:LYS:O	1:B:58:GLU:HB2	2.02	0.59
1:A:170:THR:HG23	1:A:173:ASP:H	1.68	0.58
1:A:324:THR:HG22	1:A:335:VAL:HG21	1.84	0.58
1:B:302:TYR:O	1:B:306:GLN:HG2	2.04	0.57
1:B:227:LEU:HD21	1:B:268:LEU:HB3	1.86	0.57
1:A:240:MET:HG3	1:A:241:ASN:H	1.70	0.56
1:B:255:ASN:ND2	1:B:312:LYS:HG2	2.21	0.56
1:A:215:PHE:O	1:A:218:VAL:HG12	2.06	0.56
1:A:208:ARG:O	1:A:212:ILE:HG12	2.06	0.55
1:A:170:THR:HG22	1:A:173:ASP:CG	2.26	0.55
1:B:120:THR:HG23	1:B:123:LEU:H	1.69	0.55
1:B:158:ASP:O	1:B:162:ILE:HG12	2.06	0.54
1:B:188:HIS:ND1	1:B:197:LYS:HG2	2.23	0.53
1:A:81:ILE:O	1:A:85:ILE:HG12	2.08	0.53
1:B:81:ILE:O	1:B:85:ILE:HG12	2.10	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:239:GLU:HG3	1:B:240:MET:H	1.74	0.52
1:B:112:GLY:HA2	2:D:524:LEU:HD13	1.92	0.52
2:D:519:LEU:HD21	2:D:524:LEU:HD21	1.92	0.52
1:B:55:ILE:HA	1:B:60:GLY:HA2	1.92	0.51
1:A:203:GLY:HA2	2:C:510:SER:H	1.75	0.51
1:A:110:LEU:HD23	1:B:68:GLN:HE22	1.76	0.51
2:C:505:ASN:O	2:C:509:SER:HB2	2.10	0.51
1:A:311:ASN:HD21	1:A:317:LYS:HB2	1.77	0.49
1:A:76:ASN:HD22	1:A:76:ASN:N	2.10	0.49
1:A:178:ARG:HB3	2:C:516:ARG:HD3	1.95	0.49
1:A:282:PRO:HB3	1:A:294:ASN:HD21	1.78	0.48
1:A:256:ASN:HB3	1:A:259:PHE:HD2	1.78	0.48
1:B:45:GLY:O	1:B:49:ILE:HG12	2.13	0.48
1:B:335:VAL:O	1:B:339:VAL:HG23	2.14	0.48
1:A:235:ALA:HB3	1:A:238:GLU:HG2	1.96	0.47
1:B:84:ILE:O	1:B:88:MET:HG3	2.14	0.47
1:A:230:TYR:O	1:A:286:CYS:HB2	2.15	0.47
1:A:275:GLU:O	1:A:278:ILE:HG22	2.15	0.46
1:A:54:LYS:O	1:A:58:GLU:HB2	2.15	0.46
1:B:257:LYS:HG3	1:B:313:ARG:HH22	1.79	0.46
1:A:43:GLU:OE2	3:A:355:GDP:H5'	2.16	0.46
1:A:143:SER:HA	1:A:146:TYR:CE1	2.50	0.46
1:B:311:ASN:HB2	1:B:319:ILE:HD11	1.97	0.46
1:B:183:GLY:HA2	1:B:204:GLN:NE2	2.30	0.46
1:B:52:GLN:NE2	1:B:175:LEU:HD21	2.31	0.46
1:B:250:PHE:HZ	1:B:319:ILE:HD13	1.81	0.46
2:C:519:LEU:HD23	2:C:524:LEU:HD21	1.97	0.45
1:A:55:ILE:HA	1:A:60:GLY:HA2	1.97	0.45
1:A:212:ILE:O	1:A:216:GLU:HG3	2.17	0.45
1:A:249:LEU:O	1:A:253:ILE:HG12	2.16	0.45
1:B:122:GLU:O	1:B:126:VAL:HG23	2.17	0.45
1:A:226:ALA:HB1	1:A:270:LYS:HD2	2.00	0.44
1:B:74:TYR:HE1	1:B:162:ILE:HG22	1.81	0.44
1:A:158:ASP:O	1:A:162:ILE:HG12	2.17	0.44
1:B:265:ILE:HD13	1:B:339:VAL:HG13	1.99	0.44
1:A:54:LYS:HE3	1:A:61:TYR:CZ	2.53	0.44
1:B:54:LYS:HE3	1:B:61:TYR:CZ	2.53	0.44
2:D:503:LEU:HD23	2:D:506:ARG:HH11	1.83	0.44
1:A:52:GLN:NE2	1:A:175:LEU:HD21	2.33	0.43
1:B:63:GLU:O	1:B:67:LYS:HG3	2.18	0.43
1:A:61:TYR:H	1:A:171:GLN:NE2	2.16	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:61:TYR:HA	1:A:65:GLU:OE1	2.19	0.43
1:B:164:GLN:O	1:B:167:TYR:HB3	2.19	0.43
1:B:249:LEU:O	1:B:253:ILE:HG12	2.19	0.42
2:D:527:PRO:HG2	2:D:530:LEU:HD12	2.01	0.42
1:A:184:ILE:HG12	1:A:201:VAL:HG13	2.01	0.42
1:A:250:PHE:HZ	1:A:319:ILE:HD13	1.85	0.42
1:B:257:LYS:HD2	1:B:257:LYS:N	2.34	0.42
1:B:226:ALA:HB1	1:B:270:LYS:HD2	2.02	0.41
1:A:75:SER:HA	2:C:519:LEU:CD1	2.50	0.41
1:B:80:SER:O	1:B:84:ILE:HG13	2.21	0.41
1:B:34:VAL:HG13	1:B:196:PHE:HA	2.02	0.41
1:B:41:ALA:HA	2:D:507:VAL:HG12	2.02	0.41
1:A:63:GLU:O	1:A:67:LYS:HG3	2.20	0.41
1:B:54:LYS:HE3	1:B:61:TYR:CE2	2.56	0.41
1:A:76:ASN:N	1:A:76:ASN:ND2	2.68	0.41
1:A:123:LEU:O	1:A:127:ILE:HG13	2.20	0.41
1:A:132:LYS:C	1:A:132:LYS:HD3	2.41	0.41
1:B:40:GLY:O	2:D:508:GLN:HG2	2.21	0.41
1:B:344:ILE:C	1:B:346:ASN:H	2.23	0.41
1:A:185:VAL:HG23	1:A:200:ASP:HB3	2.03	0.40
1:B:69:TYR:O	1:B:73:VAL:HG23	2.21	0.40
2:D:498:GLU:O	2:D:502:GLU:HG3	2.21	0.40
1:A:211:TRP:HB3	1:A:215:PHE:CE1	2.57	0.40
1:B:100:ARG:HA	1:B:100:ARG:NE	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	316/328 (96%)	297 (94%)	18 (6%)	1 (0%)	41 72

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	310/328 (94%)	295 (95%)	15 (5%)	0	100	100
2	C	29/36 (81%)	27 (93%)	2 (7%)	0	100	100
2	D	26/36 (72%)	24 (92%)	2 (8%)	0	100	100
All	All	681/728 (94%)	643 (94%)	37 (5%)	1 (0%)	51	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	316	THR

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	276/284 (97%)	275 (100%)	1 (0%)	91	97
1	B	274/284 (96%)	274 (100%)	0	100	100
2	C	31/32 (97%)	31 (100%)	0	100	100
2	D	28/32 (88%)	28 (100%)	0	100	100
All	All	609/632 (96%)	608 (100%)	1 (0%)	93	98

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	147	GLN
1	A	164	GLN
1	A	171	GLN
1	A	172	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	195	HIS
1	A	204	GLN
1	A	294	ASN
1	A	306	GLN
1	A	333	GLN
1	B	57	HIS
1	B	68	GLN
1	B	147	GLN
1	B	172	GLN
1	B	256	ASN
1	B	333	GLN
1	B	347	ASN
2	C	508	GLN
2	C	515	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GDP	B	356	-	24,30,30	0.95	1 (4%)	30,47,47	1.32	4 (13%)
5	GOL	A	3	-	5,5,5	0.39	0	5,5,5	0.18	0
4	SO4	A	2	-	4,4,4	0.13	0	6,6,6	0.05	0
3	GDP	A	355	-	24,30,30	0.93	1 (4%)	30,47,47	1.32	4 (13%)
4	SO4	B	1	-	4,4,4	0.14	0	6,6,6	0.05	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GDP	A	355	-	-	4/12/32/32	0/3/3/3
3	GDP	B	356	-	-	4/12/32/32	0/3/3/3
5	GOL	A	3	-	-	2/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	356	GDP	C6-N1	-2.37	1.34	1.37
3	A	355	GDP	C6-N1	-2.30	1.34	1.37

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	356	GDP	PA-O3A-PB	-3.80	119.78	132.83
3	A	355	GDP	PA-O3A-PB	-3.67	120.23	132.83
3	A	355	GDP	C3'-C2'-C1'	3.07	105.59	100.98
3	B	356	GDP	C3'-C2'-C1'	2.81	105.21	100.98
3	A	355	GDP	C8-N7-C5	2.30	107.38	102.99
3	B	356	GDP	C8-N7-C5	2.29	107.34	102.99
3	A	355	GDP	C5-C6-N1	2.27	117.96	113.95
3	B	356	GDP	C5-C6-N1	2.26	117.95	113.95

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	355	GDP	PA-O3A-PB-O2B
3	B	356	GDP	C5'-O5'-PA-O1A

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
3	B	356	GDP	C5'-O5'-PA-O2A
5	A	3	GOL	O1-C1-C2-C3
5	A	3	GOL	O1-C1-C2-O2
3	A	355	GDP	C5'-O5'-PA-O3A
3	A	355	GDP	C5'-O5'-PA-O1A
3	A	355	GDP	C5'-O5'-PA-O2A
3	B	356	GDP	C5'-O5'-PA-O3A
3	B	356	GDP	O4'-C4'-C5'-O5'

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	355	GDP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/328 (96%)	-0.12	9 (2%) 53 43	29, 53, 101, 158	0
1	B	314/328 (95%)	0.23	25 (7%) 12 6	29, 62, 128, 166	0
2	C	33/36 (91%)	0.33	1 (3%) 50 40	29, 73, 108, 127	0
2	D	30/36 (83%)	0.89	9 (30%) 0 0	39, 89, 119, 131	0
All	All	695/728 (95%)	0.10	44 (6%) 20 12	29, 57, 119, 166	0

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	315	ASP	6.6
1	B	257	LYS	6.3
1	B	208	ARG	6.0
1	B	313	ARG	5.9
1	A	314	LYS	5.6
1	B	209	LYS	5.2
1	B	316	THR	5.2
1	B	193	ASP	4.3
1	B	237	ASP	4.2
1	B	63	GLU	3.8
1	A	316	THR	3.7
2	D	498	GLU	3.6
1	A	31	ALA	3.6
1	B	236	GLU	3.6
1	B	258	TRP	3.4
1	B	197	LYS	3.4
1	B	239	GLU	3.4
1	B	312	LYS	3.1
1	B	207	GLU	3.0
1	B	59	ALA	2.9
1	B	315	ASP	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	212	ILE	2.7
2	D	514	ASP	2.7
1	A	97	ASP	2.7
1	B	188	HIS	2.7
2	D	522	GLU	2.6
1	B	202	GLY	2.6
1	A	63	GLU	2.5
1	B	254	CYS	2.5
2	D	500	LEU	2.5
2	D	521	LYS	2.5
1	A	98	SER	2.4
1	A	30	GLY	2.4
1	B	255	ASN	2.4
1	B	210	LYS	2.3
1	B	344	ILE	2.3
1	B	194	LEU	2.3
2	C	514	ASP	2.3
1	A	182	THR	2.2
1	B	314	LYS	2.1
2	D	525	VAL	2.1
2	D	506	ARG	2.1
2	D	502	GLU	2.1
2	D	503	LEU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	A	2	5/5	0.73	0.48	172,176,177,177	0
4	SO4	B	1	5/5	0.83	0.44	156,164,167,170	0
5	GOL	A	3	6/6	0.90	0.19	44,51,52,59	0
3	GDP	B	356	28/28	0.96	0.19	22,47,68,105	0
3	GDP	A	355	28/28	0.97	0.18	17,35,68,84	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.