

Full wwPDB X-ray Structure Validation Report (i)

Sep 13, 2023 – 10:51 AM EDT

PDB ID : 4QIE

Title: Crystal Structure of PduA with edge mutation K26D

Authors: Pang, A.H.; Sawaya, M.R.; Yeates, T.O.

Deposited on : 2014-05-30

Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul: 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13 EDS : 2.35.1

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

 $Refmac \quad : \quad 5.8.0158$

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries,\ resolution\ range(\AA)}) \end{array}$
R_{free}	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain					
1	A	102	5% 68% 18	8% •	14%			
1	В	102	65% 20%	6 •	15%			
1	С	102	75%	9%	17%			
1	D	102		17%	14%			
1	Е	102	75%	9%	16%			



Mol	Chain	Length	Quality of chain				
1	F	102	9%	6% • 14%			
1	G	102	74%	8% 19%			
1	Н	102	9%	7% 13%			
1	I	102	75%	9% 17%			



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5600 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Propanediol utilization protein PduA.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace	
1	A	88	Total	С	N	О	S	0	0	0	
1	A	00	624	390	110	121	3	0	0	0	
1	В	87	Total	С	N	О	S	0	2	0	
1	D	01	630	393	111	123	3	0	<u> </u>	U	
1	С	85	Total	С	N	Ο	S	0	0	0	
1		0.0	593	371	102	117	3	0	U	U	
1	D	88	Total	С	N	Ο	S	0	0	0	
1	D	00	624	390	110	121	3	O	U	U	
1	E	86	Total	С	N	Ο	S	0	1	0	
1	ш	00	611	381	107	120	3	0	1	U	
1	F	88	Total	С	N	O	S	0	0	0	
1	I.	00	616	384	108	121	3	0	U	U	
1	C	G	83	Total	\mathbf{C}	N	Ο	S	0	0	0
1	G	00	583	362	103	115	3	U	U	U	
1	Н	89	Total	С	N	Ο	S	0	0	0	
1	11	09	631	393	111	124	3		<u> </u>	0	
1	I	85	Total	С	N	О	S	0	0	0	
1	1	00	599	374	105	117	3		U	U	

There are 90 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	expression tag	UNP P0A1C7
A	-6	HIS	-	expression tag	UNP P0A1C7
A	-5	HIS	-	expression tag	UNP P0A1C7
A	-4	HIS	-	expression tag	UNP P0A1C7
A	-3	HIS	-	expression tag	UNP P0A1C7
A	-2	HIS	-	expression tag	UNP P0A1C7
A	-1	HIS	-	expression tag	UNP P0A1C7
A	0	GLY	-	expression tag	UNP P0A1C7
A	1	THR	-	expression tag	UNP P0A1C7
A	26	ASP	LYS	engineered mutation	UNP P0A1C7
В	-7	MET	-	expression tag	UNP P0A1C7



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled Modelled	Actual	Comment	Reference
В	-6	HIS	-	expression tag	UNP P0A1C7
В	-5	HIS	-	expression tag	UNP P0A1C7
В	-4	HIS	-	expression tag	UNP P0A1C7
В	-3	HIS	-	expression tag	UNP P0A1C7
В	-2	HIS	-	expression tag	UNP P0A1C7
В	-1	HIS	-	expression tag	UNP P0A1C7
В	0	GLY	-	expression tag	UNP P0A1C7
В	1	THR	-	expression tag	UNP P0A1C7
В	26	ASP	LYS	engineered mutation	UNP P0A1C7
С	-7	MET	-	expression tag	UNP P0A1C7
С	-6	HIS	-	expression tag	UNP P0A1C7
С	-5	HIS	_	expression tag	UNP P0A1C7
С	-4	HIS	-	expression tag	UNP P0A1C7
С	-3	HIS	-	expression tag	UNP P0A1C7
С	-2	HIS	-	expression tag	UNP P0A1C7
С	-1	HIS	_	expression tag	UNP P0A1C7
С	0	GLY	-	expression tag	UNP P0A1C7
С	1	THR	-	expression tag	UNP P0A1C7
С	26	ASP	LYS	engineered mutation	UNP P0A1C7
D	-7	MET	-	expression tag	UNP P0A1C7
D	-6	HIS	-	expression tag	UNP P0A1C7
D	-5	HIS	-	expression tag	UNP P0A1C7
D	-4	HIS	-	expression tag	UNP P0A1C7
D	-3	HIS	-	expression tag	UNP P0A1C7
D	-2	HIS	-	expression tag	UNP P0A1C7
D	-1	HIS	-	expression tag	UNP P0A1C7
D	0	GLY	-	expression tag	UNP P0A1C7
D	1	THR	-	expression tag	UNP P0A1C7
D	26	ASP	LYS	engineered mutation	UNP P0A1C7
Е	-7	MET	-	expression tag	UNP P0A1C7
Е	-6	HIS	-	expression tag	UNP P0A1C7
Е	-5	HIS	-	expression tag	UNP P0A1C7
Е	-4	HIS	-	expression tag	UNP P0A1C7
E	-3	HIS	-	expression tag	UNP P0A1C7
Е	-2	HIS	-	expression tag	UNP P0A1C7
Е	-1	HIS	-	expression tag	UNP P0A1C7
Е	0	GLY	-	expression tag	UNP P0A1C7
Е	1	THR	-	expression tag	UNP P0A1C7
Е	26	ASP	LYS	engineered mutation	UNP P0A1C7
F	-7	MET	-	expression tag	UNP P0A1C7
F	-6	HIS	-	expression tag	UNP P0A1C7
F	-5	HIS	-	expression tag	UNP P0A1C7



 $Continued\ from\ previous\ page...$

Chain	Residue	Modelled	Actual	Comment	Reference
F	-4	HIS	-	expression tag	UNP P0A1C7
F	-3	HIS	-	expression tag	UNP P0A1C7
F	-2	HIS	-	expression tag	UNP P0A1C7
F	-1	HIS	-	expression tag	UNP P0A1C7
F	0	GLY	-	expression tag	UNP P0A1C7
F	1	THR	-	expression tag	UNP P0A1C7
F	26	ASP	LYS	engineered mutation	UNP P0A1C7
G	-7	MET	-	expression tag	UNP P0A1C7
G	-6	HIS	-	expression tag	UNP P0A1C7
G	-5	HIS	-	expression tag	UNP P0A1C7
G	-4	HIS	-	expression tag	UNP P0A1C7
G	-3	HIS	-	expression tag	UNP P0A1C7
G	-2	HIS	_	expression tag	UNP P0A1C7
G	-1	HIS	-	expression tag	UNP P0A1C7
G	0	GLY	_	expression tag	UNP P0A1C7
G	1	THR	-	expression tag	UNP P0A1C7
G	26	ASP	LYS	engineered mutation	UNP P0A1C7
Н	-7	MET	-	expression tag	UNP P0A1C7
Н	-6	HIS	-	expression tag	UNP P0A1C7
Н	-5	HIS	-	expression tag	UNP P0A1C7
Н	-4	HIS	-	expression tag	UNP P0A1C7
Н	-3	HIS	-	expression tag	UNP P0A1C7
Н	-2	HIS	-	expression tag	UNP P0A1C7
Н	-1	HIS	-	expression tag	UNP P0A1C7
Н	0	GLY	_	expression tag	UNP P0A1C7
Н	1	THR	-	expression tag	UNP P0A1C7
Н	26	ASP	LYS	engineered mutation	UNP P0A1C7
I	-7	MET	-	expression tag	UNP P0A1C7
I	-6	HIS	-	expression tag	UNP P0A1C7
I	-5	HIS	-	expression tag	UNP P0A1C7
I	-4	HIS	-	expression tag	UNP P0A1C7
I	-3	HIS	-	expression tag	UNP P0A1C7
I	-2	HIS	-	expression tag	UNP P0A1C7
I	-1	HIS	-	expression tag	UNP P0A1C7
I	0	GLY	-	expression tag	UNP P0A1C7
I	1	THR	-	expression tag	UNP P0A1C7
I	26	ASP	LYS	engineered mutation	UNP P0A1C7


 \bullet Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: $\mathrm{C_3H_8O_3}).$





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	G	1	Total C O 6 3 3	0	0

 \bullet Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: $\mathrm{O_4S}).$



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total O S 5 4 1	0	0
3	D	1	Total O S 5 4 1	0	0



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	Н	1	Total O S 5 4 1	0	0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	7	Total O 7 7	0	0
4	В	13	Total O 13 13	0	0
4	С	2	Total O 2 2	0	0
4	D	6	Total O 6 6	0	0
4	Е	6	Total O 6 6	0	0
4	F	10	Total O 10 10	0	0
4	G	5	Total O 5 5	0	0
4	Н	4	Total O 4 4	0	0
4	I	9	Total O 9 9	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Propanediol utilization protein PduA



• Molecule 1: Propanediol utilization protein PduA







4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants	108.11Å 108.11Å 334.34Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.85 - 2.35	Depositor
rtesolution (A)	19.85 - 2.35	EDS
% Data completeness	94.6 (19.85-2.35)	Depositor
(in resolution range)	94.6 (19.85-2.35)	EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.47 (at 2.35Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1555)	Depositor
R, R_{free}	0.201 , 0.238	Depositor
It, It free	0.207 , 0.240	DCC
R_{free} test set	4645 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.014	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.37, 38.8	EDS
L-test for twinning ²	$< L > = 0.48, < L^2> = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5600	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 11.95% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond	\mathbf{angles}
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.46	0/629	0.64	0/853
1	В	0.44	0/635	0.63	0/861
1	С	0.41	0/597	0.64	0/811
1	D	0.44	0/629	0.62	0/853
1	Е	0.44	0/618	0.64	0/838
1	F	0.51	0/621	0.68	0/845
1	G	0.42	0/587	0.61	0/796
1	Н	0.42	0/636	0.65	0/864
1	I	0.46	0/603	0.63	0/818
All	All	0.45	0/5555	0.64	0/7539

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	624	0	651	18	0
1	В	630	0	654	18	0
1	С	593	0	612	5	0
1	D	624	0	651	12	0



Continued from previous page...

Mol	Chain		H(model)	H(added)	Clashes	Symm-Clashes
1	Ε	611	0	636	9	0
1	F	616	0	629	14	0
1	G	583	0	601	6	0
1	Н	631	0	656	4	0
1	I	599	0	623	9	0
2	A	6	0	8	1	0
2	G	6	0	8	0	0
3	A	5	0	0	0	0
3	D	5	0	0	0	0
3	Н	5	0	0	0	0
4	A	7	0	0	0	0
4	В	13	0	0	1	0
4	С	2	0	0	0	0
4	D	6	0	0	0	0
4	${ m E}$	6	0	0	0	0
4	F	10	0	0	0	0
4	G	5	0	0	0	0
4	Н	4	0	0	0	0
4	I	9	0	0	0	0
All	All	5600	0	5729	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:B:35:TYR:CD1	1:F:87:ILE:HG12	2.21	0.76
1:B:79:ARG:HD3	1:G:66:ARG:HB3	1.74	0.70
1:F:6:LEU:HD11	1:F:88:LEU:HD11	1.77	0.67
1:A:90:LYS:HE3	1:F:37:LYS:NZ	2.15	0.61
1:G:35:TYR:CD2	1:I:87:ILE:HB	2.36	0.60
1:A:5:ALA:HB2	1:A:51:VAL:HG22	1.83	0.59
1:A:36:GLU:OE2	1:A:90:LYS:HD2	2.02	0.59
1:G:48:ARG:NH1	1:G:85:GLU:OE1	2.36	0.59
1:B:26:ASP:HA	1:F:81:HIS:HE1	1.68	0.58
1:A:79:ARG:HB3	1:A:80:PRO:HD2	1.88	0.56
1:A:21:ALA:HB2	1:A:45:VAL:HG11	1.88	0.56
1:B:3:GLN:NE2	1:B:80:PRO:O	2.34	0.55
1:B:12[B]:LYS:HB3	1:B:70:GLU:HB2	1.89	0.54
2:A:101:GOL:H32	1:D:40:SER:OG	2.07	0.54



Continuea from pred	1 0	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}\ (\rm \mathring{A})$	overlap (Å)
1:I:33:VAL:CG2	1:I:48:ARG:HG2	2.37	0.54
1:F:33:VAL:HG21	1:F:48:ARG:HG3	1.89	0.54
1:E:23:ALA:O	1:E:26:ASP:HB2	2.07	0.54
1:F:6:LEU:HD21	1:F:88:LEU:HD13	1.89	0.54
1:B:26:ASP:HA	1:F:81:HIS:CE1	2.43	0.53
1:G:35:TYR:HD2	1:I:87:ILE:HB	1.74	0.53
1:D:21:ALA:HB2	1:D:45:VAL:HG11	1.90	0.52
1:D:79:ARG:HB3	1:D:80:PRO:HD2	1.92	0.52
1:F:37:LYS:N	1:F:37:LYS:HD2	2.26	0.51
1:I:20:ALA:HB1	1:I:24:MET:HE2	1.93	0.50
1:A:35:TYR:HE1	1:E:87:ILE:HA	1.77	0.50
1:G:24:MET:HG2	1:G:57:ALA:O	2.12	0.50
1:B:87:ILE:HB	1:D:35:TYR:CD1	2.46	0.50
1:B:33:VAL:CG2	1:B:48:ARG:HG3	2.41	0.49
1:B:33:VAL:HG21	1:B:48:ARG:HG3	1.95	0.49
1:F:33:VAL:CG2	1:F:48:ARG:HG3	2.42	0.49
1:B:20:ALA:HB2	1:B:65:ALA:HB2	1.93	0.48
1:A:81:HIS:ND1	1:A:81:HIS:N	2.60	0.48
1:A:3:GLN:O	1:A:51:VAL:HG23	2.13	0.48
1:B:24:MET:HG2	1:B:57:ALA:O	2.13	0.48
1:H:8:MET:HG2	1:H:46:ILE:HG12	1.95	0.47
1:I:33:VAL:HG21	1:I:48:ARG:HG2	1.94	0.47
1:A:90:LYS:HE3	1:F:37:LYS:HZ1	1.77	0.47
1:A:27:SER:C	1:A:29:ASN:H	2.17	0.46
1:I:25:VAL:HG11	1:I:32:LEU:HB2	1.97	0.46
1:B:35:TYR:CE1	1:F:87:ILE:HG12	2.51	0.46
1:A:41:GLY:HA2	1:E:38:ILE:HG13	1.99	0.45
1:A:27:SER:O	1:A:29:ASN:N	2.50	0.45
1:A:35:TYR:CD1	1:E:87:ILE:HG12	2.51	0.45
1:B:87:ILE:HB	1:D:35:TYR:CE1	2.52	0.45
1:A:90:LYS:HE3	1:F:37:LYS:HZ3	1.82	0.45
1:A:35:TYR:CE1	1:E:87:ILE:HA	2.52	0.44
1:A:30:VAL:HG13	1:A:47:VAL:HB	1.99	0.44
1:F:87:ILE:HG22	1:F:88:LEU:HG	2.01	0.43
1:A:24:MET:HG2	1:A:57:ALA:O	2.19	0.43
1:C:33:VAL:HG13	1:C:46:ILE:O	2.18	0.43
1:B:38:ILE:HG22	1:D:37:LYS:HD2	2.01	0.42
1:H:20:ALA:HB2	1:H:65:ALA:HB2	2.01	0.42
1:C:38:ILE:HG13	1:E:41:GLY:HA2	2.01	0.42
1:D:5:ALA:HB2	1:D:51:VAL:HG22	2.01	0.42
1:H:24:MET:HG2	1:H:57:ALA:O	2.20	0.42



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)
1:B:37:LYS:HB3	4:B:103:HOH:O	2.19	0.42
1:I:21:ALA:O	1:I:25:VAL:HG22	2.19	0.42
1:D:3:GLN:O	1:D:51:VAL:HG23	2.19	0.42
1:D:24:MET:HG2	1:D:57:ALA:O	2.20	0.41
1:D:28:ALA:HB3	1:D:57:ALA:HB2	2.01	0.41
1:F:81:HIS:N	1:F:81:HIS:CD2	2.86	0.41
1:B:15:THR:HG22	1:B:68:VAL:HG11	2.02	0.41
1:H:7:GLY:O	1:H:8:MET:HG3	2.20	0.41
1:B:3:GLN:HG3	1:B:4:GLU:N	2.34	0.41
1:I:20:ALA:HB2	1:I:65:ALA:HB2	2.01	0.41
1:C:84:VAL:O	1:C:87:ILE:HG12	2.20	0.41
1:C:20:ALA:O	1:C:24:MET:HG3	2.21	0.41
1:A:35:TYR:CE1	1:E:87:ILE:HG23	2.56	0.41
1:B:84:VAL:O	1:B:87:ILE:HG12	2.21	0.41
1:C:8:MET:HE2	1:E:18:ILE:HD13	2.02	0.41
1:D:8:MET:HG2	1:D:46:ILE:HG12	2.03	0.41
1:E:8:MET:HG2	1:E:46:ILE:HG12	2.03	0.41
1:G:35:TYR:CE2	1:I:87:ILE:HB	2.56	0.41
1:D:20:ALA:HB2	1:D:65:ALA:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	A	86/102 (84%)	85 (99%)	0	1 (1%)	13	11
1	В	87/102 (85%)	85 (98%)	2 (2%)	0	100	100
1	С	83/102 (81%)	82 (99%)	1 (1%)	0	100	100
1	D	86/102 (84%)	85 (99%)	1 (1%)	0	100	100
1	E	85/102 (83%)	84 (99%)	1 (1%)	0	100	100



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	F	86/102 (84%)	84 (98%)	2 (2%)	0	100	100
1	G	81/102 (79%)	81 (100%)	0	0	100	100
1	Н	87/102 (85%)	85 (98%)	2 (2%)	0	100	100
1	I	83/102 (81%)	82 (99%)	1 (1%)	0	100	100
All	All	$764/918 \; (83\%)$	753 (99%)	10 (1%)	1 (0%)	51	63

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	28	ALA

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	63/75 (84%)	62 (98%)	1 (2%)	62 75
1	В	64/75 (85%)	62 (97%)	2 (3%)	40 48
1	С	59/75 (79%)	58 (98%)	1 (2%)	60 72
1	D	63/75 (84%)	63 (100%)	0	100 100
1	E	62/75 (83%)	60 (97%)	2 (3%)	39 47
1	F	61/75 (81%)	60 (98%)	1 (2%)	62 75
1	G	58/75 (77%)	56 (97%)	2 (3%)	37 46
1	Н	64/75 (85%)	64 (100%)	0	100 100
1	I	60/75 (80%)	60 (100%)	0	100 100
All	All	554/675~(82%)	545 (98%)	9 (2%)	65 75

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	81	HIS
1	В	26	ASP



Continued from previous page...

Mol	Chain	Res	Type
1	В	85	GLU
1	С	83	ASP
1	Е	27[A]	SER
1	Е	27[B]	SER
1	F	87	ILE
1	G	82	THR
1	G	83	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	81	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Type Chain		Res Link	B	Bond lengths			Bond angles		
MIOI	Туре	Chain	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2	
3	SO4	D	101	-	4,4,4	0.16	0	6,6,6	0.14	0	



Mol	Type	Chain	Res	Res Link	Bond lengths			Bond angles		
MIOI					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SO4	Н	101	-	4,4,4	0.19	0	6,6,6	0.18	0
2	GOL	G	101	-	5,5,5	0.39	0	5,5,5	0.24	0
3	SO4	A	102	-	4,4,4	0.15	0	6,6,6	0.14	0
2	GOL	A	101	-	5,5,5	0.41	0	5,5,5	0.98	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	101	-	-	2/4/4/4	-
2	GOL	G	101	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	101	GOL	O1-C1-C2-C3
2	G	101	GOL	O1-C1-C2-C3
2	A	101	GOL	O1-C1-C2-O2
2	G	101	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

\mathbf{Mol}	Chain	Res	Type	Clashes	Symm-Clashes
2	A	101	GOL	1	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	$\# \mathrm{RSRZ}{>}2$	$OWAB(A^2)$	Q < 0.9
1	A	88/102 (86%)	0.22	5 (5%) 23 34	27, 38, 63, 96	0
1	В	87/102 (85%)	0.15	5 (5%) 23 34	27, 34, 69, 77	0
1	C	85/102 (83%)	0.72	10 (11%) 4 7	31, 52, 70, 82	0
1	D	88/102 (86%)	0.47	6 (6%) 17 25	29, 43, 79, 98	0
1	E	86/102 (84%)	0.17	3 (3%) 44 56	28, 37, 62, 72	0
1	F	88/102 (86%)	0.37	9 (10%) 6 11	27, 39, 81, 85	0
1	G	83/102 (81%)	0.28	5 (6%) 21 32	28, 39, 68, 82	0
1	Н	89/102 (87%)	0.36	9 (10%) 7 11	31, 38, 58, 78	0
1	I	85/102 (83%)	0.16	6 (7%) 16 23	28, 37, 67, 80	0
All	All	779/918 (84%)	0.32	58 (7%) 14 22	27, 39, 71, 98	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res Type		RSRZ
1	F	3	GLN	6.1
1	D	81	HIS	5.9
1	D	82	THR	5.5
1	G	82	THR	5.3
1	F	89	PRO	4.9
1	A	81	HIS	4.7
1	С	82	THR	4.7
1	F	82	THR	4.6
1	С	86	LYS	4.5
1	Н	1	THR	4.1
1	С	27	SER	4.0
1	D	80	PRO	4.0
1	G	83	ASP	3.9
1	I	86	LYS	3.9
1	D	3	GLN	3.8



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	В	3	GLN	3.6
1	С	31	MET	3.6
1	В	86	LYS	3.4
1	F	83	ASP	3.4
1	I	26	ASP	3.4
1	A	82	THR	3.4
1	D	83	ASP	3.0
1	В	35	TYR	3.0
1	Н	83	ASP	3.0
1	F	88	LEU	3.0
1	Е	82	THR	3.0
1	С	21	ALA	2.9
1	F	79	ARG	2.9
1	I	27	SER	2.8
1	G	79	ARG	2.8
1	С	67	ASN	2.8
1	A	35	TYR	2.7
1	A	80	PRO	2.7
1	Е	28	ALA	2.7
1	F	90	LYS	2.5
1	D	86	LYS	2.5
1	Н	2	GLN	2.5
1	I	31	MET	2.4
1	G	81	HIS	2.4
1	С	18	ILE	2.3
1	Н	29	ASN	2.3
1	Н	45	VAL	2.3
1	В	28	ALA	2.3
1	С	17	ALA	2.3
1	A	83	ASP	2.2
1	С	69	GLY	2.2
1	Н	6	LEU	2.2
1	Н	81	HIS	2.1
1	С	9	VAL	2.1
1	F	86	LYS	2.1
1	I	82	THR	2.1
1	В	82	THR	2.1
1	Н	20	ALA	2.1
1	G	50	ASP	2.0
1	Е	47	VAL	2.0
1	I	9	VAL	2.0
1	Н	31	MET	2.0



		-		
Mol	Chain	Res	Type	RSRZ
1	F	27	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q < 0.9
3	SO4	Н	101	5/5	0.75	0.29	70,72,94,102	0
3	SO4	D	101	5/5	0.82	0.36	87,95,106,108	0
2	GOL	A	101	6/6	0.89	0.20	32,35,41,51	0
3	SO4	A	102	5/5	0.92	0.30	81,87,96,97	0
2	GOL	G	101	6/6	0.94	0.15	37,39,41,43	6

6.5 Other polymers (i)

There are no such residues in this entry.

