



# Full wwPDB X-ray Structure Validation Report i

Feb 27, 2024 – 11:37 PM EST

PDB ID : 5QJ6  
Title : PanDDA analysis group deposition of models with modelled events (e.g. bound ligands) – Crystal Structure of NUDT5 in complex with Z1614545742  
Authors : Dubianok, Y.; Collins, P.; Krojer, T.; Wright, N.; Strain-Damerell, C.; Burgess-Brown, N.; Bountra, C.; Arrowsmith, C.H.; Edwards, A.; Huber, K.; von Delft, F.  
Deposited on : 2018-10-31  
Resolution : 1.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The types of validation reports are described at  
<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriaage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GQJ	D	303	-	-	-	X

## 2 Entry composition [\(i\)](#)

There are 5 unique types of molecules in this entry. The entry contains 6052 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-sugar pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	194	Total	C 1450	N 917	O 242	S 283	8	0	0
1	B	194	Total	C 1464	N 924	O 244	S 288	8	0	0
1	C	192	Total	C 1402	N 884	O 234	S 277	7	0	0
1	D	190	Total	C 1410	N 890	O 238	S 274	8	0	1


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP Q9UKK9
B	0	SER	-	expression tag	UNP Q9UKK9
C	0	SER	-	expression tag	UNP Q9UKK9
D	0	SER	-	expression tag	UNP Q9UKK9

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Mg 2	0	0
2	B	2	Total	Mg 2	0	0
2	C	2	Total	Mg 2	0	0
2	D	2	Total	Mg 2	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

- Molecule 4 is methyl (2 {S},4 {R})-1-(furan-2-ylcarbonyl)-4-oxidanyl-pyrrolidine-2-carboxylate (three-letter code: GQJ) (formula: C<sub>11</sub>H<sub>13</sub>NO<sub>5</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	N	O	
			17	11	1	5	0
4	D	1	Total	C	N	O	
			17	11	1	5	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	70	Total	O		
			70	70	0	0
5	B	79	Total	O		
			79	79	0	0
5	C	74	Total	O		
			74	74	0	0
5	D	37	Total	O		
			37	37	0	0


### 3 Residue-property plots [\(i\)](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase




- Molecule 1: ADP-sugar pyrophosphatase



- Molecule 1: ADP-sugar pyrophosphatase





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.25 Å    59.78 Å    80.09 Å 79.42°    81.68°    75.64°	Depositor
Resolution (Å)	78.30 – 1.65 31.87 – 1.65	Depositor EDS
% Data completeness (in resolution range)	96.2 (78.30-1.65) 96.3 (31.87-1.65)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	1.56 (at 1.65 Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
$R$ , $R_{free}$	0.212 , 0.249 0.228 , 0.262	Depositor DCC
$R_{free}$ test set	5059 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	26.8	Xtriage
Anisotropy	0.166	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 41.0	EDS
L-test for twinning <sup>2</sup>	$<  L  > = 0.50$ , $< L^2 > = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	6052	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.61% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GQJ, EDO, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.02	1/1478 (0.1%)	1.02	2/2019 (0.1%)
1	B	0.94	0/1492	1.03	4/2035 (0.2%)
1	C	0.96	0/1429	1.01	2/1958 (0.1%)
1	D	0.96	1/1436 (0.1%)	1.11	9/1961 (0.5%)
All	All	0.97	2/5835 (0.0%)	1.04	17/7973 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	112	GLU	CD-OE2	-5.59	1.19	1.25
1	D	199	SER	CB-OG	-5.02	1.35	1.42

All (17) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	196[A]	ARG	NE-CZ-NH2	-11.06	114.77	120.30
1	D	196[B]	ARG	NE-CZ-NH2	-11.06	114.77	120.30
1	D	196[A]	ARG	NE-CZ-NH1	9.20	124.90	120.30
1	D	196[B]	ARG	NE-CZ-NH1	9.20	124.90	120.30
1	A	70	ARG	NE-CZ-NH2	-8.85	115.88	120.30
1	C	196	ARG	NE-CZ-NH1	8.50	124.55	120.30
1	B	196	ARG	NE-CZ-NH1	7.50	124.05	120.30
1	D	133	ASP	CB-CG-OD1	6.66	124.29	118.30
1	A	54	ARG	NE-CZ-NH2	-6.30	117.15	120.30
1	B	111	ARG	NE-CZ-NH1	5.92	123.26	120.30
1	D	194	ASP	CB-CG-OD1	-5.58	113.28	118.30
1	D	38	ASP	CB-CG-OD1	5.43	123.19	118.30
1	B	194	ASP	CB-CG-OD2	-5.38	113.46	118.30
1	B	151	ASP	CB-CG-OD2	5.18	122.96	118.30
1	C	38	ASP	CB-CG-OD1	5.12	122.91	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	132	MET	CG-SD-CE	-5.06	92.11	100.20
1	D	74	TYR	CA-CB-CG	5.05	123.00	113.40

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1450	0	1405	13	0
1	B	1464	0	1414	11	0
1	C	1402	0	1298	9	0
1	D	1410	0	1349	32	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
2	C	2	0	0	0	0
2	D	2	0	0	0	0
3	A	4	0	6	0	0
3	B	8	0	12	2	0
3	C	8	0	12	0	0
3	D	4	0	6	0	0
4	B	17	0	0	0	0
4	D	17	0	0	0	0
5	A	70	0	0	0	0
5	B	79	0	0	4	0
5	C	74	0	0	4	0
5	D	37	0	0	0	0
All	All	6052	0	5502	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:196[B]:ARG:HH21	1:D:196[B]:ARG:CG	1.31	1.33
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:HG3	1.31	1.13
1:C:176:ASN:HB3	5:C:463:HOH:O	1.74	0.88
1:B:208:ASN:OD1	5:B:401:HOH:O	1.97	0.82
1:C:125:GLU:OE1	5:C:401:HOH:O	1.97	0.82
1:A:71:THR:HG23	1:A:151:ASP:OD2	1.79	0.81
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:N	2.30	0.80
1:B:92:ILE:HD11	1:B:191:LEU:HD13	1.65	0.78
1:D:120:LYS:H	1:D:155:ASN:HD21	1.33	0.76
1:C:104:THR:HG23	1:C:107:ALA:H	1.51	0.75
1:D:196[B]:ARG:CG	1:D:196[B]:ARG:NH2	2.06	0.72
1:D:194:ASP:OD1	1:D:196[A]:ARG:HD3	1.92	0.70
1:D:196[B]:ARG:HH21	1:D:196[B]:ARG:CB	2.07	0.67
1:B:104:THR:HG23	1:B:107:ALA:H	1.58	0.67
1:A:203:ALA:HB3	1:B:203:ALA:HB3	1.80	0.63
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:CB	2.62	0.62
1:D:196[B]:ARG:HH21	1:D:196[B]:ARG:HG3	0.50	0.62
1:C:203:ALA:HB3	1:D:203:ALA:HB3	1.82	0.60
1:A:104:THR:HG23	1:A:107:ALA:H	1.66	0.59
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:CA	2.67	0.57
1:D:208:ASN:ND2	1:D:208:ASN:C	2.57	0.57
1:C:200:TYR:OH	1:D:206:HIS:HD2	1.89	0.55
5:C:401:HOH:O	1:D:206:HIS:HE1	1.89	0.55
1:D:196[B]:ARG:NH2	1:D:196[B]:ARG:H	2.03	0.55
1:C:203:ALA:HB2	1:D:200:TYR:CD1	2.42	0.54
1:C:15:GLN:N	5:C:403:HOH:O	2.41	0.54
1:B:189:GLU:HB3	1:B:191:LEU:HD12	1.91	0.52
3:B:301:EDO:C2	5:B:409:HOH:O	2.56	0.52
1:D:207:ALA:O	1:D:208:ASN:ND2	2.38	0.51
1:A:125:GLU:HG3	1:A:145:THR:HB	1.92	0.51
1:A:125:GLU:OE1	1:B:206:HIS:HE1	1.93	0.51
1:B:125:GLU:HG2	1:B:145:THR:HB	1.93	0.51
1:D:71:THR:O	1:D:73:HIS:ND1	2.37	0.51
1:D:113:LEU:HD22	1:D:146:VAL:HG11	1.93	0.51
1:D:104:THR:HG23	1:D:107:ALA:H	1.77	0.50
1:A:200:TYR:CD1	1:B:203:ALA:HB2	2.46	0.50
1:A:54:ARG:HD3	1:A:60:ASP:OD1	2.11	0.50
1:B:125:GLU:HG3	5:B:470:HOH:O	2.12	0.50
1:A:200:TYR:OH	1:B:206:HIS:HD2	1.96	0.49
1:A:191:LEU:C	1:A:191:LEU:HD23	2.36	0.47
1:D:191:LEU:C	1:D:191:LEU:HD12	2.35	0.46
1:A:104:THR:CG2	1:A:107:ALA:H	2.28	0.46

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:38:ASP:OD1	1:D:40:THR:N	2.49	0.45
1:D:119:TYR:CD1	1:D:148:ILE:HG21	2.52	0.45
1:D:113:LEU:CD2	1:D:146:VAL:HG11	2.47	0.44
1:D:196[B]:ARG:N	1:D:196[B]:ARG:HH22	2.12	0.43
1:C:179:LEU:HD11	1:C:198:TYR:CZ	2.53	0.43
1:A:64:VAL:O	1:A:66:PRO:HD3	2.19	0.43
3:B:301:EDO:H22	5:B:409:HOH:O	2.18	0.42
1:D:208:ASN:C	1:D:208:ASN:HD22	2.23	0.42
1:B:14:LYS:HE3	1:B:39:PRO:HB2	2.02	0.42
1:D:64:VAL:O	1:D:66:PRO:HD3	2.20	0.42
1:A:71:THR:O	1:A:73:HIS:HD2	2.03	0.41
1:D:195:ALA:HB3	1:D:196[B]:ARG:HH12	1.85	0.41
1:C:200:TYR:CD1	1:D:203:ALA:HB2	2.56	0.41
1:D:67:VAL:O	1:D:67:VAL:HG13	2.21	0.41
1:D:68:LEU:HA	1:D:148:ILE:HB	2.03	0.40
1:D:38:ASP:HB2	1:D:39:PRO:CD	2.52	0.40
1:A:65:ILE:HA	1:A:66:PRO:HD3	1.88	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	192/209 (92%)	188 (98%)	3 (2%)	1 (0%)	29 11
1	B	190/209 (91%)	186 (98%)	2 (1%)	2 (1%)	14 2
1	C	188/209 (90%)	184 (98%)	3 (2%)	1 (0%)	29 11
1	D	187/209 (90%)	183 (98%)	3 (2%)	1 (0%)	29 11
All	All	757/836 (91%)	741 (98%)	11 (2%)	5 (1%)	22 6

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	66	PRO
1	A	66	PRO
1	C	66	PRO
1	D	66	PRO
1	B	161	LYS

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	151/179 (84%)	146 (97%)	5 (3%)	38 12
1	B	154/179 (86%)	151 (98%)	3 (2%)	57 34
1	C	139/179 (78%)	135 (97%)	4 (3%)	42 16
1	D	144/179 (80%)	140 (97%)	4 (3%)	43 18
All	All	588/716 (82%)	572 (97%)	16 (3%)	44 19

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	LYS
1	A	38	ASP
1	A	43	THR
1	A	125	GLU
1	A	191	LEU
1	B	31	LEU
1	B	38	ASP
1	B	43	THR
1	C	31	LEU
1	C	38	ASP
1	C	71	THR
1	C	164	ASP
1	D	24	SER
1	D	43	THR
1	D	72	LEU
1	D	208	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	73	HIS
1	A	180	GLN
1	A	206	HIS
1	B	73	HIS
1	B	190	HIS
1	B	206	HIS
1	D	155	ASN
1	D	206	HIS

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [\(i\)](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	B	305	-	3,3,3	0.49	0	2,2,2	0.22	0
3	EDO	D	304	-	3,3,3	0.53	0	2,2,2	0.36	0
3	EDO	C	301	-	3,3,3	0.65	0	2,2,2	0.33	0
3	EDO	A	303	-	3,3,3	0.49	0	2,2,2	0.63	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	GQJ	D	303	-	14,18,18	1.81	5 (35%)	19,25,25	2.04	4 (21%)
4	GQJ	B	304	-	14,18,18	1.50	3 (21%)	19,25,25	2.19	7 (36%)
3	EDO	C	304	-	3,3,3	0.53	0	2,2,2	0.47	0
3	EDO	B	301	-	3,3,3	0.78	0	2,2,2	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	305	-	-	0/1/1/1	-
3	EDO	D	304	-	-	0/1/1/1	-
3	EDO	C	301	-	-	1/1/1/1	-
3	EDO	A	303	-	-	0/1/1/1	-
4	GQJ	D	303	-	-	2/10/26/26	0/2/2/2
4	GQJ	B	304	-	-	2/10/26/26	0/2/2/2
3	EDO	C	304	-	-	0/1/1/1	-
3	EDO	B	301	-	-	0/1/1/1	-

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	D	303	GQJ	C11-C10	-3.11	1.45	1.49
4	B	304	GQJ	O02-C03	2.85	1.40	1.33
4	D	303	GQJ	C04-C03	2.84	1.57	1.52
4	D	303	GQJ	C10-N09	2.77	1.41	1.34
4	D	303	GQJ	O02-C03	2.59	1.39	1.33
4	B	304	GQJ	C11-C10	-2.27	1.46	1.49
4	D	303	GQJ	C05-C06	2.20	1.56	1.52
4	B	304	GQJ	C10-N09	2.17	1.39	1.34

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	304	GQJ	C01-O02-C03	5.43	128.21	115.94
4	D	303	GQJ	C01-O02-C03	4.86	126.92	115.94
4	B	304	GQJ	O02-C03-C04	4.36	118.86	110.89
4	D	303	GQJ	O02-C03-C04	4.16	118.48	110.89
4	D	303	GQJ	O02-C03-O17	-3.79	116.42	123.84
4	B	304	GQJ	O02-C03-O17	-2.93	118.10	123.84

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	304	GQJ	C06-C08-N09	2.63	105.87	103.08
4	B	304	GQJ	C06-C05-C04	2.54	107.22	103.97
4	B	304	GQJ	C05-C06-C08	2.26	106.09	103.31
4	B	304	GQJ	C05-C04-N09	2.24	106.06	103.10
4	D	303	GQJ	O07-C06-C08	-2.22	105.53	110.39

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	B	304	GQJ	O17-C03-O02-C01
4	D	303	GQJ	O17-C03-O02-C01
4	B	304	GQJ	C04-C03-O02-C01
4	D	303	GQJ	C04-C03-O02-C01
3	C	301	EDO	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	301	EDO	2	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	194/209 (92%)	0.19	14 (7%) <span style="background-color: red; border: 1px solid black; padding: 2px;">15</span> <span style="background-color: red; border: 1px solid black; padding: 2px;">14</span>	19, 29, 54, 65	5 (2%)
1	B	194/209 (92%)	0.45	21 (10%) <span style="background-color: red; border: 1px solid black; padding: 2px;">5</span> <span style="background-color: red; border: 1px solid black; padding: 2px;">5</span>	20, 33, 59, 73	10 (5%)
1	C	192/209 (91%)	0.34	20 (10%) <span style="background-color: red; border: 1px solid black; padding: 2px;">6</span> <span style="background-color: red; border: 1px solid black; padding: 2px;">5</span>	20, 34, 58, 67	1 (0%)
1	D	190/209 (90%)	0.77	29 (15%) <span style="background-color: red; border: 1px solid black; padding: 2px;">2</span> <span style="background-color: red; border: 1px solid black; padding: 2px;">1</span>	22, 37, 60, 72	15 (7%)
All	All	770/836 (92%)	0.44	84 (10%) <span style="background-color: red; border: 1px solid black; padding: 2px;">5</span> <span style="background-color: red; border: 1px solid black; padding: 2px;">4</span>	19, 33, 59, 73	31 (4%)

All (84) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	196[A]	ARG	16.3
1	D	207	ALA	8.9
1	B	67	VAL	8.6
1	B	66	PRO	8.0
1	A	204	LEU	8.0
1	B	68	LEU	7.1
1	D	145	THR	6.9
1	D	67	VAL	6.5
1	D	208	ASN	5.8
1	B	145	THR	5.8
1	D	66	PRO	5.6
1	B	127	SER	5.5
1	D	40	THR	5.5
1	B	126	CYS	5.3
1	D	153	ALA	5.1
1	C	40	THR	4.9
1	C	18	ILE	4.8
1	A	206	HIS	4.7
1	B	157	ARG	4.6
1	D	146	VAL	4.5
1	B	102	GLY	4.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	C	16	TYR	4.2
1	B	163	GLY	4.2
1	D	68	LEU	4.2
1	B	190	HIS	4.0
1	C	22	LEU	4.0
1	A	163	GLY	3.9
1	B	69	GLN	3.9
1	B	164	ASP	3.8
1	A	167	PHE	3.7
1	A	205	LYS	3.6
1	D	158	PRO	3.6
1	D	157	ARG	3.6
1	C	102	GLY	3.5
1	D	148	ILE	3.4
1	D	28	TRP	3.4
1	C	101	ASP	3.3
1	A	164	ASP	3.2
1	B	165	GLY	3.2
1	D	16	TYR	3.2
1	C	41	GLY	3.1
1	C	206	HIS	3.1
1	B	175	LYS	3.0
1	D	147	THR	3.0
1	A	195	ALA	3.0
1	D	152	ASP	3.0
1	D	156	ALA	2.9
1	B	162	PRO	2.9
1	A	190	HIS	2.9
1	B	167	PHE	2.7
1	A	56	GLU	2.7
1	D	51	ARG	2.6
1	D	101	ASP	2.6
1	C	197	VAL	2.6
1	C	100	ASP	2.6
1	A	157	ARG	2.6
1	C	195	ALA	2.6
1	D	125	GLU	2.6
1	D	160	PRO	2.6
1	B	159	LYS	2.5
1	B	156	ALA	2.4
1	D	100	ASP	2.4
1	C	164	ASP	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	58	THR	2.4
1	B	158	PRO	2.3
1	B	195	ALA	2.3
1	D	167	PHE	2.3
1	D	159	LYS	2.3
1	C	39	PRO	2.2
1	A	22	LEU	2.2
1	D	121	GLY	2.2
1	D	74	TYR	2.2
1	A	162	PRO	2.2
1	A	156	ALA	2.2
1	C	36	TYR	2.2
1	C	15	GLN	2.2
1	C	190	HIS	2.1
1	C	21	GLU	2.1
1	C	19	SER	2.1
1	D	102	GLY	2.1
1	C	104	THR	2.1
1	B	28	TRP	2.1
1	D	195	ALA	2.1
1	C	53	THR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GQJ	D	303	17/17	0.57	0.45	50,53,59,60	17
4	GQJ	B	304	17/17	0.63	0.39	37,43,48,53	17

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	B	303	1/1	0.76	0.18	105,105,105,105	0
2	MG	C	302	1/1	0.80	0.08	65,65,65,65	0
3	EDO	D	304	4/4	0.89	0.15	35,39,40,41	0
2	MG	A	302	1/1	0.91	0.11	60,60,60,60	0
2	MG	D	301	1/1	0.91	0.04	33,33,33,33	0
3	EDO	B	301	4/4	0.91	0.20	42,43,45,48	0
3	EDO	C	304	4/4	0.92	0.24	40,40,41,43	0
3	EDO	A	303	4/4	0.93	0.17	28,28,30,30	0
3	EDO	B	305	4/4	0.93	0.16	32,35,36,36	0
3	EDO	C	301	4/4	0.94	0.07	36,51,53,62	0
2	MG	D	302	1/1	0.96	0.11	38,38,38,38	0
2	MG	C	303	1/1	0.97	0.03	31,31,31,31	0
2	MG	B	302	1/1	0.99	0.04	32,32,32,32	0
2	MG	A	301	1/1	0.99	0.10	27,27,27,27	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.