



## wwPDB EM Validation Summary Report ⓘ

Jan 29, 2022 – 06:20 am GMT

PDB ID : 7QJ1  
EMDB ID : EMD-14006  
Title : Structure of the recombinant human gamma-Tubulin Ring Complex 6-spoked assembly intermediate (spokes 7-12, homogeneous dataset)  
Authors : Zupa, E.; Pfeffer, S.  
Deposited on : 2021-12-16  
Resolution : 7.00 Å (reported)  
Based on initial models : 7AS4, 6X0U, 6L81, 6V6S

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.0.dev97  
MolProbity : 4.02b-467  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.26


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*ELECTRON MICROSCOPY*

The reported resolution of this entry is 7.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.











Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion  $< 40\%$ ). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	J	1024	
1	l	1024	
2	b	82	
2	m	82	
3	H	907	
3	a	907	
4	G	902	
5	I	667	

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
5	K	667	 74% 9% 16%
6	L	1819	 28% 69%
7	U	451	 9% 68% 21% 7%
7	V	451	 67% 21% 5% 7%
7	W	451	 67% 21% 5% 7%
7	X	451	 68% 21% 5% 7%
7	Y	451	 6% 67% 21% 5% 7%
7	Z	451	 11% 68% 21% 7%

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 50899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gamma-tubulin complex component 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	l	108	847	539	150	157	1	0	0
1	J	534	4429	2893	737	776	23	0	0

- Molecule 2 is a protein called Mitotic-spindle organizing protein 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
2	m	65	484	299	85	96	4	0	0
2	b	65	484	299	85	96	4	0	0

- Molecule 3 is a protein called Gamma-tubulin complex component 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	a	116	933	591	171	169	2	0	0
3	H	594	4907	3130	864	888	25	0	0

- Molecule 4 is a protein called Gamma-tubulin complex component 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	G	636	5186	3342	871	940	33	0	0

- Molecule 5 is a protein called Gamma-tubulin complex component 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	I	521	4225	2737	720	750	18	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	K	562	4579	2964	781	816	18	0	0

- Molecule 6 is a protein called Gamma-tubulin complex component 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	L	566	4587	3000	773	789	25	0	0

- Molecule 7 is a protein called Tubulin gamma-1 chain.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
7	U	420	3373	2134	586	638	15	0	0
7	V	420	3373	2134	586	638	15	0	0
7	W	420	3373	2134	586	638	15	0	0
7	X	420	3373	2134	586	638	15	0	0
7	Y	420	3373	2134	586	638	15	0	0
7	Z	420	3373	2134	586	638	15	0	0

















• Molecule 5: Gamma-tubulin complex component 4




• Molecule 5: Gamma-tubulin complex component 4







• Molecule 6: Gamma-tubulin complex component 6









• Molecule 7: Tubulin gamma-1 chain




• Molecule 7: Tubulin gamma-1 chain







• Molecule 7: Tubulin gamma-1 chain



• Molecule 7: Tubulin gamma-1 chain



• Molecule 7: Tubulin gamma-1 chain



## 4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	59861	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ( $e^-/\text{\AA}^2$ )	35	Depositor
Minimum defocus (nm)	2000	Depositor
Maximum defocus (nm)	2500	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.338	Depositor
Minimum map value	-0.142	Depositor
Average map value	0.001	Depositor
Map value standard deviation	0.008	Depositor
Recommended contour level	0.0571	Depositor
Map size (Å)	532.0, 532.0, 532.0	wwPDB
Map dimensions	200, 200, 200	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	2.66, 2.66, 2.66	Depositor



## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	J	0.34	0/4525	0.70	6/6119 (0.1%)
1	l	0.34	0/863	0.66	3/1166 (0.3%)
2	b	0.42	0/484	0.98	4/653 (0.6%)
2	m	0.42	0/484	0.98	4/653 (0.6%)
3	H	0.35	0/5009	0.67	7/6761 (0.1%)
3	a	0.38	0/948	0.72	1/1277 (0.1%)
4	G	0.36	0/5295	0.71	5/7147 (0.1%)
5	I	0.42	3/4322 (0.1%)	0.67	5/5853 (0.1%)
5	K	0.36	0/4683	0.66	6/6338 (0.1%)
6	L	0.35	0/4697	0.65	7/6348 (0.1%)
7	U	0.33	0/3441	0.71	10/4661 (0.2%)
7	V	0.33	0/3441	0.71	10/4661 (0.2%)
7	W	0.33	0/3441	0.71	10/4661 (0.2%)
7	X	0.33	0/3441	0.71	10/4661 (0.2%)
7	Y	0.33	0/3441	0.71	10/4661 (0.2%)
7	Z	0.33	0/3441	0.71	10/4661 (0.2%)
All	All	0.35	3/51956 (0.0%)	0.70	108/70281 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	J	0	7
3	H	0	1
4	G	0	2
5	I	0	4
5	K	0	2
6	L	0	1
All	All	0	17

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	I	361	TYR	CB-CG	-7.70	1.40	1.51
5	I	530	TYR	CB-CG	-5.12	1.44	1.51
5	I	361	TYR	CD2-CE2	-5.05	1.31	1.39

The worst 5 of 108 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	G	557	LEU	CB-CG-CD2	14.86	136.26	111.00
4	G	557	LEU	CB-CG-CD1	-12.97	88.95	111.00
7	X	343	ARG	NE-CZ-NH2	11.06	125.83	120.30
7	W	343	ARG	NE-CZ-NH2	11.06	125.83	120.30
7	V	343	ARG	NE-CZ-NH2	11.02	125.81	120.30

There are no chirality outliers.

5 of 17 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	G	240	GLY	Peptide
4	G	580	HIS	Peptide
3	H	454	VAL	Peptide
5	I	407	ASP	Peptide
5	I	503	LYS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	J	4429	0	4482	29	0
1	l	847	0	789	0	0
2	b	484	0	512	0	0
2	m	484	0	512	0	0
3	H	4907	0	4896	46	0
3	a	933	0	953	0	0
4	G	5186	0	5219	44	0
5	I	4225	0	4259	48	0
5	K	4579	0	4586	43	0
6	L	4587	0	4636	31	0
7	U	3373	0	3325	45	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	V	3373	0	3325	47	0
7	W	3373	0	3325	46	0
7	X	3373	0	3325	43	0
7	Y	3373	0	3325	47	0
7	Z	3373	0	3325	45	0
All	All	50899	0	50794	484	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 484 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:I:527:ASN:O	5:I:530:TYR:HB3	1.70	0.91
5:K:4:GLU:OE1	5:K:17:PHE:CE1	2.27	0.87
6:L:284:GLU:HA	6:L:287:LEU:HG	1.64	0.79
5:K:4:GLU:OE1	5:K:17:PHE:HE1	1.70	0.74
5:K:4:GLU:CD	5:K:17:PHE:CE1	2.61	0.73

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	J	506/1024 (49%)	469 (93%)	34 (7%)	3 (1%)	25 66
1	l	104/1024 (10%)	98 (94%)	5 (5%)	1 (1%)	15 54
2	b	63/82 (77%)	61 (97%)	2 (3%)	0	100 100
2	m	63/82 (77%)	61 (97%)	2 (3%)	0	100 100
3	H	584/907 (64%)	572 (98%)	11 (2%)	1 (0%)	47 81

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	a	112/907 (12%)	108 (96%)	4 (4%)	0	100	100
4	G	624/902 (69%)	597 (96%)	22 (4%)	5 (1%)	19	60
5	I	511/667 (77%)	489 (96%)	20 (4%)	2 (0%)	34	72
5	K	548/667 (82%)	530 (97%)	17 (3%)	1 (0%)	47	81
6	L	540/1819 (30%)	509 (94%)	27 (5%)	4 (1%)	22	63
7	U	408/451 (90%)	392 (96%)	16 (4%)	0	100	100
7	V	408/451 (90%)	391 (96%)	17 (4%)	0	100	100
7	W	408/451 (90%)	392 (96%)	16 (4%)	0	100	100
7	X	408/451 (90%)	392 (96%)	16 (4%)	0	100	100
7	Y	408/451 (90%)	392 (96%)	16 (4%)	0	100	100
7	Z	408/451 (90%)	392 (96%)	16 (4%)	0	100	100
All	All	6103/10787 (57%)	5845 (96%)	241 (4%)	17 (0%)	44	77

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	l	121	PRO
4	G	241	ARG
4	G	581	ASP
3	H	455	LYS
5	I	601	LEU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	J	498/933 (53%)	493 (99%)	5 (1%)	76	86
1	l	84/933 (9%)	84 (100%)	0	100	100
2	b	53/62 (86%)	40 (76%)	13 (24%)	0	3
2	m	53/62 (86%)	40 (76%)	13 (24%)	0	3
3	H	539/798 (68%)	535 (99%)	4 (1%)	84	90

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	a	101/798 (13%)	100 (99%)	1 (1%)	76	86
4	G	572/791 (72%)	568 (99%)	4 (1%)	84	90
5	I	472/594 (80%)	465 (98%)	7 (2%)	65	80
5	K	509/594 (86%)	506 (99%)	3 (1%)	86	92
6	L	501/1546 (32%)	496 (99%)	5 (1%)	76	86
7	U	376/400 (94%)	310 (82%)	66 (18%)	2	11
7	V	376/400 (94%)	310 (82%)	66 (18%)	2	11
7	W	376/400 (94%)	310 (82%)	66 (18%)	2	11
7	X	376/400 (94%)	310 (82%)	66 (18%)	2	11
7	Y	376/400 (94%)	310 (82%)	66 (18%)	2	11
7	Z	376/400 (94%)	310 (82%)	66 (18%)	2	11
All	All	5638/9511 (59%)	5187 (92%)	451 (8%)	16	35

5 of 451 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
7	W	323	ILE
7	Z	384	ILE
7	X	260	LEU
7	Z	369	SER
7	Z	125	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 93 such sidechains are listed below:

Mol	Chain	Res	Type
7	U	187	ASN
7	W	322	ASN
7	U	229	ASN
7	V	229	ASN
7	X	187	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


## 6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-14006. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

### 6.1 Orthogonal projections [i](#)


#### 6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

### 6.2 Central slices [i](#)


#### 6.2.1 Primary map



The images above show central slices of the map in three orthogonal directions.

## 6.3 Largest variance slices [i](#)

### 6.3.1 Primary map



X Index: 62



Y Index: 101



Z Index: 77

The images above show the largest variance slices of the map in three orthogonal directions.

## 6.4 Orthogonal surface views [i](#)

### 6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 0.0571. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.




## 6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

## 7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

### 7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


## 7.2 Volume estimate [i](#)



The volume at the recommended contour level is 646 nm<sup>3</sup>; this corresponds to an approximate mass of 584 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

### 7.3 Rotationally averaged power spectrum [i](#)



\*Reported resolution corresponds to spatial frequency of 0.143 Å<sup>-1</sup>


## 8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

## 9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-14006 and PDB model 7QJ1. Per-residue inclusion information can be found in section [3](#) on page [6](#).

### 9.1 Map-model overlay [i](#)



The images above show the 3D surface view of the map at the recommended contour level 0.0571 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

## 9.2 Atom inclusion [i](#)



At the recommended contour level, 95% of all backbone atoms, 71% of all non-hydrogen atoms, are inside the map.