

Full wwPDB X-ray Structure Validation Report (i)

May 16, 2020 - 11:17 am BST

PDB ID	:	3QKB
Title	:	Crystal structure of a Protein with unknown function which belongs to Pfam
		DUF74 family (PEPE_0654) from Pediococcus pentosaceus ATCC 25745 at
		2.73 A resolution
Authors	:	Joint Center for Structural Genomics (JCSG)
Deposited on		
Resolution	:	2.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.11
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25 th 2019)
Refmac	:	5.8.0158
CCP4	:	$7.0.044 (\mathrm{Gargrove})$
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.11

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} {f Whole archive}\ (\#{f Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries},{ m resolution\ range}({ m \AA}))$
R _{free}	130704	1271(2.76-2.72)
Clashscore	141614	1322(2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	A	111	% 8 2%	•	15%
1	В	111	80%	•	16%
1	С	111	3% 81%	•	17%
1	D	111	78%	5%	16%
1	Е	111	4% 80%	•	17%



2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 3653 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Λ	94	Total	С	Ν	0	Se	0	1	0
	A	94	717	461	110	143	3	0		0
1	В	93	Total	С	Ν	0	Se	0	1	0
	D	90	716	460	110	143	3	0	T	0
1	С	92	Total	С	Ν	Ο	Se	0	4	0
	U	92	728	470	110	145	3	0	4	
1	D	93	Total	С	Ν	Ο	Se	0	2	0
	D	90	722	465	109	145	3	0	2	0
1	Е	92	Total	С	Ν	Ο	Se	0	3	0
		92	715	460	108	143	4	U	J	0

• Molecule 1 is a protein called Uncharacterized protein.

There are 95 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	EXPRESSION TAG	UNP Q03GF5
A	-17	GLY	-	EXPRESSION TAG	UNP Q03GF5
A	-16	SER	-	EXPRESSION TAG	UNP Q03GF5
A	-15	ASP	-	EXPRESSION TAG	UNP Q03GF5
A	-14	LYS	-	EXPRESSION TAG	UNP Q03GF5
A	-13	ILE	-	EXPRESSION TAG	UNP Q03GF5
A	-12	HIS	-	EXPRESSION TAG	UNP Q03GF5
A	-11	HIS	-	EXPRESSION TAG	UNP Q03GF5
A	-10	HIS	-	EXPRESSION TAG	UNP Q03GF5
A	-9	HIS	-	EXPRESSION TAG	UNP Q03GF5
A	-8	HIS	-	EXPRESSION TAG	UNP Q03GF5
A	-7	HIS	-	EXPRESSION TAG	UNP Q03GF5
A	-6	GLU	-	EXPRESSION TAG	UNP Q03GF5
A	-5	ASN	-	EXPRESSION TAG	UNP Q03GF5
A	-4	LEU	-	EXPRESSION TAG	UNP Q03GF5
А	-3	TYR	-	EXPRESSION TAG	UNP Q03GF5
A	-2	PHE	-	EXPRESSION TAG	UNP Q03GF5
A	-1	GLN	-	EXPRESSION TAG	UNP Q03GF5
А	0	GLY	-	EXPRESSION TAG	UNP Q03GF5

Continued on next page...



3QKB	
------	--

Chain	Residue	vious page Modelled	Actual	Comment	Reference
В	-18	MSE	_	EXPRESSION TAG	UNP Q03GF5
В	-17	GLY	_	EXPRESSION TAG	UNP Q03GF5
В	-16	SER	-	EXPRESSION TAG	UNP Q03GF5
В	-15	ASP	-	EXPRESSION TAG	UNP Q03GF5
В	-14	LYS	-	EXPRESSION TAG	UNP Q03GF5
В	-13	ILE	-	EXPRESSION TAG	UNP Q03GF5
В	-12	HIS	-	EXPRESSION TAG	UNP Q03GF5
В	-11	HIS	-	EXPRESSION TAG	UNP Q03GF5
В	-10	HIS	-	EXPRESSION TAG	UNP Q03GF5
В	-9	HIS	-	EXPRESSION TAG	UNP Q03GF5
В	-8	HIS	-	EXPRESSION TAG	UNP Q03GF5
В	-7	HIS	-	EXPRESSION TAG	UNP Q03GF5
В	-6	GLU	-	EXPRESSION TAG	UNP Q03GF5
В	-5	ASN	-	EXPRESSION TAG	UNP Q03GF5
В	-4	LEU	-	EXPRESSION TAG	UNP Q03GF5
В	-3	TYR	-	EXPRESSION TAG	UNP Q03GF5
В	-2	PHE	-	EXPRESSION TAG	UNP Q03GF5
В	-1	GLN	-	EXPRESSION TAG	UNP Q03GF5
В	0	GLY	-	EXPRESSION TAG	UNP Q03GF5
С	-18	MSE	-	EXPRESSION TAG	UNP Q03GF5
С	-17	GLY	-	EXPRESSION TAG	UNP Q03GF5
С	-16	SER	-	EXPRESSION TAG	UNP Q03GF5
С	-15	ASP	-	EXPRESSION TAG	UNP Q03GF5
С	-14	LYS	-	EXPRESSION TAG	UNP Q03GF5
С	-13	ILE	-	EXPRESSION TAG	UNP Q03GF5
С	-12	HIS	-	EXPRESSION TAG	UNP Q03GF5
С	-11	HIS	-	EXPRESSION TAG	UNP Q03GF5
С	-10	HIS	-	EXPRESSION TAG	UNP Q03GF5
С	-9	HIS	-	EXPRESSION TAG	UNP Q03GF5
С	-8	HIS	-	EXPRESSION TAG	UNP Q03GF5
С	-7	HIS	-	EXPRESSION TAG	UNP Q03GF5
С	-6	GLU	-	EXPRESSION TAG	UNP Q03GF5
С	-5	ASN	-	EXPRESSION TAG	UNP Q03GF5
С	-4	LEU	-	EXPRESSION TAG	UNP Q03GF5
С	-3	TYR	-	EXPRESSION TAG	UNP Q03GF5
С	-2	PHE	-	EXPRESSION TAG	UNP Q03GF5
С	-1	GLN	-	EXPRESSION TAG	UNP Q03GF5
С	0	GLY	-	EXPRESSION TAG	UNP Q03GF5
D	-18	MSE	-	EXPRESSION TAG	UNP Q03GF5
D	-17	GLY	-	EXPRESSION TAG	UNP Q03GF5
D	-16	SER	-	EXPRESSION TAG	UNP Q03GF5
D	-15	ASP	-	EXPRESSION TAG	UNP Q03GF5

Continued from previous page...

Continued on next page...



Chain	Residue	Modelled	Actual	Comment	Reference
D	-14	LYS	-	EXPRESSION TAG	UNP Q03GF5
D	-13	ILE	-	EXPRESSION TAG	UNP Q03GF5
D	-12	HIS	-	EXPRESSION TAG	UNP Q03GF5
D	-11	HIS	-	EXPRESSION TAG	UNP Q03GF5
D	-10	HIS	-	EXPRESSION TAG	UNP Q03GF5
D	-9	HIS	-	EXPRESSION TAG	UNP Q03GF5
D	-8	HIS	-	EXPRESSION TAG	UNP Q03GF5
D	-7	HIS	-	EXPRESSION TAG	UNP Q03GF5
D	-6	GLU	-	EXPRESSION TAG	UNP Q03GF5
D	-5	ASN	-	EXPRESSION TAG	UNP Q03GF5
D	-4	LEU	-	EXPRESSION TAG	UNP Q03GF5
D	-3	TYR	-	EXPRESSION TAG	UNP Q03GF5
D	-2	PHE	-	EXPRESSION TAG	UNP Q03GF5
D	-1	GLN	-	EXPRESSION TAG	UNP Q03GF5
D	0	GLY	-	EXPRESSION TAG	UNP Q03GF5
Ε	-18	MSE	-	EXPRESSION TAG	UNP Q03GF5
Ε	-17	GLY	-	EXPRESSION TAG	UNP Q03GF5
Ε	-16	SER	-	EXPRESSION TAG	UNP Q03GF5
Ε	-15	ASP	-	EXPRESSION TAG	UNP Q03GF5
Е	-14	LYS	-	EXPRESSION TAG	UNP Q03GF5
Ε	-13	ILE	-	EXPRESSION TAG	UNP Q03GF5
Е	-12	HIS	-	EXPRESSION TAG	UNP Q03GF5
Ε	-11	HIS	-	EXPRESSION TAG	UNP Q03GF5
Ε	-10	HIS	-	EXPRESSION TAG	UNP Q03GF5
Е	-9	HIS	-	EXPRESSION TAG	UNP Q03GF5
Ε	-8	HIS	-	EXPRESSION TAG	UNP Q03GF5
Е	-7	HIS	-	EXPRESSION TAG	UNP Q03GF5
Е	-6	GLU	-	EXPRESSION TAG	UNP Q03GF5
Ε	-5	ASN	-	EXPRESSION TAG	UNP Q03GF5
Е	-4	LEU	-	EXPRESSION TAG	UNP Q03GF5
Ε	-3	TYR	-	EXPRESSION TAG	UNP Q03GF5
Е	-2	PHE	-	EXPRESSION TAG	UNP Q03GF5
Е	-1	GLN	-	EXPRESSION TAG	UNP Q03GF5
Е	0	GLY	-	EXPRESSION TAG	UNP Q03GF5

Continued from previous page...

• Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	9	Total O 9 9	0	0
2	В	10	Total O 10 10	0	0

Continued on next page...




Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	С	21	TotalO2222	0	1
2	D	9	Total O 9 9	0	0
2	Ε	5	Total O 5 5	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Uncharacterized protein



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants	110.30Å 67.30 Å 85.84 Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.73 – 2.73	Depositor
Resolution (A)	47.74 - 2.73	EDS
% Data completeness	99.0 (47.73-2.73)	Depositor
(in resolution range)	98.9(47.74-2.73)	EDS
R _{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.10 (at 2.73 Å)	Xtriage
Refinement program	REFMAC $5.5.0110$	Depositor
D D.	0.202 , 0.239	Depositor
R, R_{free}	0.196 , 0.226	DCC
R_{free} test set	884 reflections $(5.08%)$	wwPDB-VP
Wilson B-factor (Å ²)	53.2	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.31 , 54.6	EDS
L-test for twinning ²	$ L > = 0.49, < L^2 > = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3653	wwPDB-VP
Average B, all atoms $(Å^2)$	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.15% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Chain		lengths	Bond angles	
	Ullalli	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.76	0/725	0.66	0/972
1	В	0.77	0/724	0.63	0/970
1	С	0.79	0/746	0.68	0/999
1	D	0.78	0/734	0.67	0/984
1	Е	0.75	0/730	0.64	0/979
All	All	0.77	0/3659	0.66	0/4904

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	В	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	В	-1	GLN	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	717	0	711	2	0
1	В	716	0	712	2	0
1	С	728	0	744	2	0
1	D	722	0	727	5	0
1	Е	715	0	713	3	0
2	А	9	0	0	0	0
2	В	10	0	0	0	0
2	С	22	0	0	0	0
2	D	9	0	0	0	0
2	Е	5	0	0	0	0
All	All	3653	0	3607	9	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (9) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:62:GLY:O	1:E:5:THR:HG21	1.88	0.73
1:E:36[A]:MSE:HA	1:E:36[A]:MSE:HE2	1.72	0.70
1:A:62:GLY:O	1:B:5:THR:HG21	1.96	0.65
1:B:62:GLY:O	1:C:5:THR:HG21	1.97	0.63
1:C:62:GLY:O	1:D:5:THR:HG21	2.01	0.61
1:A:5:THR:HG21	1:E:62:GLY:O	2.01	0.59
1:D:25:MSE:HE3	1:D:78:PHE:CZ	2.47	0.49
1:D:25:MSE:HE3	1:D:78:PHE:CE1	2.52	0.45
1:D:68:GLU:OE1	1:D:82:HIS:NE2	2.50	0.41

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	93/111~(84%)	90~(97%)	2(2%)	1 (1%)	14	26
1	В	92/111~(83%)	90 (98%)	2(2%)	0	100	100
1	С	94/111~(85%)	93~(99%)	1 (1%)	0	100	100
1	D	93/111~(84%)	90~(97%)	3~(3%)	0	100	100
1	Ε	93/111~(84%)	91 (98%)	2(2%)	0	100	100
All	All	465/555 (84%)	454 (98%)	10 (2%)	1 (0%)	47	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	Type
1	А	-1	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	5
1	А	75/91~(82%)	75~(100%)	0	100 100	
1	В	75/91~(82%)	74 (99%)	1 (1%)	69 82	
1	С	80/91~(88%)	80 (100%)	0	100 100	
1	D	78/91~(86%)	78 (100%)	0	100 100	
1	Ε	76/91~(84%)	76~(100%)	0	100 100	
All	All	384/455~(84%)	383 (100%)	1 (0%)	92 95	

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	\mathbf{Res}	Type
1	В	42	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	$\# RSRZ {>}2$	$OWAB(A^2)$	Q<0.9
1	А	91/111~(81%)	-0.19	1 (1%) 80 85	29, 59, 96, 122	0
1	В	90/111~(81%)	0.04	3 (3%) 46 52	26, 52, 83, 101	0
1	С	89/111~(80%)	-0.14	3 (3%) 45 50	24, 43, 73, 89	0
1	D	90/111~(81%)	0.03	5 (5%) 24 27	23, 47, 87, 97	0
1	Ε	89/111~(80%)	0.07	4 (4%) 33 36	31, 55, 95, 112	0
All	All	449/555~(80%)	-0.04	16 (3%) 42 47	23, 51, 92, 122	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	В	0	GLY	4.7
1	D	8	ILE	3.5
1	С	9	ASN	3.3
1	Е	0	GLY	3.0
1	В	-1	GLN	3.0
1	В	56	GLU	2.8
1	С	0	GLY	2.8
1	Е	89	LEU	2.7
1	D	59	GLY	2.5
1	С	8	ILE	2.5
1	А	0	GLY	2.4
1	D	89	LEU	2.3
1	D	90	ILE	2.3
1	Е	9	ASN	2.3
1	D	0	GLY	2.2
1	Е	56	GLU	2.1



6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.

