



# wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 24, 2024 – 10:21 am BST

PDB ID : 8QMC  
Title : High resolution structure of the Streptococcus pneumoniae topoisomerase IV-complex with the V-site 18mer dsDNA and novel fluoroquinolone Delafloxacin  
Authors : Najmudin, S.; Pan, X.S.; Wang, B.; Chayen, N.E.; Fisher, L.M.; Sanderson, M.R.  
Deposited on : 2023-09-21  
Resolution : 2.40 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 3.0  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.002 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.38.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	742	
1	B	742	
2	E	7	
3	F	11	

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	H	11	 <p>36% 27% 55% 18%</p>
5	G	7	 <p>14% 14% 71% 14%</p>

## 2 Entry composition

There are 12 unique types of molecules in this entry. The entry contains 24995 atoms, of which 12198 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA topoisomerase (ATP-hydrolyzing), DNA topoisomerase 4 subunit A.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	724	11662	3651	5882	1007	1097	25	325	3	0
1	B	724	11620	3641	5859	1000	1096	24	321	1	0

There are 38 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	403	MET	-	initiating methionine	UNP J0V1V8
A	404	LYS	-	expression tag	UNP J0V1V8
A	405	ASN	-	expression tag	UNP J0V1V8
A	406	LYS	-	expression tag	UNP J0V1V8
A	407	LYS	-	expression tag	UNP J0V1V8
A	408	ASP	-	expression tag	UNP J0V1V8
A	409	LYS	-	expression tag	UNP J0V1V8
A	410	GLY	-	expression tag	UNP J0V1V8
A	411	LEU	-	expression tag	UNP J0V1V8
A	648	HIS	-	linker	UNP J0V1V8
A	1257	THR	ILE	conflict	UNP P72525
A	1489	LEU	-	expression tag	UNP P72525
A	1490	GLU	-	expression tag	UNP P72525
A	1491	HIS	-	expression tag	UNP P72525
A	1492	HIS	-	expression tag	UNP P72525
A	1493	HIS	-	expression tag	UNP P72525
A	1494	HIS	-	expression tag	UNP P72525
A	1495	HIS	-	expression tag	UNP P72525
A	1496	HIS	-	expression tag	UNP P72525
B	403	MET	-	initiating methionine	UNP J0V1V8
B	404	LYS	-	expression tag	UNP J0V1V8
B	405	ASN	-	expression tag	UNP J0V1V8
B	406	LYS	-	expression tag	UNP J0V1V8
B	407	LYS	-	expression tag	UNP J0V1V8

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	408	ASP	-	expression tag	UNP J0V1V8
B	409	LYS	-	expression tag	UNP J0V1V8
B	410	GLY	-	expression tag	UNP J0V1V8
B	411	LEU	-	expression tag	UNP J0V1V8
B	648	HIS	-	linker	UNP J0V1V8
B	1257	THR	ILE	conflict	UNP P72525
B	1489	LEU	-	expression tag	UNP P72525
B	1490	GLU	-	expression tag	UNP P72525
B	1491	HIS	-	expression tag	UNP P72525
B	1492	HIS	-	expression tag	UNP P72525
B	1493	HIS	-	expression tag	UNP P72525
B	1494	HIS	-	expression tag	UNP P72525
B	1495	HIS	-	expression tag	UNP P72525
B	1496	HIS	-	expression tag	UNP P72525

- Molecule 2 is a DNA chain called DNA (5'-D(\*T)-R(P\*G)-D(P\*T)-R(P\*GP\*GP\*A)-D(P\*T)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
2	E	7	225	70	81	26	42	6	1	0	0

- Molecule 3 is a DNA chain called DNA/RNA (5'-R(P\*GP\*G)-D(P\*TP\*T)-R(P\*A)-D(P\*T)-R(P\*CP\*CP\*AP\*CP\*A)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
3	F	11	349	107	125	40	66	11	1	0	0


- Molecule 4 is a DNA chain called DNA/RNA (5'-R(P\*AP\*AP\*CP\*CP\*G)-D(P\*T)-R(P\*A)-D(P\*TP\*T)-R(P\*AP\*C)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
4	H	11	348	107	125	40	65	11	1	0	0

- Molecule 5 is a DNA chain called DNA/RNA (5'-R(\*G)-D(P\*T)-R(P\*AP\*A)-D(P\*T)-R(P\*AP\*C)-3').


Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P			
5	G	7	221	69	80	27	39	6	1	0	0

- Molecule 6 is MALONIC ACID (three-letter code: MLA) (formula:  $C_3H_4O_4$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
6	A	1	Total	C	H	O	0	0
			9	3	2	4		
6	A	1	Total	C	H	O	0	0
			9	3	2	4		

- Molecule 7 is ACETATE ION (three-letter code: ACT) (formula:  $C_2H_3O_2$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			7	2	3	2		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	A	1	Total	C	H	O	0	0
			7	2	3	2		


- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	3	Total	Mg	0	0
			3	3		
8	B	3	Total	Mg	0	0
			3	3		

- Molecule 9 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total	Cl	0	0
			1	1		
9	B	1	Total	Cl	0	0
			1	1		

- Molecule 10 is (4S)-2-METHYL-2,4-PENTANEDIOL (three-letter code: MPD) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
10	B	1	Total	C	H	O	1	0
			22	6	14	2		

- Molecule 11 is delafloxacin (three-letter code: TE9) (formula:  $C_{18}H_{12}ClF_3N_4O_4$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	
			Total	C	Cl	F	H	N			O
11	F	1	41	18	1	3	11	4	4	1	0
11	H	1	41	18	1	3	11	4	4	1	0

- Molecule 12 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
12	A	193	Total	O	0	0
			193	193		
12	B	199	Total	O	0	0
			199	199		
12	E	8	Total	O	0	0
			8	8		
12	F	9	Total	O	0	0
			9	9		
12	H	10	Total	O	0	0
			10	10		
12	G	7	Total	O	0	0
			7	7		




### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA topoisomerase (ATP-hydrolyzing),DNA topoisomerase 4 subunit A





- Molecule 1: DNA topoisomerase (ATP-hydrolyzing),DNA topoisomerase 4 subunit A






- Molecule 2: DNA (5'-D(\*T)-R(P\*G)-D(P\*T)-R(P\*GP\*GP\*A)-D(P\*T)-3')



- Molecule 3: DNA/RNA (5'-R(P\*GP\*G)-D(P\*TP\*T)-R(P\*A)-D(P\*T)-R(P\*CP\*CP\*AP\*CP\*A)-3')




- Molecule 4: DNA/RNA (5'-R(P\*AP\*AP\*CP\*CP\*G)-D(P\*T)-R(P\*A)-D(P\*TP\*T)-R(P\*AP\*C)-3')



- Molecule 5: DNA/RNA (5'-R(\*G)-D(P\*T)-R(P\*AP\*A)-D(P\*T)-R(P\*AP\*C)-3')





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	157.14Å 157.14Å 211.64Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	64.86 – 2.40 64.86 – 2.40	Depositor EDS
% Data completeness (in resolution range)	50.9 (64.86-2.40) 50.9 (64.86-2.40)	Depositor EDS
$R_{merge}$	0.34	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.13 (at 2.40Å)	Xtrriage
Refinement program	REFMAC 5.8.0352, PDB-REDO	Depositor
R, $R_{free}$	0.241 , 0.299 0.241 , 0.300	Depositor DCC
$R_{free}$ test set	3069 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	24.4	Xtrriage
Anisotropy	0.010	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 23.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.52$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.009 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.87	EDS
Total number of atoms	24995	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MPD, ACT, CL, MG, TE9, MLA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.37	0/5880	0.70	0/7927
1	B	0.36	0/5858	0.68	0/7900
2	E	0.81	0/161	1.40	4/248 (1.6%)
3	F	0.89	0/250	1.52	7/383 (1.8%)
4	H	0.73	0/249	1.55	8/381 (2.1%)
5	G	0.78	0/158	1.51	2/242 (0.8%)
All	All	0.41	0/12556	0.78	21/17081 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	6
1	B	0	2
All	All	0	8

There are no bond length outliers.

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	H	10	DA	P-O3'-C3'	-8.23	109.82	119.70
4	H	3	DC	P-O3'-C3'	-8.07	110.02	119.70
4	H	4	DC	P-O3'-C3'	-7.73	110.43	119.70
4	H	5	DG	P-O3'-C3'	-6.85	111.48	119.70
3	F	1	DG	O4'-C1'-N9	6.70	112.69	108.00

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1276	ARG	Sidechain
1	A	1353	ARG	Sidechain
1	A	537	ILE	Peptide
1	A	550	GLY	Peptide
1	A	551	LYS	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5780	5882	5871	121	0
1	B	5761	5859	5848	133	0
2	E	144	81	82	6	0
3	F	224	125	125	2	0
4	H	223	125	125	6	0
5	G	141	80	81	11	0
6	A	14	4	4	1	0
7	A	8	6	6	0	0
8	A	3	0	0	0	0
8	B	3	0	0	0	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
10	B	8	14	14	3	0
11	F	30	11	0	4	0
11	H	30	11	0	4	0
12	A	193	0	0	26	0
12	B	199	0	0	20	0
12	E	8	0	0	1	0
12	F	9	0	0	1	0
12	G	7	0	0	0	0
12	H	10	0	0	0	0
All	All	12797	12198	12156	270	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 270 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
11:H:101:TE9:CL1	11:H:101:TE9:N2	2.25	1.05
1:B:623:GLY:O	1:B:629:ARG:NH2	1.94	0.99
11:F:101:TE9:CL1	11:F:101:TE9:N2	2.33	0.98
1:B:1001:MET:HB3	1:B:1005:GLN:HE21	1.26	0.98
11:F:101:TE9:CL1	11:F:101:TE9:C2	2.54	0.93

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	725/742 (98%)	657 (91%)	49 (7%)	19 (3%)	<b>4</b> <b>4</b>
1	B	723/742 (97%)	660 (91%)	47 (6%)	16 (2%)	<b>5</b> <b>6</b>
All	All	1448/1484 (98%)	1317 (91%)	96 (7%)	35 (2%)	<b>5</b> <b>5</b>

5 of 35 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	489	ALA
1	A	549	LYS
1	A	610	ASP
1	B	569	LYS
1	B	588	ALA

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	622/635 (98%)	578 (93%)	44 (7%)	12	20
1	B	620/635 (98%)	582 (94%)	38 (6%)	15	27
All	All	1242/1270 (98%)	1160 (93%)	82 (7%)	14	23

5 of 82 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	569	LYS
1	B	1139	VAL
1	B	579	ARG
1	B	1078	ASP
1	B	1236	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	515	GLN
1	B	1049	ASN
1	B	578	GLN
1	B	1053	ASN
1	B	1005	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 8 are monoatomic - leaving 7 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	MLA	A	1501	-	6,6,6	1.25	0	7,7,7	1.08	0
10	MPD	B	1501	-	7,7,7	0.21	0	9,10,10	0.56	0
7	ACT	A	1503	-	3,3,3	0.98	0	3,3,3	1.10	0
7	ACT	A	1504	-	3,3,3	1.13	0	3,3,3	0.91	0
6	MLA	A	1502	-	6,6,6	1.29	0	7,7,7	1.06	0
11	TE9	H	101	8	31,33,33	1.15	3 (9%)	37,51,51	1.20	2 (5%)
11	TE9	F	101	8	31,33,33	1.09	3 (9%)	37,51,51	1.21	3 (8%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	MLA	A	1501	-	-	0/4/4/4	-
10	MPD	B	1501	-	-	0/5/5/5	-
11	TE9	F	101	8	-	3/11/20/20	0/4/4/4
6	MLA	A	1502	-	-	2/4/4/4	-
11	TE9	H	101	8	-	7/11/20/20	0/4/4/4

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
11	H	101	TE9	C2-C16	-2.90	1.39	1.42
11	H	101	TE9	C1-C18	2.52	1.42	1.40
11	F	101	TE9	C2-C16	-2.40	1.39	1.42
11	F	101	TE9	C1-C18	2.27	1.42	1.40
11	H	101	TE9	C3-N3	2.20	1.42	1.37

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	H	101	TE9	C17-C18-C1	-3.28	119.56	121.70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
11	F	101	TE9	C17-C18-C1	-3.24	119.59	121.70
11	H	101	TE9	C4-C3-N3	-2.60	120.55	123.75
11	F	101	TE9	C4-C3-N3	-2.39	120.80	123.75
11	F	101	TE9	C1-N2-C2	2.10	122.34	116.99

There are no chirality outliers.

5 of 12 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
11	F	101	TE9	C16-C2-N3-C15
11	F	101	TE9	N2-C2-N3-C3
11	H	101	TE9	C16-C2-N3-C15
11	H	101	TE9	N2-C2-N3-C3
11	H	101	TE9	C6-C4-C5-O2


There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
10	B	1501	MPD	3	0
6	A	1502	MLA	1	0
11	H	101	TE9	4	0
11	F	101	TE9	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	724/742 (97%)	0.81	133 (18%) 4 4	6, 21, 65, 93	3 (0%)
1	B	724/742 (97%)	0.61	107 (14%) 7 6	6, 19, 57, 87	1 (0%)
2	E	7/7 (100%)	0.45	1 (14%) 7 6	16, 16, 36, 52	0
3	F	11/11 (100%)	0.71	1 (9%) 16 14	20, 28, 51, 55	0
4	H	11/11 (100%)	1.56	4 (36%) 1 1	37, 47, 64, 65	0
5	G	7/7 (100%)	0.15	1 (14%) 7 6	12, 14, 36, 54	0
All	All	1484/1520 (97%)	0.72	247 (16%) 5 5	6, 21, 61, 93	4 (0%)

The worst 5 of 247 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	644	ALA	8.6
1	A	571	PHE	8.3
1	B	555	VAL	8.2
1	B	572	GLY	7.8
1	A	574	GLY	7.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
8	MG	B	1504	1/1	0.77	0.11	9,9,9,9	0
6	MLA	A	1502	7/7	0.79	0.14	24,26,28,28	0
8	MG	B	1502	1/1	0.81	0.10	15,15,15,15	0
7	ACT	A	1504	4/4	0.83	0.13	22,22,23,23	0
9	CL	A	1508	1/1	0.86	0.14	17,17,17,17	0
11	TE9	H	101	30/30	0.88	0.15	30,42,45,45	1
11	TE9	F	101	30/30	0.89	0.12	22,25,29,30	1
6	MLA	A	1501	7/7	0.89	0.11	23,23,24,24	0
8	MG	A	1507	1/1	0.90	0.06	17,17,17,17	0
9	CL	B	1505	1/1	0.91	0.06	15,15,15,15	0
8	MG	A	1505	1/1	0.91	0.04	11,11,11,11	0
8	MG	A	1506	1/1	0.91	0.12	8,8,8,8	0
7	ACT	A	1503	4/4	0.93	0.11	13,13,13,14	0
10	MPD	B	1501	8/8	0.94	0.11	11,12,14,30	1
8	MG	B	1503	1/1	0.96	0.07	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around TE9 H 101:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.