



Full wwPDB X-ray Structure Validation Report i

Jun 20, 2024 – 04:22 pm BST

PDB ID : 8QNF
Title : Crystal structure of the Condensation domain TomBC from the Tomaymycin non-ribosomal peptide synthetase
Authors : Karanth, M.; Schmelz, S.; Kirkpatrick, J.; Krausze, J.; Scrima, A.; Carluccio, T.
Deposited on : 2023-09-26
Resolution : 1.65 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The types of validation reports are described at
<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.37.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	130704	1827 (1.66-1.66)
Clashscore	141614	1931 (1.66-1.66)
Ramachandran outliers	138981	1891 (1.66-1.66)
Sidechain outliers	138945	1891 (1.66-1.66)
RSRZ outliers	127900	1791 (1.66-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	612	-	-	X	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	GOL	A	613	-	-	X	-

2 Entry composition [\(i\)](#)


There are 6 unique types of molecules in this entry. The entry contains 7933 atoms, of which 3764 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Condensation domain TomBC from the Tomaymycin non-ribosomal peptide synthetase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	448	7493	2348	3738	708	690	9	0	53	0

- Molecule 2 is FORMIC ACID (three-letter code: FMT) (formula: CH₂O₂).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	4	1	1	2	0	0
2	A	1	4	1	1	2	0	0
2	A	1	4	1	1	2	0	0
2	A	1	4	1	1	2	0	0
2	A	1	4	1	1	2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C H O 4 1 1 2	0	0
2	A	1	Total C H O 4 1 1 2	0	0
2	A	1	Total C H O 4 1 1 2	0	0
2	A	1	Total C H O 4 1 1 2	0	0
2	A	1	Total C H O 4 1 1 2	0	0
2	A	1	Total C H O 4 1 1 2	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H O 14 3 8 3	0	0
3	A	1	Total C H O 13 3 7 3	0	0

- Molecule 4 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	7	Total Na 7 7	0	0

- Molecule 5 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total K 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	360	Total O 360 360	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Condensation domain TomBC from the Tomaymycin non-ribosomal peptide synthetase



4 Data and refinement statistics i

Property	Value	Source
Space group	F 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	121.65 Å 160.06 Å 163.90 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.94 – 1.65 19.94 – 1.65	Depositor EDS
% Data completeness (in resolution range)	85.4 (19.94-1.65) 85.4 (19.94-1.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.17 (at 1.65 Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R , R_{free}	0.177 , 0.197 0.177 , 0.196	Depositor DCC
R_{free} test set	4067 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	30.6	Xtriage
Anisotropy	0.273	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.43 , 46.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.017 for -h,l,k	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	7933	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NA, FMT, GOL, K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.72	2/4040 (0.0%)	0.91	9/5494 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	117	TYR	CG-CD2	5.86	1.46	1.39
1	A	377	TYR	CZ-OH	5.46	1.47	1.37

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	470	LEU	CA-CB-CG	6.81	130.96	115.30
1	A	235	LEU	CA-CB-CG	-6.75	99.79	115.30
1	A	323	ASP	CB-CG-OD1	5.64	123.38	118.30
1	A	313	LEU	CB-CG-CD2	-5.62	101.44	111.00
1	A	323	ASP	CB-CG-OD2	-5.58	113.28	118.30
1	A	490	ASP	CB-CG-OD1	5.42	123.17	118.30
1	A	227	ASP	CB-CG-OD1	5.31	123.08	118.30
1	A	184[A]	ASP	CB-CG-OD1	5.02	122.82	118.30
1	A	184[B]	ASP	CB-CG-OD1	5.02	122.82	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3755	3738	3508	47	0
2	A	33	11	11	4	0
3	A	12	15	15	6	0
4	A	7	0	0	0	0
5	A	2	0	0	0	0
6	A	360	0	0	14	0
All	All	4169	3764	3534	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123[B]:VAL:HG21	1:A:235:LEU:HD13	1.41	1.01
1:A:181:LYS:NZ	6:A:701:HOH:O	1.99	0.94
1:A:123[B]:VAL:HG21	1:A:235:LEU:CD1	2.10	0.81
1:A:477:ASP:N	1:A:480:GLY:O	2.20	0.73
1:A:192[B]:ASN:OD1	6:A:702:HOH:O	2.08	0.70
1:A:356[B]:ARG:HD2	1:A:385:GLN:OE1	1.93	0.68
1:A:243[B]:ASP:OD1	6:A:703:HOH:O	2.11	0.67
1:A:90:ARG:NH2	6:A:708:HOH:O	2.22	0.67
1:A:330:ARG:NH1	6:A:711:HOH:O	2.29	0.66
1:A:236[C]:ARG:NH1	1:A:373:GLU:OE1	2.25	0.65
1:A:414:ALA:O	6:A:704:HOH:O	2.15	0.64
3:A:612:GOL:O3	3:A:613:GOL:H11	1.97	0.64
2:A:605:FMT:H	6:A:798:HOH:O	2.01	0.60
1:A:476:ARG:HA	1:A:481:LEU:HA	1.84	0.60
1:A:240[B]:ASP:OD1	6:A:705:HOH:O	2.16	0.59
3:A:612:GOL:O2	3:A:613:GOL:H11	2.02	0.59
1:A:235:LEU:HD12	1:A:458[B]:ILE:CD1	2.36	0.56
1:A:482:HIS:HD2	6:A:1019:HOH:O	1.91	0.54
1:A:176[B]:ARG:NH1	6:A:715:HOH:O	2.39	0.53
1:A:347:PHE:CE1	1:A:351:HIS:CE1	2.97	0.53
1:A:471:SER:OG	2:A:610:FMT:O1	2.19	0.52
1:A:476:ARG:HA	1:A:480:GLY:O	2.10	0.52
1:A:295:ASP:OD1	3:A:613:GOL:H12	2.09	0.52
3:A:612:GOL:C3	3:A:613:GOL:H11	2.40	0.52
1:A:356[A]:ARG:HG2	1:A:387[A]:GLU:HG2	1.93	0.51
1:A:476:ARG:O	1:A:476:ARG:HG3	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:SER:HB3	1:A:414:ALA:HB2	1.94	0.49
1:A:286[B]:ARG:HH11	1:A:385:GLN:HB3	1.78	0.49
3:A:612:GOL:O2	3:A:613:GOL:C1	2.60	0.49
1:A:295:ASP:OD1	3:A:613:GOL:C1	2.61	0.48
1:A:91:ASP:OD1	1:A:93:ARG:HD3	2.13	0.47
1:A:377:TYR:CE1	1:A:379[C]:VAL:CG2	2.97	0.47
1:A:403:GLU:OE1	6:A:706:HOH:O	2.21	0.46
1:A:176[B]:ARG:NH2	2:A:607:FMT:O1	2.47	0.46
1:A:147[B]:VAL:HG12	1:A:225:ALA:HB1	1.97	0.46
1:A:414:ALA:HB3	6:A:704:HOH:O	2.16	0.46
1:A:235:LEU:HD12	1:A:458[B]:ILE:HD11	1.98	0.46
1:A:382[A]:VAL:HG11	1:A:408:ALA:HB1	1.98	0.46
1:A:326:GLU:OE2	1:A:476:ARG:HD3	2.15	0.45
1:A:377:TYR:CZ	1:A:379[C]:VAL:HG23	2.52	0.45
1:A:281:ASP:OD1	1:A:422:ARG:NH2	2.36	0.45
1:A:377:TYR:CE1	1:A:379[B]:VAL:HG23	2.53	0.44
1:A:265[B]:GLN:CD	1:A:268:ARG:HH21	2.23	0.42
1:A:235:LEU:CD1	1:A:458[B]:ILE:CD1	2.98	0.41
1:A:438[A]:MET:SD	1:A:440[A]:VAL:CG2	3.09	0.41
1:A:197:LEU:HD21	1:A:222:HIS:CD2	2.55	0.41
1:A:377:TYR:CZ	1:A:379[C]:VAL:CG2	3.04	0.40
1:A:430:GLY:O	6:A:707:HOH:O	2.22	0.40
1:A:257[B]:ARG:HD3	6:A:842:HOH:O	2.22	0.40
2:A:606:FMT:H	2:A:611:FMT:O2	2.20	0.40
1:A:235:LEU:HD12	1:A:458[B]:ILE:HD12	2.02	0.40
1:A:123[B]:VAL:CG2	1:A:235:LEU:CD1	2.90	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	503/537 (94%)	491 (98%)	12 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	410/434 (94%)	398 (97%)	12 (3%)	42 16

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	265[A]	GLN
1	A	265[B]	GLN
1	A	267	TYR
1	A	356[A]	ARG
1	A	356[B]	ARG
1	A	389	ASP
1	A	422	ARG
1	A	428	SER
1	A	476	ARG
1	A	484	LEU
1	A	514[B]	ARG
1	A	514[C]	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 22 ligands modelled in this entry, 9 are monoatomic - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FMT	A	611	-	2,2,2	0.50	0	1,1,1	0.52	0
3	GOL	A	613	-	5,5,5	1.31	1 (20%)	5,5,5	0.68	0
2	FMT	A	608	-	2,2,2	0.52	0	1,1,1	0.48	0
3	GOL	A	612	-	5,5,5	1.11	0	5,5,5	1.08	0
2	FMT	A	605	-	2,2,2	0.71	0	1,1,1	0.57	0
2	FMT	A	601	-	2,2,2	1.04	0	1,1,1	0.26	0
2	FMT	A	604	-	2,2,2	0.29	0	1,1,1	0.65	0
2	FMT	A	610	-	2,2,2	0.18	0	1,1,1	0.20	0
2	FMT	A	606	-	2,2,2	0.55	0	1,1,1	0.06	0
2	FMT	A	607	-	2,2,2	0.70	0	1,1,1	0.25	0
2	FMT	A	603	-	2,2,2	0.29	0	1,1,1	0.31	0
2	FMT	A	609	-	2,2,2	0.75	0	1,1,1	0.25	0
2	FMT	A	602	-	2,2,2	0.64	0	1,1,1	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	613	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	612	-	-	2/4/4/4	-

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	613	GOL	O2-C2	-2.29	1.36	1.43

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	612	GOL	C1-C2-C3-O3
3	A	613	GOL	O1-C1-C2-C3
3	A	613	GOL	O1-C1-C2-O2
3	A	612	GOL	O2-C2-C3-O3

There are no ring outliers.

7 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	611	FMT	1	0
3	A	613	GOL	6	0
3	A	612	GOL	4	0
2	A	605	FMT	1	0
2	A	610	FMT	1	0
2	A	606	FMT	1	0
2	A	607	FMT	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	448/537 (83%)	0.37	41 (9%) 9 8	23, 33, 70, 104	0

All (41) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	429	SER	8.7
1	A	428	SER	8.2
1	A	430	GLY	8.1
1	A	107	ILE	8.0
1	A	478	SER	8.0
1	A	106	LEU	6.8
1	A	479	GLU	6.5
1	A	527	ALA	6.2
1	A	91	ASP	6.0
1	A	156	ASP	5.8
1	A	92	GLY	5.8
1	A	426	PRO	5.4
1	A	93	ARG	5.2
1	A	526	GLY	5.1
1	A	427	ARG	4.7
1	A	480	GLY	4.7
1	A	453	ASP	4.6
1	A	110	TYR	4.4
1	A	477	ASP	4.1
1	A	476	ARG	4.1
1	A	431	TYR	3.9
1	A	425	PRO	3.9
1	A	80	ILE	3.8
1	A	81	PRO	3.4
1	A	424	GLN	3.4
1	A	108	ASP	3.3
1	A	155	SER	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	452	ASP	3.1
1	A	524	ALA	3.0
1	A	320	ALA	2.8
1	A	247	ARG	2.7
1	A	89	PRO	2.6
1	A	147[A]	VAL	2.4
1	A	273	PRO	2.3
1	A	263	PHE	2.3
1	A	274	GLU	2.2
1	A	454	THR	2.2
1	A	511	PHE	2.2
1	A	260	TYR	2.2
1	A	158	THR	2.2
1	A	470	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	FMT	A	606	3/3	0.58	0.21	66,70,70,84	0
3	GOL	A	613	6/6	0.72	0.34	57,63,76,76	0
2	FMT	A	608	3/3	0.80	0.46	60,72,72,75	0
2	FMT	A	609	3/3	0.83	0.25	60,63,67,76	0
2	FMT	A	611	3/3	0.83	0.17	53,60,63,64	0
2	FMT	A	607	3/3	0.83	0.32	58,61,64,78	0
2	FMT	A	610	3/3	0.87	0.15	45,50,53,55	0
3	GOL	A	612	6/6	0.89	0.17	34,57,73,76	0
2	FMT	A	604	3/3	0.90	0.11	39,47,55,57	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	K	A	621	1/1	0.90	0.18	84,84,84,84	0
2	FMT	A	605	3/3	0.93	0.11	45,54,57,69	0
2	FMT	A	603	3/3	0.94	0.17	41,49,52,62	0
4	NA	A	618	1/1	0.95	0.07	47,47,47,47	0
2	FMT	A	602	3/3	0.96	0.06	42,48,49,59	0
4	NA	A	619	1/1	0.97	0.08	43,43,43,43	0
4	NA	A	620	1/1	0.98	0.12	29,29,29,29	0
2	FMT	A	601	3/3	0.98	0.06	31,34,39,42	0
5	K	A	622	1/1	0.98	0.12	50,50,50,50	0
4	NA	A	616	1/1	0.99	0.11	33,33,33,33	0
4	NA	A	614	1/1	0.99	0.15	33,33,33,33	0
4	NA	A	615	1/1	0.99	0.08	36,36,36,36	0
4	NA	A	617	1/1	1.00	0.09	40,40,40,40	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.