



# Full wwPDB X-ray Structure Validation Report

Oct 18, 2022 – 10:54 am BST

PDB ID : 7QQX  
Title : SpCas9 bound to FANCF off-target5 DNA substrate  
Authors : Pacesa, M.; Jinek, M.  
Deposited on : 2022-01-10  
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0267  
CCP4 : 7.1.010 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)
RNA backbone	3102	1174 (2.80-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	84	 5% 73% 25%
2	B	1368	 2% 88% 9%
3	C	28	 4% 82% 18%
4	D	12	 8% 58% 33% 8%

## 2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 14001 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called FANCF sgRNA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	A	84	1763	789	317	574	83	0	0	0

- Molecule 2 is a protein called CRISPR-associated endonuclease Cas9/Csn1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	1336	10919	6957	1895	2045	22	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	10	ALA	ASP	engineered mutation	UNP Q99ZW2
B	840	ALA	HIS	engineered mutation	UNP Q99ZW2

- Molecule 3 is a DNA chain called FANCF off-target5 target strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
3	C	28	577	275	109	166	27	0	0	0

- Molecule 4 is a DNA chain called FANCF off-target5 non-target strand.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
4	D	11	207	98	37	62	10	0	0	1

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Mg	0	0
			2	2		

- Molecule 6 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	2	Total K 2 2	0	0
6	B	7	Total K 7 7	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	146	Total O 146 146	0	0
7	B	335	Total O 335 335	0	0
7	C	32	Total O 32 32	0	0
7	D	11	Total O 11 11	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: FANCF sgRNA




- Molecule 2: CRISPR-associated endonuclease Cas9/Csn1




- Molecule 3: FANCF off-target5 target strand





- Molecule 4: FANCF off-target5 non-target strand



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	178.53Å 67.38Å 188.39Å 90.00° 110.90° 90.00°	Depositor
Resolution (Å)	48.00 – 2.40 48.00 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.1 (48.00-2.40) 99.1 (48.00-2.40)	Depositor EDS
$R_{merge}$	0.12	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.34 (at 2.39Å)	Xtriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.182 , 0.216 0.182 , 0.216	Depositor DCC
$R_{free}$ test set	4091 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.7	Xtriage
Anisotropy	0.328	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	14001	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	69.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/1973	0.84	0/3073
2	B	0.27	0/11111	0.44	0/14928
3	C	0.65	0/648	0.99	0/1000
4	D	0.67	0/231	0.94	0/356
All	All	0.32	0/13963	0.57	0/19357

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1763	0	888	12	0
2	B	10919	0	11099	78	0
3	C	577	0	317	4	0
4	D	207	0	114	2	0
5	A	2	0	0	0	0
6	A	2	0	0	0	0
6	B	7	0	0	0	0
7	A	146	0	0	0	0
7	B	335	0	0	6	0

*Continued on next page...*



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	C	32	0	0	2	0
7	D	11	0	0	0	0
All	All	14001	0	12418	90	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (90) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:80:CYS:SG	7:B:1775:HOH:O	2.37	0.83
2:B:574:CYS:SG	7:B:1665:HOH:O	2.47	0.72
2:B:1004:LEU:HD11	2:B:1042:ILE:HD11	1.73	0.71
3:C:-7:DC:H2'	3:C:-6:DA:C8	2.28	0.68
1:A:59:U:OP1	2:B:467:ARG:NH2	2.29	0.66
2:B:1042:ILE:HG23	2:B:1043:MET:HG2	1.79	0.65
2:B:967:ARG:NH1	2:B:986:ASP:OD1	2.31	0.64
3:C:8:DG:N7	7:C:101:HOH:O	2.31	0.61
2:B:1212:ARG:NH2	2:B:1280:VAL:O	2.35	0.59
1:A:49:A:N3	2:B:1122:ARG:NH2	2.48	0.59
2:B:654:ARG:HD2	2:B:656:TYR:CZ	2.40	0.57
2:B:780:ARG:NH1	2:B:806:LEU:O	2.38	0.57
2:B:763:MET:HE1	2:B:931:VAL:HG21	1.86	0.56
2:B:266:LEU:HD22	2:B:294:LYS:HD2	1.88	0.56
2:B:967:ARG:NH2	7:B:1501:HOH:O	2.31	0.56
2:B:271:TYR:HA	2:B:274:ASP:HB2	1.88	0.56
2:B:258:LEU:HD11	2:B:282:ILE:HD11	1.89	0.55
2:B:870:VAL:HG23	2:B:908:LEU:HG	1.87	0.55
2:B:530:VAL:HG22	2:B:537:PRO:HA	1.88	0.55
4:D:-2:DC:H2''	4:D:-1:DC:H5''	1.89	0.55
2:B:790:GLU:HG2	2:B:889:ALA:HA	1.89	0.54
1:A:20:C:OP2	2:B:403:ARG:NH1	2.40	0.54
2:B:1110:ILE:HG23	2:B:1122:ARG:HD2	1.90	0.54
2:B:1205:GLU:OE1	2:B:1359:ARG:NH2	2.41	0.54
1:A:33:G:H5'	1:A:34:A:OP2	2.08	0.54
2:B:898:ASP:O	2:B:905:ARG:NH2	2.41	0.53
2:B:784:ILE:O	2:B:788:ILE:HG12	2.10	0.51
2:B:184:LEU:HD12	2:B:299:ALA:HB2	1.93	0.50
2:B:250:PRO:HD2	2:B:264:LEU:O	2.12	0.49
2:B:936:ASP:OD1	2:B:951:ARG:NH1	2.43	0.49
2:B:1243:GLU:HG3	2:B:1244:LYS:H	1.76	0.49

Continued on next page...

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:557:ARG:NH2	2:B:596:ASP:OD1	2.46	0.49
2:B:265:GLN:HG2	2:B:267:SER:H	1.79	0.48
2:B:400:ARG:NH2	2:B:406:ASP:OD2	2.45	0.48
2:B:1117:ASP:OD1	2:B:1117:ASP:N	2.46	0.47
2:B:226:ILE:HG13	2:B:234:LYS:HA	1.97	0.47
2:B:175:ASN:N	2:B:176:PRO:HD3	2.30	0.47
2:B:501:ASN:HB3	2:B:708:ILE:HD12	1.96	0.47
1:A:33:G:H5''	1:A:33:G:H8	1.80	0.47
2:B:901:THR:O	2:B:904:GLU:HG2	2.15	0.47
2:B:838:VAL:HG12	2:B:855:LYS:HE2	1.96	0.46
2:B:1245:LEU:O	2:B:1245:LEU:HD12	2.15	0.46
2:B:468:LYS:HG3	2:B:483:ASP:HB2	1.98	0.45
2:B:870:VAL:HG13	2:B:871:PRO:HD2	1.97	0.45
2:B:1306:ALA:O	2:B:1310:ILE:HG12	2.16	0.45
1:A:32:A:H2'	1:A:33:G:O4'	2.16	0.45
2:B:839:ASP:O	2:B:856:VAL:HG12	2.17	0.45
2:B:294:LYS:NZ	7:B:1521:HOH:O	2.48	0.45
2:B:962:LEU:HD23	2:B:962:LEU:HA	1.76	0.45
2:B:882:TYR:CZ	2:B:886:LEU:HD11	2.52	0.45
2:B:89:GLU:HG3	2:B:432:PHE:CD1	2.52	0.45
2:B:673:LYS:HE3	2:B:673:LYS:HB2	1.86	0.45
2:B:285:GLN:H	2:B:285:GLN:CD	2.21	0.44
2:B:178:ASN:HA	2:B:184:LEU:HD11	1.99	0.44
2:B:822:MET:HG3	2:B:883:TRP:HE1	1.82	0.44
1:A:4:A:OP1	2:B:661:ARG:NE	2.29	0.44
1:A:33:G:N2	1:A:36:A:OP2	2.51	0.44
1:A:41:A:OP2	2:B:340:ARG:NH2	2.49	0.44
2:B:258:LEU:CD1	2:B:282:ILE:HD11	2.48	0.43
2:B:1151:LYS:HD2	2:B:1158:LYS:HD2	1.99	0.43
2:B:1210:ARG:HA	2:B:1280:VAL:HG22	2.00	0.43
2:B:78:ARG:NH2	7:B:1531:HOH:O	2.51	0.43
3:C:-1:DC:H2'	3:C:0:DT:C6	2.53	0.43
1:A:34:A:OP1	1:A:34:A:H8	2.01	0.43
2:B:704:PHE:O	2:B:708:ILE:HG12	2.19	0.43
2:B:721:HIS:HB2	7:B:1815:HOH:O	2.18	0.43
2:B:923:GLU:OE2	2:B:925:ARG:NH1	2.51	0.43
2:B:958:LEU:HD22	2:B:962:LEU:HD12	2.00	0.43
1:A:17:C:OP2	2:B:74:ARG:NH1	2.52	0.43
2:B:240:ASN:ND2	2:B:255:ASN:OD1	2.40	0.43
2:B:256:PHE:HB2	2:B:258:LEU:HD13	2.00	0.43
2:B:276:ASP:HB3	2:B:599:LYS:NZ	2.33	0.43

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:1284:ASP:N	2:B:1284:ASP:OD1	2.52	0.43
1:A:70:C:H2'	1:A:71:U:C6	2.54	0.42
2:B:357:ASN:ND2	2:B:371:GLU:HB3	2.34	0.42
2:B:174:LEU:HD21	2:B:302:LEU:HD21	2.00	0.42
2:B:1044:ASN:HB3	2:B:1050:ILE:HG21	2.02	0.42
2:B:1204:PHE:HE1	2:B:1214:LEU:HD13	1.84	0.42
4:D:6:DT:H2''	4:D:7:DG:H5''	2.01	0.42
2:B:253:LYS:HD2	2:B:261:ASP:OD1	2.20	0.42
2:B:182:ASP:OD2	2:B:209:LYS:HB2	2.20	0.41
2:B:1210:ARG:NH2	2:B:1341:GLU:OE2	2.54	0.41
2:B:952:GLU:H	2:B:952:GLU:HG3	1.72	0.41
2:B:1339:THR:O	2:B:1342:VAL:HG22	2.21	0.41
2:B:733:ILE:HD12	2:B:733:ILE:HA	1.93	0.40
2:B:830:ILE:O	2:B:833:LEU:HG	2.21	0.40
2:B:1163:LEU:HD23	2:B:1163:LEU:HA	1.91	0.40
2:B:850:ASP:O	2:B:855:LYS:HD2	2.22	0.40
2:B:1163:LEU:HG	2:B:1343:LEU:HD21	2.02	0.40
3:C:8:DG:H5''	7:C:124:HOH:O	2.21	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [i](#)

#### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	B	1328/1368 (97%)	1292 (97%)	36 (3%)	0	<b>100</b>   <b>100</b>

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar

resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	1198/1225 (98%)	1188 (99%)	10 (1%)	81 91

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	55	SER
2	B	174	LEU
2	B	301	LEU
2	B	409	SER
2	B	419	LEU
2	B	512	SER
2	B	514	LEU
2	B	706	GLU
2	B	853	ASP
2	B	1154	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	309	ASN
2	B	497	ASN
2	B	1041	ASN
2	B	1044	ASN

### 5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	A	82/84 (97%)	13 (15%)	0

All (13) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	A	0	G
1	A	1	G
1	A	2	G
1	A	8	C

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	20	C
1	A	28	A
1	A	29	G
1	A	51	A
1	A	56	U
1	A	59	U
1	A	68	A
1	A	72	U
1	A	77	A

There are no RNA pucker outliers to report.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 11 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	84/84 (100%)	-0.09	4 (4%) 30 29	37, 59, 178, 217	0
2	B	1336/1368 (97%)	-0.03	30 (2%) 62 60	36, 63, 111, 154	0
3	C	28/28 (100%)	-0.13	1 (3%) 42 42	50, 57, 147, 161	0
4	D	11/12 (91%)	0.05	1 (9%) 9 8	53, 76, 142, 147	0
All	All	1459/1492 (97%)	-0.04	36 (2%) 57 55	36, 63, 115, 217	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	-1	G	4.9
2	B	715	GLY	4.6
1	A	74	A	4.5
2	B	1052	LEU	4.3
1	A	0	G	4.1
2	B	1032	ALA	3.8
1	A	75	A	3.7
2	B	802	GLU	3.3
2	B	1058	ARG	3.3
2	B	1034	ALA	3.3
2	B	1053	ALA	3.1
2	B	338	LEU	2.9
2	B	205	GLY	2.8
2	B	778	ARG	2.8
2	B	177	ASP	2.8
2	B	291	LEU	2.8
2	B	381	GLU	2.7
2	B	339	VAL	2.6
2	B	1244	LYS	2.5
3	C	-7	DC	2.5
2	B	716	GLN	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	1012	ASP	2.3
2	B	1302	ILE	2.3
2	B	293	ALA	2.3
2	B	1243	GLU	2.3
2	B	286	TYR	2.3
2	B	311	GLU	2.3
2	B	290	PHE	2.3
2	B	1258	PHE	2.3
2	B	806	LEU	2.2
2	B	343	LEU	2.1
2	B	188	LEU	2.1
4	D	7	DG	2.0
2	B	287	ALA	2.0
2	B	804	THR	2.0
2	B	1151	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	MG	A	101	1/1	0.86	0.11	77,77,77,77	0
6	K	A	104	1/1	0.90	0.14	62,62,62,62	0
6	K	B	1401	1/1	0.90	0.21	115,115,115,115	0
6	K	B	1406	1/1	0.92	0.10	102,102,102,102	0
6	K	B	1403	1/1	0.94	0.10	77,77,77,77	0
6	K	B	1404	1/1	0.95	0.14	83,83,83,83	0
5	MG	A	102	1/1	0.97	0.12	33,33,33,33	0
6	K	B	1405	1/1	0.98	0.04	87,87,87,87	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	K	B	1402	1/1	0.98	0.10	56,56,56,56	0
6	K	B	1407	1/1	0.98	0.05	56,56,56,56	0
6	K	A	103	1/1	0.99	0.13	53,53,53,53	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.