



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 21, 2024 – 09:44 PM EST

PDB ID : 4QRM
Title : crystal structure of a binary complex of FliM-FliG middle domains from T.maritima
Authors : Crane, B.R.; Sircar, R.
Deposited on : 2014-07-01
Resolution : 4.32 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.32 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1018 (4.84-3.80)
Clashscore	141614	1081 (4.84-3.80)
Ramachandran outliers	138981	1033 (4.84-3.80)
Sidechain outliers	138945	1016 (4.84-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	183	64% (green), 32% (yellow), 2% (orange), 2% (red), 0% (grey)
1	C	183	67% (green), 31% (yellow), 1% (orange), 1% (red), 0% (grey)
1	E	183	64% (green), 32% (yellow), 2% (orange), 2% (red), 0% (grey)
1	G	183	65% (green), 32% (yellow), 1% (orange), 1% (red), 0% (grey)
1	I	183	68% (green), 31% (yellow), 1% (orange), 1% (red), 0% (grey)
1	K	183	74% (green), 23% (yellow), 1% (orange), 1% (red), 0% (grey)
1	M	183	69% (green), 29% (yellow), 1% (orange), 1% (red), 0% (grey)

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	O	183	 71% 26% .
1	Q	183	 67% 32% ..
1	S	183	 67% 30% ..
1	U	183	 64% 32% .
2	B	75	 61% 36% ..
2	D	75	 80% 20%
2	F	75	 75% 23% .
2	H	75	 77% 23%
2	J	75	 81% 17% .
2	L	75	 69% 28% .
2	N	75	 81% 17% .
2	P	75	 71% 25% ..
2	R	75	 67% 32% .
2	T	75	 69% 27% .
2	V	75	 53% 41% 5%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 22777 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Flagellar motor switch protein FliM.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	183	1483	968	231	276	8	0	0	0
1	C	183	1483	968	231	276	8	0	0	0
1	E	183	1483	968	231	276	8	0	0	0
1	G	183	1483	968	231	276	8	0	0	0
1	I	183	1483	968	231	276	8	0	0	0
1	K	183	1483	968	231	276	8	0	0	0
1	M	183	1483	968	231	276	8	0	0	0
1	O	183	1483	968	231	276	8	0	0	0
1	Q	183	1483	968	231	276	8	0	0	0
1	S	183	1483	968	231	276	8	0	0	0
1	U	183	1483	968	231	276	8	0	0	0

- Molecule 2 is a protein called Flagellar motor switch protein FliG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	74	584	375	98	110	1	0	0	0
2	D	75	588	377	99	111	1	0	0	0
2	F	75	588	377	99	111	1	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	75	Total	C	N	O	S	0	0	0
			588	377	99	111	1			
2	J	75	Total	C	N	O	S	0	0	0
			588	377	99	111	1			
2	L	75	Total	C	N	O	S	0	0	0
			588	377	99	111	1			
2	N	75	Total	C	N	O	S	0	0	0
			588	377	99	111	1			
2	P	75	Total	C	N	O	S	0	0	0
			588	377	99	111	1			
2	R	75	Total	C	N	O	S	0	0	0
			588	377	99	111	1			
2	T	75	Total	C	N	O	S	0	0	0
			588	377	99	111	1			
2	V	75	Total	C	N	O	S	0	0	0
			588	377	99	111	1			

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	113	GLY	-	expression tag	UNP Q9WY63
B	114	SER	-	expression tag	UNP Q9WY63
B	115	HIS	-	expression tag	UNP Q9WY63
B	116	MET	-	expression tag	UNP Q9WY63
D	113	GLY	-	expression tag	UNP Q9WY63
D	114	SER	-	expression tag	UNP Q9WY63
D	115	HIS	-	expression tag	UNP Q9WY63
D	116	MET	-	expression tag	UNP Q9WY63
F	113	GLY	-	expression tag	UNP Q9WY63
F	114	SER	-	expression tag	UNP Q9WY63
F	115	HIS	-	expression tag	UNP Q9WY63
F	116	MET	-	expression tag	UNP Q9WY63
H	113	GLY	-	expression tag	UNP Q9WY63
H	114	SER	-	expression tag	UNP Q9WY63
H	115	HIS	-	expression tag	UNP Q9WY63
H	116	MET	-	expression tag	UNP Q9WY63
J	113	GLY	-	expression tag	UNP Q9WY63
J	114	SER	-	expression tag	UNP Q9WY63
J	115	HIS	-	expression tag	UNP Q9WY63
J	116	MET	-	expression tag	UNP Q9WY63
L	113	GLY	-	expression tag	UNP Q9WY63
L	114	SER	-	expression tag	UNP Q9WY63
L	115	HIS	-	expression tag	UNP Q9WY63

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
L	116	MET	-	expression tag	UNP Q9WY63
N	113	GLY	-	expression tag	UNP Q9WY63
N	114	SER	-	expression tag	UNP Q9WY63
N	115	HIS	-	expression tag	UNP Q9WY63
N	116	MET	-	expression tag	UNP Q9WY63
P	113	GLY	-	expression tag	UNP Q9WY63
P	114	SER	-	expression tag	UNP Q9WY63
P	115	HIS	-	expression tag	UNP Q9WY63
P	116	MET	-	expression tag	UNP Q9WY63
R	113	GLY	-	expression tag	UNP Q9WY63
R	114	SER	-	expression tag	UNP Q9WY63
R	115	HIS	-	expression tag	UNP Q9WY63
R	116	MET	-	expression tag	UNP Q9WY63
T	113	GLY	-	expression tag	UNP Q9WY63
T	114	SER	-	expression tag	UNP Q9WY63
T	115	HIS	-	expression tag	UNP Q9WY63
T	116	MET	-	expression tag	UNP Q9WY63
V	113	GLY	-	expression tag	UNP Q9WY63
V	114	SER	-	expression tag	UNP Q9WY63
V	115	HIS	-	expression tag	UNP Q9WY63
V	116	MET	-	expression tag	UNP Q9WY63


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Flagellar motor switch protein FliM




- Molecule 1: Flagellar motor switch protein FliM




- Molecule 1: Flagellar motor switch protein FliM



- Molecule 1: Flagellar motor switch protein FliM








• Molecule 1: Flagellar motor switch protein FliM




• Molecule 1: Flagellar motor switch protein FliM



• Molecule 1: Flagellar motor switch protein FliM



• Molecule 1: Flagellar motor switch protein FliM




• Molecule 1: Flagellar motor switch protein FliM





- Molecule 1: Flagellar motor switch protein FliM




- Molecule 1: Flagellar motor switch protein FliM



- Molecule 2: Flagellar motor switch protein FliG




- Molecule 2: Flagellar motor switch protein FliG



- Molecule 2: Flagellar motor switch protein FliG





- Molecule 2: Flagellar motor switch protein FliG






- Molecule 2: Flagellar motor switch protein FliG




- Molecule 2: Flagellar motor switch protein FliG



- Molecule 2: Flagellar motor switch protein FliG



- Molecule 2: Flagellar motor switch protein FliG



- Molecule 2: Flagellar motor switch protein FliG



- Molecule 2: Flagellar motor switch protein FliG



- Molecule 2: Flagellar motor switch protein FliG





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	105.40Å 216.25Å 262.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	44.18 – 4.32 49.98 – 4.31	Depositor EDS
% Data completeness (in resolution range)	89.7 (44.18-4.32) 88.5 (49.98-4.31)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 4.29Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, R_{free}	0.209 , 0.291 0.213 , 0.289	Depositor DCC
R_{free} test set	2000 reflections (4.92%)	wwPDB-VP
Wilson B-factor (Å ²)	118.5	Xtrriage
Anisotropy	0.874	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 110.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.88	EDS
Total number of atoms	22777	wwPDB-VP
Average B, all atoms (Å ²)	168.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/1521	0.50	0/2069
1	C	0.29	0/1521	0.46	0/2069
1	E	0.27	0/1521	0.46	0/2069
1	G	0.27	0/1521	0.46	0/2069
1	I	0.29	0/1521	0.44	0/2069
1	K	0.27	0/1521	0.47	0/2069
1	M	0.29	0/1521	0.47	0/2069
1	O	0.27	0/1521	0.44	0/2069
1	Q	0.27	0/1521	0.44	0/2069
1	S	0.30	0/1521	0.48	0/2069
1	U	0.27	0/1521	0.49	0/2069
2	B	0.43	0/593	0.53	0/805
2	D	0.26	0/597	0.46	0/810
2	F	0.26	0/597	0.45	0/810
2	H	0.25	0/597	0.49	0/810
2	J	0.26	0/597	0.48	0/810
2	L	0.27	0/597	0.48	0/810
2	N	0.26	0/597	0.44	0/810
2	P	0.26	0/597	0.54	0/810
2	R	0.28	0/597	0.44	0/810
2	T	0.29	0/597	0.52	0/810
2	V	0.25	0/597	0.51	0/810
All	All	0.28	0/23294	0.47	0/31664

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
1	G	0	1

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
All	All	0	3

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	225	GLU	Peptide
1	E	59	HIS	Peptide
1	G	176	ILE	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1483	0	1471	47	0
1	C	1483	0	1471	40	0
1	E	1483	0	1471	38	0
1	G	1483	0	1471	32	0
1	I	1483	0	1471	28	0
1	K	1483	0	1471	28	0
1	M	1483	0	1471	32	0
1	O	1483	0	1471	32	0
1	Q	1483	0	1471	34	0
1	S	1483	0	1471	34	0
1	U	1483	0	1471	39	0
2	B	584	0	610	22	0
2	D	588	0	613	8	0
2	F	588	0	613	12	0
2	H	588	0	613	9	0
2	J	588	0	613	7	0
2	L	588	0	613	18	0
2	N	588	0	613	12	0
2	P	588	0	613	19	0
2	R	588	0	613	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	T	588	0	613	20	0
2	V	588	0	613	23	0
All	All	22777	0	22921	514	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 514 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:48:SER:HB3	1:I:51:GLN:HG2	1.49	0.95
1:E:170:GLN:HG3	1:E:223:LEU:HD11	1.61	0.83
1:A:221:GLU:HA	1:A:224:LEU:HB2	1.63	0.81
1:C:195:VAL:HG22	1:C:216:PRO:HA	1.66	0.78
1:Q:68:THR:HG21	1:S:68:THR:HG21	1.66	0.77

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	181/183 (99%)	164 (91%)	14 (8%)	3 (2%)	9	43
1	C	181/183 (99%)	165 (91%)	11 (6%)	5 (3%)	5	33
1	E	181/183 (99%)	159 (88%)	16 (9%)	6 (3%)	4	30
1	G	181/183 (99%)	159 (88%)	19 (10%)	3 (2%)	9	43
1	I	181/183 (99%)	170 (94%)	9 (5%)	2 (1%)	14	52
1	K	181/183 (99%)	161 (89%)	17 (9%)	3 (2%)	9	43
1	M	181/183 (99%)	163 (90%)	13 (7%)	5 (3%)	5	33
1	O	181/183 (99%)	157 (87%)	19 (10%)	5 (3%)	5	33

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	181/183 (99%)	166 (92%)	10 (6%)	5 (3%)	5	33
1	S	181/183 (99%)	161 (89%)	13 (7%)	7 (4%)	3	26
1	U	181/183 (99%)	158 (87%)	17 (9%)	6 (3%)	4	30
2	B	72/75 (96%)	67 (93%)	3 (4%)	2 (3%)	5	33
2	D	73/75 (97%)	66 (90%)	6 (8%)	1 (1%)	11	47
2	F	73/75 (97%)	69 (94%)	3 (4%)	1 (1%)	11	47
2	H	73/75 (97%)	62 (85%)	9 (12%)	2 (3%)	5	34
2	J	73/75 (97%)	69 (94%)	3 (4%)	1 (1%)	11	47
2	L	73/75 (97%)	66 (90%)	7 (10%)	0	100	100
2	N	73/75 (97%)	69 (94%)	3 (4%)	1 (1%)	11	47
2	P	73/75 (97%)	62 (85%)	7 (10%)	4 (6%)	2	21
2	R	73/75 (97%)	65 (89%)	6 (8%)	2 (3%)	5	34
2	T	73/75 (97%)	67 (92%)	5 (7%)	1 (1%)	11	47
2	V	73/75 (97%)	63 (86%)	7 (10%)	3 (4%)	3	25
All	All	2793/2838 (98%)	2508 (90%)	217 (8%)	68 (2%)	6	36

5 of 68 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	60	GLU
1	C	90	GLU
2	D	181	GLU
1	E	60	GLU
1	G	177	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	170/170 (100%)	160 (94%)	10 (6%)	19	47
1	C	170/170 (100%)	162 (95%)	8 (5%)	26	53

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	E	170/170 (100%)	159 (94%)	11 (6%)	17	44
1	G	170/170 (100%)	159 (94%)	11 (6%)	17	44
1	I	170/170 (100%)	158 (93%)	12 (7%)	14	41
1	K	170/170 (100%)	161 (95%)	9 (5%)	22	50
1	M	170/170 (100%)	160 (94%)	10 (6%)	19	47
1	O	170/170 (100%)	162 (95%)	8 (5%)	26	53
1	Q	170/170 (100%)	161 (95%)	9 (5%)	22	50
1	S	170/170 (100%)	154 (91%)	16 (9%)	8	30
1	U	170/170 (100%)	159 (94%)	11 (6%)	17	44
2	B	67/67 (100%)	64 (96%)	3 (4%)	27	54
2	D	67/67 (100%)	67 (100%)	0	100	100
2	F	67/67 (100%)	65 (97%)	2 (3%)	41	63
2	H	67/67 (100%)	64 (96%)	3 (4%)	27	54
2	J	67/67 (100%)	65 (97%)	2 (3%)	41	63
2	L	67/67 (100%)	64 (96%)	3 (4%)	27	54
2	N	67/67 (100%)	66 (98%)	1 (2%)	65	80
2	P	67/67 (100%)	64 (96%)	3 (4%)	27	54
2	R	67/67 (100%)	63 (94%)	4 (6%)	19	46
2	T	67/67 (100%)	64 (96%)	3 (4%)	27	54
2	V	67/67 (100%)	63 (94%)	4 (6%)	19	46
All	All	2607/2607 (100%)	2464 (94%)	143 (6%)	21	49

5 of 143 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	S	88	THR
1	S	180	GLU
1	U	86	GLN
1	I	53	ARG
2	H	160	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 17 such sidechains are listed below:

Mol	Chain	Res	Type
2	P	115	HIS
2	T	145	GLN
1	E	86	GLN
1	I	157	ASN
1	K	51	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.