



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 17, 2023 – 10:55 PM EST

PDB ID : 2QTQ  
Title : Crystal structure of a predicted dna-binding transcriptional regulator (saro\_1072) from novosphingobium aromaticivorans dsm at 1.85 Å resolution  
Authors : Joint Center for Structural Genomics (JCSG)  
Deposited on : 2007-08-02  
Resolution : 1.85 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.31.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.31.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2469 (1.86-1.86)
Clashscore	141614	2625 (1.86-1.86)
Ramachandran outliers	138981	2592 (1.86-1.86)
Sidechain outliers	138945	2592 (1.86-1.86)
RSRZ outliers	127900	2436 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	 4% 88% 7% . .
1	B	213	 5% 84% 8% . 8%
1	C	213	 11% 83% 9% 8%
1	D	213	 9% 83% 10% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	D	213	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7418 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcriptional regulator, TetR family.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	204	1694	1070	298	319	2	5	0	9	0
1	B	197	1714	1078	314	315	2	5	0	15	0
1	C	197	1596	1017	283	288	3	5	0	5	0
1	D	198	1620	1033	288	291	2	6	0	6	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	expression tag	UNP Q2G9F6
B	0	GLY	-	expression tag	UNP Q2G9F6
C	0	GLY	-	expression tag	UNP Q2G9F6
D	0	GLY	-	expression tag	UNP Q2G9F6

- Molecule 2 is NITRATE ION (three-letter code: NO3) (formula: NO<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	N	O	0	0
			4	1	3		
2	B	1	Total	N	O	0	0
			4	1	3		
2	C	1	Total	N	O	0	0
			4	1	3		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	B	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	C	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0
3	D	1	Total C O 4 2 2	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	205	Total	O	0	4
			209	209		
4	B	209	Total	O	0	3
			212	212		
4	C	145	Total	O	0	2
			147	147		
4	D	113	Total	O	0	5
			118	118		



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: Transcriptional regulator, TetR family





- Molecule 1: Transcriptional regulator, TetR family



- Molecule 1: Transcriptional regulator, TetR family



- Molecule 1: Transcriptional regulator, TetR family





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	66.47Å 98.76Å 164.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.50 – 1.85 29.50 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.0 (29.50-1.85) 98.0 (29.50-1.85)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	0.06	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.96 (at 1.85Å)	Xtrriage
Refinement program	REFMAC 5.2.0019, PHENIX	Depositor
R, $R_{free}$	0.185 , 0.216 0.188 , 0.216	Depositor DCC
$R_{free}$ test set	4544 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	31.5	Xtrriage
Anisotropy	0.524	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 51.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7418	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NO3, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.63	0/1718	0.70	1/2312 (0.0%)
1	B	0.69	1/1735 (0.1%)	0.73	0/2326
1	C	0.58	1/1618 (0.1%)	0.62	0/2173
1	D	0.50	0/1643	0.62	1/2207 (0.0%)
All	All	0.60	2/6714 (0.0%)	0.67	2/9018 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	28	MSE	SE-CE	-9.64	1.38	1.95
1	C	28	MSE	SE-CE	-6.67	1.56	1.95

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	198	ARG	NE-CZ-NH1	5.40	123.00	120.30
1	D	115	MSE	CG-SE-CE	-5.17	87.52	98.90

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1694	0	1655	13	0
1	B	1714	0	1713	16	0
1	C	1596	0	1593	19	0
1	D	1620	0	1613	15	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
3	A	40	0	60	2	0
3	B	28	0	42	1	0
3	C	12	0	18	2	0
3	D	16	0	24	0	0
4	A	209	0	0	6	0
4	B	212	0	0	3	0
4	C	147	0	0	2	0
4	D	118	0	0	2	0
All	All	7418	0	6718	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (63) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:87:MSE:HE2	1:B:91:ALA:HB1	1.39	1.00
1:C:34:VAL:HG12	1:C:34:VAL:O	1.74	0.84
1:B:87:MSE:HE2	1:B:91:ALA:CB	2.08	0.83
1:A:172:ARG:HD3	1:B:165:THR:HA	1.66	0.77
1:B:87:MSE:CE	1:B:91:ALA:HB1	2.18	0.71
1:C:115:MSE:HE3	1:C:119:ARG:HH21	1.56	0.70
1:C:115:MSE:HE2	4:C:347:HOH:O	1.92	0.70
1:C:115:MSE:CE	1:C:119:ARG:HH21	2.04	0.69
1:D:87[A]:MSE:HE1	1:D:95[A]:ARG:NH1	2.14	0.63
1:D:90:GLU:HG3	1:D:206:MSE:HE2	1.79	0.63
1:D:90:GLU:HG3	1:D:206:MSE:CE	2.30	0.61
1:C:115:MSE:HE3	1:C:119:ARG:NH2	2.13	0.61
1:D:87[A]:MSE:HE1	1:D:95[A]:ARG:HH11	1.65	0.61
1:C:34:VAL:O	1:C:34:VAL:CG1	2.47	0.60
1:D:115:MSE:HE1	1:D:119:ARG:CZ	2.32	0.59
1:D:35:ASP:OD1	1:D:59[B]:LYS:HE3	2.02	0.58
1:B:15:GLY:N	3:B:220:EDO:HO1	2.01	0.58
1:B:33:VAL:CG1	4:B:274:HOH:O	2.51	0.58
1:D:211:ALA:HA	1:D:212:HIS:HB2	1.86	0.58

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:318:HOH:O	1:B:161:LEU:HD22	2.05	0.57
1:D:81:LEU:HD13	1:D:95[A]:ARG:NE	2.23	0.54
1:C:112:ARG:CD	3:C:215:EDO:O1	2.56	0.54
3:A:215:EDO:H11	4:A:243:HOH:O	2.08	0.52
1:B:195:ASP:OD1	1:B:198[B]:ARG:NH2	2.44	0.51
1:A:212:HIS:ND1	4:A:344:HOH:O	2.34	0.50
1:D:81:LEU:HD13	1:D:95[A]:ARG:CZ	2.42	0.49
1:C:145:ILE:O	1:C:149:VAL:HG13	2.12	0.49
1:B:33:VAL:HG13	4:B:274:HOH:O	2.12	0.49
1:B:33:VAL:HG12	4:B:274:HOH:O	2.11	0.48
1:C:90:GLU:OE2	1:C:94:ARG:NH1	2.46	0.48
1:A:172:ARG:NH1	4:A:285:HOH:O	2.22	0.48
1:B:143:ARG:O	1:B:147:GLU:HG3	2.13	0.48
1:C:182:HIS:HD2	4:D:251:HOH:O	1.96	0.48
1:C:146:GLY:HA2	1:C:149:VAL:HG22	1.97	0.47
1:D:47:LEU:HB3	1:D:51:LEU:HD12	1.97	0.47
1:C:206:MSE:HA	1:C:209:ILE:HG22	1.97	0.46
1:D:181:LYS:NZ	4:D:303:HOH:O	2.46	0.46
1:A:102[B]:ASP:OD1	1:A:198:ARG:HD3	2.15	0.46
1:C:112:ARG:HD2	3:C:215:EDO:O1	2.15	0.46
1:C:157:ILE:CD1	1:C:209:ILE:HD13	2.46	0.45
1:A:80:ALA:HB2	4:A:274:HOH:O	2.16	0.45
1:C:115:MSE:CE	1:C:119:ARG:NH2	2.73	0.45
1:C:115:MSE:HE1	1:C:119:ARG:HH21	1.81	0.45
1:B:85:ASP:O	1:B:86:ASP:CB	2.65	0.44
1:C:34:VAL:HG13	1:C:113:LEU:HD13	1.99	0.44
1:D:67:LEU:HD22	1:D:117:LEU:HD12	2.00	0.43
1:B:33:VAL:HG22	1:B:35[A]:ASP:H	1.83	0.43
1:D:26:ASN:O	1:D:30[A]:GLU:HG2	2.19	0.43
1:B:28:MSE:HE2	1:B:109:TYR:CG	2.54	0.42
1:B:35[A]:ASP:OD1	1:B:59:LYS:NZ	2.40	0.42
1:D:206:MSE:HA	1:D:209:ILE:HG22	2.02	0.42
1:A:67:LEU:HD22	1:A:117[A]:LEU:HD12	2.02	0.42
1:D:187:ASP:O	1:D:193:LEU:HD12	2.20	0.42
1:B:135:LEU:HD13	1:B:139[A]:ARG:NH2	2.35	0.42
1:A:117[B]:LEU:N	1:A:117[B]:LEU:HD12	2.36	0.41
1:A:127:LYS:NZ	1:A:131:ASP:OD2	2.52	0.41
1:A:172:ARG:HD2	4:A:247:HOH:O	2.21	0.41
1:A:195:ASP:O	1:A:199:GLU:HG2	2.20	0.41
1:C:86:ASP:CG	1:C:87:MSE:HE3	2.41	0.41
1:C:116[B]:ARG:HG3	4:C:347:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9[B]:ASP:OD1	1:A:9[B]:ASP:N	2.54	0.40
1:A:19:LEU:HD11	3:A:217:EDO:H12	2.04	0.40
1:A:34:VAL:HG21	1:A:112:ARG:HB3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	210/213 (99%)	209 (100%)	1 (0%)	0	100	100
1	B	209/213 (98%)	204 (98%)	5 (2%)	0	100	100
1	C	200/213 (94%)	197 (98%)	3 (2%)	0	100	100
1	D	202/213 (95%)	199 (98%)	2 (1%)	1 (0%)	29	15
All	All	821/852 (96%)	809 (98%)	11 (1%)	1 (0%)	51	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	84	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/177 (100%)	174 (98%)	3 (2%)	60	47
1	B	182/177 (103%)	180 (99%)	2 (1%)	73	65
1	C	164/177 (93%)	162 (99%)	2 (1%)	71	62
1	D	167/177 (94%)	167 (100%)	0	100	100
All	All	690/708 (98%)	683 (99%)	7 (1%)	78	69

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9[A]	ASP
1	A	9[B]	ASP
1	A	189	LEU
1	B	79	ASP
1	B	195	ASP
1	C	143	ARG
1	C	171	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

27 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	EDO	D	213	-	3,3,3	0.59	0	2,2,2	0.15	0
3	EDO	A	222	-	3,3,3	0.42	0	2,2,2	0.33	0
3	EDO	B	220	-	3,3,3	0.44	0	2,2,2	0.36	0
3	EDO	B	218	-	3,3,3	0.53	0	2,2,2	0.15	0
3	EDO	A	221	-	3,3,3	0.46	0	2,2,2	0.29	0
3	EDO	B	219	-	3,3,3	0.59	0	2,2,2	0.05	0
2	NO3	C	213	-	1,3,3	3.53	1 (100%)	0,3,3	-	-
2	NO3	B	213	-	1,3,3	3.15	1 (100%)	0,3,3	-	-
3	EDO	C	215	-	3,3,3	0.53	0	2,2,2	0.28	0
3	EDO	B	217	-	3,3,3	0.57	0	2,2,2	0.18	0
3	EDO	A	220	-	3,3,3	0.48	0	2,2,2	0.36	0
3	EDO	D	215	-	3,3,3	0.45	0	2,2,2	0.17	0
2	NO3	A	213	-	1,3,3	3.83	1 (100%)	0,3,3	-	-
3	EDO	B	215	-	3,3,3	0.48	0	2,2,2	0.19	0
3	EDO	A	218	-	3,3,3	0.66	0	2,2,2	0.37	0
3	EDO	A	215	-	3,3,3	0.58	0	2,2,2	0.31	0
3	EDO	B	216	-	3,3,3	0.64	0	2,2,2	0.27	0
3	EDO	C	216	-	3,3,3	0.48	0	2,2,2	0.19	0
3	EDO	A	216	-	3,3,3	0.49	0	2,2,2	0.44	0
3	EDO	D	216	-	3,3,3	0.51	0	2,2,2	0.08	0
3	EDO	D	214	-	3,3,3	0.55	0	2,2,2	0.21	0
3	EDO	A	219	-	3,3,3	0.54	0	2,2,2	0.35	0
3	EDO	B	214	-	3,3,3	0.73	0	2,2,2	0.05	0
3	EDO	C	214	-	3,3,3	0.51	0	2,2,2	0.23	0
3	EDO	A	214	-	3,3,3	0.50	0	2,2,2	0.18	0
3	EDO	A	223	-	3,3,3	0.36	0	2,2,2	0.50	0
3	EDO	A	217	-	3,3,3	0.37	0	2,2,2	0.56	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	D	213	-	-	0/1/1/1	-
3	EDO	A	222	-	-	1/1/1/1	-
3	EDO	B	220	-	-	1/1/1/1	-

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	218	-	-	0/1/1/1	-
3	EDO	A	221	-	-	1/1/1/1	-
3	EDO	B	219	-	-	1/1/1/1	-
3	EDO	C	215	-	-	0/1/1/1	-
3	EDO	B	217	-	-	1/1/1/1	-
3	EDO	A	220	-	-	1/1/1/1	-
3	EDO	D	215	-	-	0/1/1/1	-
3	EDO	B	215	-	-	1/1/1/1	-
3	EDO	A	218	-	-	1/1/1/1	-
3	EDO	A	215	-	-	0/1/1/1	-
3	EDO	B	216	-	-	1/1/1/1	-
3	EDO	C	216	-	-	1/1/1/1	-
3	EDO	A	216	-	-	0/1/1/1	-
3	EDO	D	216	-	-	1/1/1/1	-
3	EDO	D	214	-	-	0/1/1/1	-
3	EDO	A	219	-	-	1/1/1/1	-
3	EDO	B	214	-	-	1/1/1/1	-
3	EDO	C	214	-	-	0/1/1/1	-
3	EDO	A	214	-	-	1/1/1/1	-
3	EDO	A	223	-	-	0/1/1/1	-
3	EDO	A	217	-	-	1/1/1/1	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	213	NO3	O1-N	3.83	1.41	1.24
2	C	213	NO3	O1-N	3.53	1.40	1.24
2	B	213	NO3	O1-N	3.15	1.38	1.24

There are no bond angle outliers.

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	D	216	EDO	O1-C1-C2-O2
3	A	218	EDO	O1-C1-C2-O2
3	A	221	EDO	O1-C1-C2-O2
3	A	222	EDO	O1-C1-C2-O2
3	B	214	EDO	O1-C1-C2-O2
3	B	217	EDO	O1-C1-C2-O2
3	A	214	EDO	O1-C1-C2-O2

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
3	A	217	EDO	O1-C1-C2-O2
3	A	220	EDO	O1-C1-C2-O2
3	B	216	EDO	O1-C1-C2-O2
3	B	220	EDO	O1-C1-C2-O2
3	C	216	EDO	O1-C1-C2-O2
3	A	219	EDO	O1-C1-C2-O2
3	B	215	EDO	O1-C1-C2-O2
3	B	219	EDO	O1-C1-C2-O2

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	220	EDO	1	0
3	C	215	EDO	2	0
3	A	215	EDO	1	0
3	A	217	EDO	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	199/213 (93%)	0.07	8 (4%) 38 36	22, 34, 46, 57	0
1	B	192/213 (90%)	0.15	11 (5%) 23 23	26, 35, 50, 67	0
1	C	192/213 (90%)	0.39	23 (11%) 4 4	30, 38, 49, 55	0
1	D	193/213 (90%)	0.43	20 (10%) 6 6	31, 40, 51, 65	0
All	All	776/852 (91%)	0.26	62 (7%) 12 12	22, 37, 50, 67	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	83	ALA	4.9
1	B	166[A]	VAL	4.8
1	C	152	GLY	4.6
1	D	212	HIS	4.6
1	D	153	VAL	4.5
1	C	149	VAL	4.4
1	C	153	VAL	4.4
1	D	152	GLY	4.2
1	D	210	LEU	4.2
1	D	172[A]	ARG	4.1
1	C	124	ALA	3.9
1	D	211	ALA	3.9
1	C	169	ALA	3.9
1	C	83	ALA	3.8
1	D	80	ALA	3.7
1	C	85	ASP	3.7
1	B	165	THR	3.4
1	D	151	ALA	3.4
1	C	84	LYS	3.3
1	C	80	ALA	3.2
1	A	172	ARG	3.2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	84	LYS	3.2
1	D	169	ALA	3.1
1	A	101	ILE	3.1
1	C	165	THR	3.1
1	C	150	LYS	3.1
1	C	81	LEU	3.1
1	A	169	ALA	3.1
1	B	85	ASP	2.9
1	B	168	GLY	2.9
1	C	168	GLY	2.9
1	B	205	ILE	2.9
1	A	86	ASP	2.8
1	C	166	VAL	2.8
1	C	210	LEU	2.8
1	C	167	THR	2.8
1	C	170	ALA	2.7
1	C	151	ALA	2.7
1	A	167	THR	2.7
1	D	54	TYR	2.7
1	C	163	TYR	2.6
1	C	156	PRO	2.6
1	B	164	PHE	2.6
1	C	211	ALA	2.6
1	D	86	ASP	2.6
1	D	89	PRO	2.6
1	D	95[A]	ARG	2.6
1	D	184	PHE	2.5
1	B	169	ALA	2.5
1	C	86	ASP	2.5
1	A	166	VAL	2.4
1	B	167	THR	2.3
1	D	85	ASP	2.3
1	B	86	ASP	2.2
1	A	205	ILE	2.2
1	B	161	LEU	2.2
1	B	84	LYS	2.2
1	D	185	ASP	2.1
1	D	168	GLY	2.1
1	C	88	SER	2.1
1	D	205	ILE	2.1
1	A	204	PHE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	B	217	4/4	0.62	0.27	73,73,75,75	0
3	EDO	B	219	4/4	0.66	0.19	69,69,69,70	0
3	EDO	A	214	4/4	0.73	0.38	60,64,65,66	0
3	EDO	A	219	4/4	0.74	0.20	61,63,67,70	0
3	EDO	A	221	4/4	0.74	0.18	81,81,82,82	0
3	EDO	A	223	4/4	0.76	0.27	65,67,67,74	0
3	EDO	B	216	4/4	0.76	0.24	56,68,70,71	0
3	EDO	A	215	4/4	0.78	0.37	56,62,67,71	0
3	EDO	C	216	4/4	0.78	0.17	64,67,70,73	0
3	EDO	A	218	4/4	0.79	0.27	45,57,58,59	0
3	EDO	D	213	4/4	0.79	0.41	62,65,66,71	0
3	EDO	D	215	4/4	0.81	0.25	58,61,66,71	0
3	EDO	A	220	4/4	0.82	0.18	68,70,71,73	0
2	NO3	A	213	4/4	0.83	0.23	59,63,64,64	0
3	EDO	B	218	4/4	0.84	0.19	59,63,65,70	0
3	EDO	A	216	4/4	0.85	0.19	51,56,58,59	0
3	EDO	C	214	4/4	0.86	0.42	71,71,72,73	0
2	NO3	C	213	4/4	0.87	0.19	67,68,68,68	0
3	EDO	B	214	4/4	0.87	0.29	49,52,53,54	0
3	EDO	A	217	4/4	0.88	0.15	53,58,59,65	0
3	EDO	B	215	4/4	0.88	0.31	68,70,73,73	0
3	EDO	D	216	4/4	0.89	0.20	56,67,70,73	0
3	EDO	C	215	4/4	0.92	0.17	45,46,48,48	0
3	EDO	B	220	4/4	0.94	0.29	63,68,68,73	0
3	EDO	D	214	4/4	0.95	0.09	49,51,54,55	0
3	EDO	A	222	4/4	0.96	0.09	33,39,44,45	0
2	NO3	B	213	4/4	0.97	0.12	43,44,44,44	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.