



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 19, 2023 – 07:16 pm BST

PDB ID : 8QT5
Title : Crystal structure of Arabidopsis thaliana 14-3-3 isoform lambda in complex with a phosphopeptide from the transcription factor BZR1.
Authors : Hothorn, M.; Obergfell, E.
Deposited on : 2023-10-12
Resolution : 2.69 Å (reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.69 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	255	 4% 91% 6%
1	B	255	 4% 92% 5%
1	C	255	 2% 93%
1	D	255	 2% 93%
1	E	255	 5% 91% 5%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	255	 <p>17% 87% 9%</p>
1	G	255	 <p>60% 86% 9% 5%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 27061 atoms, of which 13407 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 14-3-3-like protein G-BOX factor 14 lambda, Protein BRASSI NAZOLE-RESISTANT 1.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
			Total	C	H	N	O	P	S			
1	A	248	3900	1230	1933	326	397	1	13	0	1	0
1	B	247	3877	1222	1922	326	394	1	12	0	0	0
1	C	246	3862	1217	1916	325	391	1	12	0	0	0
1	D	247	3870	1221	1917	324	395	1	12	0	0	0
1	E	247	3885	1226	1927	324	395	1	12	0	1	0
1	F	245	3839	1211	1904	321	390	1	12	0	0	0
1	G	242	3792	1197	1882	317	384	1	11	0	0	0

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1174	ALA	CYS	engineered mutation	UNP Q8S307
B	1174	ALA	CYS	engineered mutation	UNP Q8S307
C	1174	ALA	CYS	engineered mutation	UNP Q8S307
D	1174	ALA	CYS	engineered mutation	UNP Q8S307
E	1174	ALA	CYS	engineered mutation	UNP Q8S307
F	1174	ALA	CYS	engineered mutation	UNP Q8S307
G	1174	ALA	CYS	engineered mutation	UNP Q8S307

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			7	2	3	2		
2	F	1	Total	C	H	O	0	0
			7	2	3	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	7	Total	O	0	0
			7	7		
3	B	6	Total	O	0	0
			6	6		
3	C	2	Total	O	0	0
			2	2		
3	D	1	Total	O	0	0
			1	1		
3	E	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 14-3-3-like protein G-BOX factor 14 lambda, Protein BRASSINAZOLE-RESISTANT 1




- Molecule 1: 14-3-3-like protein G-BOX factor 14 lambda, Protein BRASSINAZOLE-RESISTANT 1



- Molecule 1: 14-3-3-like protein G-BOX factor 14 lambda, Protein BRASSINAZOLE-RESISTANT 1




- Molecule 1: 14-3-3-like protein G-BOX factor 14 lambda, Protein BRASSINAZOLE-RESISTANT 1






- Molecule 1: 14-3-3-like protein G-BOX factor 14 lambda, Protein BRASSINAZOLE-RESISTANT 1










- Molecule 1: 14-3-3-like protein G-BOX factor 14 lambda, Protein BRASSINAZOLE-RESISTANT 1



- Molecule 1: 14-3-3-like protein G-BOX factor 14 lambda, Protein BRASSINAZOLE-RESISTANT 1



4 Data and refinement statistics i

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	176.05Å 176.05Å 172.03Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	74.92 – 2.69 78.36 – 2.69	Depositor EDS
% Data completeness (in resolution range)	98.3 (74.92-2.69) 98.5 (78.36-2.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.18 (at 2.69Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, R_{free}	0.241 , 0.262 0.243 , 0.264	Depositor DCC
R_{free} test set	4235 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	86.3	Xtrriage
Anisotropy	0.268	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 57.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.018 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	27061	wwPDB-VP
Average B, all atoms (Å ²)	128.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1988	0.43	0/2676
1	B	0.24	0/1973	0.43	0/2656
1	C	0.24	0/1964	0.42	0/2644
1	D	0.24	0/1971	0.42	0/2654
1	E	0.25	0/1979	0.43	0/2665
1	F	0.25	0/1953	0.44	0/2630
1	G	0.24	0/1928	0.42	0/2597
All	All	0.25	0/13756	0.43	0/18522

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1967	1933	1933	4	1
1	B	1955	1922	1921	6	0
1	C	1946	1916	1914	5	0
1	D	1953	1917	1916	5	0
1	E	1958	1927	1927	6	1
1	F	1935	1904	1901	13	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	1910	1882	1880	12	0
2	A	4	3	3	0	0
2	F	4	3	3	0	0
3	A	7	0	0	1	0
3	B	6	0	0	0	0
3	C	2	0	0	0	0
3	D	1	0	0	0	0
3	E	6	0	0	0	0
All	All	13654	13407	13398	48	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 48 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:125:VAL:HG21	1:D:169:MET:SD	2.16	0.85
1:F:236:LEU:O	1:G:231:ARG:NH1	2.25	0.69
1:G:92:ARG:NH1	1:G:96:GLU:OE2	2.25	0.69
1:E:125:VAL:HG21	1:E:169:MET:SD	2.36	0.65
1:A:146:GLU:HB2	3:A:1302:HOH:O	1.99	0.62

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:ASP:OD1	1:E:79:ARG:NH1[3_454]	2.11	0.09

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/255 (96%)	237 (97%)	7 (3%)	0	100	100
1	B	242/255 (95%)	239 (99%)	3 (1%)	0	100	100
1	C	241/255 (94%)	236 (98%)	5 (2%)	0	100	100
1	D	242/255 (95%)	239 (99%)	3 (1%)	0	100	100
1	E	243/255 (95%)	239 (98%)	4 (2%)	0	100	100
1	F	240/255 (94%)	235 (98%)	5 (2%)	0	100	100
1	G	237/255 (93%)	233 (98%)	4 (2%)	0	100	100
All	All	1689/1785 (95%)	1658 (98%)	31 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	212/215 (99%)	204 (96%)	8 (4%)	33	62
1	B	210/215 (98%)	208 (99%)	2 (1%)	76	91
1	C	209/215 (97%)	208 (100%)	1 (0%)	88	96
1	D	210/215 (98%)	210 (100%)	0	100	100
1	E	211/215 (98%)	208 (99%)	3 (1%)	67	86
1	F	208/215 (97%)	206 (99%)	2 (1%)	76	91
1	G	205/215 (95%)	201 (98%)	4 (2%)	55	81
All	All	1465/1505 (97%)	1445 (99%)	20 (1%)	67	86

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	196	LYS
1	G	81	ASN
1	G	158	TYR
1	G	86	SER
1	A	241	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	81	ASN
1	A	244	GLN
1	B	81	ASN
1	G	81	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

7 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	SEP	E	1173	1	8,9,10	1.54	2 (25%)	8,12,14	2.19	2 (25%)
1	SEP	G	1173	1	8,9,10	1.56	1 (12%)	8,12,14	1.59	2 (25%)
1	SEP	A	1173	1	8,9,10	1.64	1 (12%)	8,12,14	2.02	2 (25%)
1	SEP	C	1173	1	8,9,10	1.55	1 (12%)	8,12,14	1.98	2 (25%)
1	SEP	F	1173	1	8,9,10	1.59	1 (12%)	8,12,14	1.73	2 (25%)
1	SEP	D	1173	1	8,9,10	1.60	1 (12%)	8,12,14	2.06	2 (25%)
1	SEP	B	1173	1	8,9,10	1.63	1 (12%)	8,12,14	1.99	2 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	E	1173	1	-	0/5/8/10	-
1	SEP	G	1173	1	-	0/5/8/10	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SEP	A	1173	1	-	0/5/8/10	-
1	SEP	C	1173	1	-	0/5/8/10	-
1	SEP	F	1173	1	-	0/5/8/10	-
1	SEP	D	1173	1	-	0/5/8/10	-
1	SEP	B	1173	1	-	0/5/8/10	-

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1173	SEP	P-O1P	3.68	1.62	1.50
1	A	1173	SEP	P-O1P	3.61	1.62	1.50
1	D	1173	SEP	P-O1P	3.52	1.61	1.50
1	F	1173	SEP	P-O1P	3.49	1.61	1.50
1	G	1173	SEP	P-O1P	3.40	1.61	1.50

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	1173	SEP	OG-CB-CA	4.40	112.43	108.14
1	E	1173	SEP	OG-CB-CA	4.22	112.25	108.14
1	D	1173	SEP	OG-CB-CA	4.19	112.22	108.14
1	B	1173	SEP	P-OG-CB	-4.19	106.75	118.30
1	A	1173	SEP	OG-CB-CA	3.90	111.94	108.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	E	1173	SEP	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ACT	A	1201	-	3,3,3	1.16	0	3,3,3	1.41	0
2	ACT	F	1201	-	3,3,3	1.15	0	3,3,3	1.47	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/255 (96%)	0.77	11 (4%) 33 31	64, 84, 130, 160	0
1	B	246/255 (96%)	0.63	11 (4%) 33 31	70, 93, 146, 221	0
1	C	245/255 (96%)	0.53	5 (2%) 65 67	74, 91, 148, 187	0
1	D	246/255 (96%)	0.49	4 (1%) 72 74	82, 104, 147, 176	0
1	E	246/255 (96%)	0.83	13 (5%) 26 25	69, 91, 135, 195	0
1	F	244/255 (95%)	1.01	43 (17%) 1 1	68, 120, 166, 220	0
1	G	241/255 (94%)	3.31	152 (63%) 0 0	143, 213, 253, 294	0
All	All	1715/1785 (96%)	1.07	239 (13%) 2 2	64, 101, 221, 294	0

The worst 5 of 239 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	79	ARG	15.5
1	G	80	LYS	11.1
1	G	91	TYR	10.8
1	G	127	TYR	10.8
1	G	204	ALA	10.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SEP	G	1173	10/11	0.86	0.15	173,186,215,215	0
1	SEP	D	1173	10/11	0.89	0.21	80,87,106,108	0
1	SEP	E	1173	10/11	0.91	0.25	75,82,99,104	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
1	SEP	F	1173	10/11	0.93	0.19	81,94,115,120	0
1	SEP	A	1173	10/11	0.93	0.26	62,76,93,99	0
1	SEP	B	1173	10/11	0.94	0.20	79,81,100,102	0
1	SEP	C	1173	10/11	0.94	0.18	75,78,97,98	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ACT	F	1201	4/4	0.82	0.53	103,112,136,136	0
2	ACT	A	1201	4/4	0.85	0.38	89,91,108,108	0

6.5 Other polymers [i](#)

There are no such residues in this entry.