



wwPDB X-ray Structure Validation Summary Report ⓘ

May 14, 2020 – 04:29 am BST

PDB ID : 1QUN
Title : X-RAY STRUCTURE OF THE FIMC-FIMH CHAPERONE ADHESIN
COMPLEX FROM UROPATHOGENIC E.COLI
Authors : Choudhury, D.; Thompson, A.; Stojanoff, V.; Langerman, S.; Pinkner, J.;
Hultgren, S.J.; Knight, S.
Deposited on : 1999-07-01
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	205	51% 35% 11% ..
1	C	205	53% 34% 9% ..
1	E	205	53% 34% 9% ..
1	G	205	52% 35% 9% ..
1	I	205	29% 41% 21% 5% .
1	K	205	30% 41% 20% 6% .
1	M	205	30% 40% 21% 5% .
1	O	205	28% 43% 21% . .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	279	 50% 35% 8% 8%
2	D	279	 49% 35% 8% 7%
2	F	279	 49% 36% 8% 8%
2	H	279	 50% 35% 8% 7%
2	J	279	 49% 32% 15% .
2	L	279	 48% 33% 16% .
2	N	279	 48% 33% 15% .
2	P	279	 48% 33% 16% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 28864 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PAPD-LIKE CHAPERONE FIMC.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	200	1553	985	268	294	6	5	0	0
1	C	200	1553	985	268	294	6	5	0	0
1	E	200	1553	985	268	294	6	5	0	0
1	G	200	1553	985	268	294	6	5	0	0
1	I	198	1543	979	266	292	6	177	0	0
1	K	198	1543	979	266	292	6	176	0	0
1	M	198	1543	979	266	292	6	177	0	0
1	O	198	1543	979	266	292	6	177	0	0

- Molecule 2 is a protein called MANNOSE-SPECIFIC ADHESIN FIMH.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	279	2052	1297	342	409	4	42	0	0
2	D	279	2052	1297	342	409	4	42	0	0
2	F	279	2052	1297	342	409	4	42	0	0
2	H	279	2052	1297	342	409	4	42	0	0
2	J	279	2052	1297	342	409	4	67	0	0
2	L	279	2052	1297	342	409	4	67	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	N	279	Total	C	N	O	S	67	0	0
			2052	1297	342	409	4			
2	P	279	Total	C	N	O	S	67	0	0
			2052	1297	342	409	4			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	5	Total	O	0	0
			5	5		
3	B	10	Total	O	0	0
			10	10		
3	C	6	Total	O	0	0
			6	6		
3	D	11	Total	O	0	0
			11	11		
3	E	5	Total	O	0	0
			5	5		
3	F	12	Total	O	0	0
			12	12		
3	G	5	Total	O	0	0
			5	5		
3	H	10	Total	O	0	0
			10	10		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: PAPD-LIKE CHAPERONE FIMC

Chain A: 




- Molecule 1: PAPD-LIKE CHAPERONE FIMC


Chain C: 




- Molecule 1: PAPD-LIKE CHAPERONE FIMC

Chain E: 








• Molecule 1: PAPD-LIKE CHAPERONE FIMC




• Molecule 1: PAPD-LIKE CHAPERONE FIMC




• Molecule 1: PAPD-LIKE CHAPERONE FIMC




• Molecule 1: PAPD-LIKE CHAPERONE FIMC

Chain M: 




• Molecule 1: PAPD-LIKE CHAPERONE FIMC

Chain O: 




• Molecule 2: MANNOSE-SPECIFIC ADHESIN FIMH

Chain B: 




• Molecule 2: MANNOSE-SPECIFIC ADHESIN FIMH


Chain D: 



• Molecule 2: MANNOSE-SPECIFIC ADHESIN FIMH




• Molecule 2: MANNOSE-SPECIFIC ADHESIN FIMH




• Molecule 2: MANNOSE-SPECIFIC ADHESIN FIMH






• Molecule 2: MANNOSE-SPECIFIC ADHESIN FIMH




• Molecule 2: MANNOSE-SPECIFIC ADHESIN FIMH



• Molecule 2: MANNOSE-SPECIFIC ADHESIN FIMH





4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	139.41Å 139.57Å 215.72Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 2.80	Depositor
% Data completeness (in resolution range)	(Not available) (20.00-2.80)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	REFMAC	Depositor
R, R_{free}	0.240 , 0.268	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	28864	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.92	3/1580 (0.2%)	2.40	60/2146 (2.8%)
1	C	0.94	3/1580 (0.2%)	2.43	65/2146 (3.0%)
1	E	0.94	3/1580 (0.2%)	2.43	65/2146 (3.0%)
1	G	0.92	3/1580 (0.2%)	2.40	60/2146 (2.8%)
1	I	2.03	14/1568 (0.9%)	2.88	63/2126 (3.0%)
1	K	2.17	13/1568 (0.8%)	2.87	63/2126 (3.0%)
1	M	2.03	14/1568 (0.9%)	2.88	64/2126 (3.0%)
1	O	2.17	13/1568 (0.8%)	2.87	63/2126 (3.0%)
2	B	1.44	6/2097 (0.3%)	2.57	116/2881 (4.0%)
2	D	1.39	7/2097 (0.3%)	2.57	117/2881 (4.1%)
2	F	1.39	7/2097 (0.3%)	2.57	117/2881 (4.1%)
2	H	1.44	6/2097 (0.3%)	2.57	116/2881 (4.0%)
2	J	1.14	6/2096 (0.3%)	2.54	63/2878 (2.2%)
2	L	1.20	4/2097 (0.2%)	2.45	56/2881 (1.9%)
2	N	1.14	6/2096 (0.3%)	2.54	64/2878 (2.2%)
2	P	1.20	4/2097 (0.2%)	2.45	56/2881 (1.9%)
All	All	1.45	112/29366 (0.4%)	2.58	1208/40130 (3.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	C	0	4
1	E	0	4
1	G	0	4
1	I	1	9
1	K	0	10
1	M	1	9
1	O	0	10
2	B	0	7

Continued on next page...

Continued from previous page...

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	1	7
2	F	1	7
2	H	0	7
2	J	1	4
2	L	1	3
2	N	1	4
2	P	1	3
All	All	8	96

The worst 5 of 112 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	I	26	ASP	CG-OD1	43.75	2.25	1.25
1	M	26	ASP	CG-OD1	43.74	2.25	1.25
1	K	26	ASP	CG-OD1	40.57	2.18	1.25
1	O	26	ASP	CG-OD1	40.57	2.18	1.25
2	L	167	ASP	C-O	-35.59	0.55	1.23

The worst 5 of 1208 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	J	165	ALA	O-C-N	-61.51	24.29	122.70
2	N	165	ALA	O-C-N	-61.50	24.30	122.70
1	K	26	ASP	CB-CG-OD1	-61.39	63.05	118.30
1	O	26	ASP	CB-CG-OD1	-61.38	63.06	118.30
2	L	165	ALA	O-C-N	-60.60	25.74	122.70

5 of 8 chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	D	176	PRO	CA
2	F	176	PRO	CA
1	I	156	ASN	CA
2	J	215	PHE	CA
2	L	215	PHE	CA

5 of 96 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	25	ASN	Mainchain
1	A	45	ASP	Mainchain
1	A	51	THR	Mainchain,Peptide

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
2	B	24	LEU	Mainchain
2	B	35	VAL	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1553	0	1587	48	0
1	C	1553	0	1587	42	0
1	E	1553	0	1587	41	0
1	G	1553	0	1587	44	0
1	I	1543	0	1576	110	0
1	K	1543	0	1576	107	0
1	M	1543	0	1576	109	0
1	O	1543	0	1576	105	0
2	B	2052	0	2006	83	0
2	D	2052	0	2006	80	0
2	F	2052	0	2006	82	0
2	H	2052	0	2006	81	0
2	J	2052	0	2003	114	0
2	L	2052	0	2004	118	0
2	N	2052	0	2003	114	0
2	P	2052	0	2004	114	0
3	A	5	0	0	0	0
3	B	10	0	0	1	0
3	C	6	0	0	0	0
3	D	11	0	0	1	0
3	E	5	0	0	0	0
3	F	12	0	0	1	0
3	G	5	0	0	0	0
3	H	10	0	0	1	0
All	All	28864	0	28690	1311	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 24.

The worst 5 of 1311 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:O:197:THR:HB	1:O:198:PRO:HD2	1.45	0.98
2:N:53:THR:H	2:N:136:ASN:HD21	1.08	0.98
2:J:53:THR:H	2:J:136:ASN:HD21	1.08	0.98
2:B:226:THR:HG22	2:B:253:THR:HB	1.45	0.98
1:I:197:THR:HB	1:I:198:PRO:HD2	1.46	0.98

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	196/205 (96%)	185 (94%)	11 (6%)	0	100	100
1	C	196/205 (96%)	184 (94%)	12 (6%)	0	100	100
1	E	196/205 (96%)	184 (94%)	12 (6%)	0	100	100
1	G	196/205 (96%)	185 (94%)	11 (6%)	0	100	100
1	I	190/205 (93%)	135 (71%)	37 (20%)	18 (10%)	0	1
1	K	190/205 (93%)	134 (70%)	37 (20%)	19 (10%)	0	1
1	M	190/205 (93%)	135 (71%)	37 (20%)	18 (10%)	0	1
1	O	190/205 (93%)	134 (70%)	37 (20%)	19 (10%)	0	1
2	B	277/279 (99%)	253 (91%)	23 (8%)	1 (0%)	34	66
2	D	277/279 (99%)	254 (92%)	22 (8%)	1 (0%)	34	66
2	F	277/279 (99%)	254 (92%)	22 (8%)	1 (0%)	34	66
2	H	277/279 (99%)	253 (91%)	23 (8%)	1 (0%)	34	66
2	J	275/279 (99%)	238 (86%)	25 (9%)	12 (4%)	2	8
2	L	277/279 (99%)	242 (87%)	24 (9%)	11 (4%)	3	9
2	N	275/279 (99%)	238 (86%)	25 (9%)	12 (4%)	2	8
2	P	277/279 (99%)	242 (87%)	24 (9%)	11 (4%)	3	9

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
All	All	3756/3872 (97%)	3250 (86%)	382 (10%)	124 (3%)	4	13

5 of 124 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	176	PRO
2	D	176	PRO
2	F	176	PRO
2	H	176	PRO
1	I	68	LEU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/175 (97%)	134 (79%)	35 (21%)	1	3
1	C	169/175 (97%)	132 (78%)	37 (22%)	1	3
1	E	169/175 (97%)	132 (78%)	37 (22%)	1	3
1	G	169/175 (97%)	134 (79%)	35 (21%)	1	3
1	I	169/175 (97%)	121 (72%)	48 (28%)	0	1
1	K	169/175 (97%)	121 (72%)	48 (28%)	0	1
1	M	169/175 (97%)	121 (72%)	48 (28%)	0	1
1	O	169/175 (97%)	121 (72%)	48 (28%)	0	1
2	B	226/226 (100%)	181 (80%)	45 (20%)	1	4
2	D	226/226 (100%)	181 (80%)	45 (20%)	1	4
2	F	226/226 (100%)	181 (80%)	45 (20%)	1	4
2	H	226/226 (100%)	181 (80%)	45 (20%)	1	4
2	J	226/226 (100%)	188 (83%)	38 (17%)	2	6
2	L	226/226 (100%)	189 (84%)	37 (16%)	2	7
2	N	226/226 (100%)	188 (83%)	38 (17%)	2	6
2	P	226/226 (100%)	189 (84%)	37 (16%)	2	7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	3160/3208 (98%)	2494 (79%)	666 (21%)	1 3

5 of 666 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	H	138	ASN
2	J	17	SER
1	O	132	ARG
2	H	183	LEU
1	I	61	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 113 such sidechains are listed below:

Mol	Chain	Res	Type
1	I	144	ASN
1	K	24	ASN
2	P	96	ASN
2	J	41	GLN
2	J	206	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	K	5
1	I	5
1	O	5
1	M	5
2	P	2
2	J	2
2	D	2
2	H	2
2	B	2
2	N	2
2	L	2
2	F	2

The worst 5 of 36 chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	J	173:PRO	C	174:ASP	N	2.75
1	N	173:PRO	C	174:ASP	N	2.75
1	I	74:GLN	C	75:LEU	N	2.13
1	M	74:GLN	C	75:LEU	N	2.13
1	K	74:GLN	C	75:LEU	N	2.05

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.