

Full wwPDB NMR Structure Validation Report (i)

Nov 7, 2023 – 06:59 pm GMT

PDB ID	:	6QYS
BMRB ID	:	34370
Title	:	Solution NMR of synthetic analogues of nisin and mutacin ring A and ring B
		- Nisin Ring B
Authors	:	Dickman, R.; Mitchell, S.A.; Figueiredo, A.; Hansen, D.F.; Tabor, A.B.
Deposited on	:	2019-03-09

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


~		
Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as 541 be (2020)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
wwPDB-RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
wwPDB-ShiftChecker	:	v1.2
BMRB Restraints Analysis	:	v1.2
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 43%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f NMR} { m archive} \ (\#{ m Entries})$		
Clashscore	158937	12864		
Ramachandran outliers	154571	11451		
Sidechain outliers	154315	11428		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain			
1	А	5	20%	40%	40%	



2 Ensemble composition and analysis (i)

This entry contains 15 models.

Cyrange was unable to find well-defined residues.

Error message: The number of core atoms (2) was below the domain threshold value (8).

NmrClust was unable to cluster the ensemble.

Error message: Wrapper check: not enough residues in core to run NmrClust



3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 68 atoms, of which 35 are hydrogens and 0 are deuteriums.

• Molecule 1 is a protein called DBB-PRO-GLY-CYS-LYS.

Mol	Chain	Residues		Atoms					Trace					
1	А	A 5	Δ	٨	٨	٨	1	Total	С	Η	Ν	0	S	0
			68	20	35	6	6	1	0					




4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


• Molecule 1: DBB-PRO-GLY-CYS-LYS



4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1



4.2.3 Score per residue for model 3

• Molecule 1: DBB-PRO-GLY-CYS-LYS



Chain A:	20%	60%		20%
11 83 63 75 85				
4.2.4 Scor	e per residu	ue for model 4		
	DBB-PRO-	GLY-CYS-LYS		
Chain A:	40%		60%	
11 P2 G3 K5 K5				
4.2.5 Scor	e per residu	ue for model 5		
• Molecule 1:	DBB-PRO-	GLY-CYS-LYS		
Chain A:	20%	60%		20%
11 22 83 85 85				
4.2.6 Scor	e per residu	ue for model 6		
• Molecule 1:	DBB-PRO-	GLY-CYS-LYS		
Chain A:	20%	40%	40%	
T1 G3 K5 K5				
4.2.7 Scor	e per residu	ue for model 7		
• Molecule 1:	DBB-PRO-	GLY-CYS-LYS		
Chain A:	20%	40%	40%	
T1 P2 G3 K5 K5				
4.2.8 Scor	e per residu	ue for model 8		
• Molecule 1:	DBB-PRO-	GLY-CYS-LYS		
Chain A:	20%	40%	40%	
		WORLDW PROTEIN DAT		

T1 P2 G3 C4 K5

4.2.9	Score per	residue fo	or model 9			
• Mole	ecule 1: DBF	3-PRO-GLY	CYS-LYS			
Chain	A: 20%		40%		40%	
T1 B2 C4 K5						
4.2.10	Score pe	er residue :	for model	10		
• Mole	ecule 1: DBE	B-PRO-GLY	CYS-LYS			
Chain	A: 20%		6	0%		20%
11 83 85 85						
4.2.11	Score pe	er residue :	for model	11		
• Mole	ecule 1: DBE	B-PRO-GLY	CYS-LYS			
Chain	A:	40%	2	0%	40%	
T1 P2 G3 K5						
4.2.12	Score pe	er residue :	for model	12		
• Mole	ecule 1: DBE	B-PRO-GLY	CYS-LYS			
Chain	A: 20%		6	0%		20%
11 63 63 85						
4.2.13	Score pe	er residue :	for model	13		
• Mole	ecule 1: DBE	8-PRO-GLY	-CYS-LYS			
Chain	A:	40%	2	0%	40%	
T1 P2 C4 K5	l					

W O R L D W I D E PROTEIN DATA BANK

4.2.14 Score per residue for model 14

• Molecule 1: DBB-PRO-GLY-CYS-LYS

Chain A	: 20%	40%	40%
11 83 63 85 85			
4.2.15	Score per re	esidue for model 15	

• Molecule 1: DBB-PRO-GLY-CYS-LYS

Chain A:	20%	20%	60%
T 1 2 2 6 3 7 5 7 5 7 5			



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 150 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Xplor-NIH	structure calculation	2.45
Xplor-NIH	refinement	2.45

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	24
Number of shifts mapped to atoms	24
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	43%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: DBB

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	А	33	35	34	7 ± 2
All	All	495	525	510	108

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 107.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Mod	dels
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:1:DBB:HG3	1:A:4:CYS:H	0.99	1.17	1	15
1:A:1:DBB:HG3	1:A:4:CYS:N	0.91	1.81	1	15
1:A:1:DBB:CG	1:A:4:CYS:H	0.71	1.98	7	15
1:A:5:LYS:CD	1:A:5:LYS:H	0.62	2.07	9	2
1:A:4:CYS:O	1:A:5:LYS:C	0.62	2.38	4	4
1:A:3:GLY:O	1:A:4:CYS:O	0.59	2.20	2	12
1:A:4:CYS:O	1:A:5:LYS:O	0.59	2.21	7	2
1:A:3:GLY:O	1:A:4:CYS:C	0.57	2.42	4	15
1:A:4:CYS:O	1:A:5:LYS:OXT	0.55	2.24	6	3
1:A:5:LYS:CD	1:A:5:LYS:N	0.53	2.70	9	2
1:A:3:GLY:O	1:A:5:LYS:N	0.45	2.49	7	1
1:A:1:DBB:HA	1:A:4:CYS:H	0.45	1.72	10	15

All unique clashes are listed below, sorted by their clash magnitude.

Continued on next page...



Atom 1	Atom 2	Clash(Å)	Distance(Å)	Moo	dels
Atom-1	Atom-2	Clash(A)	Distance(A)	Worst	Total
1:A:5:LYS:N	1:A:5:LYS:HD3	0.44	2.28	9	2
1:A:1:DBB:HA	1:A:2:PRO:C	0.43	2.32	1	3
1:A:4:CYS:C	1:A:5:LYS:OXT	0.41	2.57	6	1
1:A:1:DBB:HA	1:A:4:CYS:N	0.40	2.32	4	1

Continued from previous page...

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	3/5~(60%)	$2\pm0~(67\pm0\%)$	0±0 (0±0%)	$1\pm0~(33\pm0\%)$	0 0
All	All	45/75~(60%)	30~(67%)	0 (0%)	15 (33%)	0 0

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	А	4	CYS	15

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the side chain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	А	3/3~(100%)	2 ± 0 (78 $\pm16\%$)	$1\pm0~(22\pm16\%)$	3	30
All	All	45/45~(100%)	35~(78%)	10 (22%)	3	30

All 1 unique residues with a non-rotameric sidechain are listed below.

Mol	Chain	Res	Type	Models (Total)
1	А	5	LYS	10



6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mal	Tuno	Chain	Dog	Link		Bond leng	gths
	туре	Ullaili	nes	Link	Counts	RMSZ	#Z>2
1	DBB	А	1	1	$4,\!5,\!6$	$0.53 {\pm} 0.01$	0±0 (0±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mal	Turne	Chain	Dec	Tink		Bond an	ngles
10101	туре	Chain	nes	LIIIK	Counts	RMSZ	#Z>2
1	DBB	А	1	1	$1,\!5,\!7$	2.03 ± 0.04	1 ± 0 (66±47%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	DBB	А	1	1	-	$0\pm 0,3,4,6$	-

There are no bond-length outliers.

All unique angle outliers are listed below.



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$\mathrm{Ideal}(^{o})$	Moo Worst	
1	А	1	DBB	CG-CB-CA	2.12	108.57	113.42	15	10

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 43% for the well-defined parts and 43% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: NMRStarshifts.txt

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	24
Number of shifts mapped to atoms	24
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 43%, i.e. 19 atoms were assigned a chemical shift out of a possible 44. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	15 N
Backbone	16/19~(84%)	8/8~(100%)	8/8~(100%)	0/3~(0%)
Sidechain	3/25~(12%)	0/16~(0%)	3/8~(38%)	0/1~(0%)
Overall	19/44~(43%)	8/24 (33%)	11/16~(69%)	0/4~(0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 43%, i.e. 19 atoms were assigned a chemical shift out of a possible 44. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.



	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	15 N
Backbone	16/19~(84%)	8/8 (100%)	8/8~(100%)	0/3~(0%)
Sidechain	3/25~(12%)	0/16~(0%)	3/8~(38%)	0/1~(0%)
Overall	19/44~(43%)	8/24 (33%)	11/16~(69%)	0/4~(0%)


7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:





8 NMR restraints analysis (i)

8.1 Conformationally restricting restraints (i)

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	22
Intra-residue $(i-j =0)$	0
Sequential (i-j =1)	15
Medium range ($ i-j >1$ and $ i-j <5$)	7
Long range $(i-j \ge 5)$	0
Inter-chain	0
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	3
Number of unmapped restraints	0
Number of restraints per residue	5.0
Number of long range restraints per residue ¹	0.0

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations (i)

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model (i)

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	1.2	0.2
0.2-0.5 (Medium)	2.7	0.46
>0.5 (Large)	None	None



8.2.2 Average number of dihedral-angle violations per model (i)

Dihedral-angle violations less than 1° are not included in the calculation.

Bins $(^{\circ})$	Average number of violations per model	Max ($^{\circ}$)
1.0-10.0 (Small)	1.0	4.55
10.0-20.0 (Medium)	None	None
>20.0 (Large)	None	None



9 Distance violation analysis (i)


9.1 Summary of distance violations (i)


The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Destroints type	Count	$\%^1$	${f Violated^3}$			Consis	tently	$Violated^4$
Restraints type	Count	70-	Count	$\%^2$	$\%^{1}$	Count	$\%^2$	$\%^1$
Intra-residue (i-j =0)	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sequential (i-j =1)	15	68.2	2	13.3	9.1	1	6.7	4.5
Backbone-Backbone	7	31.8	2	28.6	9.1	1	14.3	4.5
Backbone-Sidechain	7	31.8	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	1	4.5	0	0.0	0.0	0	0.0	0.0
Medium range ($ i-j > 1 \& i-j < 5$)	7	31.8	2	28.6	9.1	2	28.6	9.1
Backbone-Backbone	3	13.6	1	33.3	4.5	1	33.3	4.5
Backbone-Sidechain	3	13.6	1	33.3	4.5	1	33.3	4.5
Sidechain-Sidechain	1	4.5	0	0.0	0.0	0	0.0	0.0
Long range $(i-j \ge 5)$	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Inter-chain	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	0	0.0	0	0.0	0.0	0	0.0	0.0
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	22	100.0	4	18.2	18.2	3	13.6	13.6
Backbone-Backbone	10	45.5	3	30.0	13.6	2	20.0	9.1
Backbone-Sidechain	10	45.5	1	10.0	4.5	1	10.0	4.5
Sidechain-Sidechain	2	9.1	0	0.0	0.0	0	0.0	0.0

 1 percentage calculated with respect to the total number of distance restraints, 2 percentage calculated with respect to the number of restraints in a particular restraint category, 3 violated in at least one model, 4 violated in all the models







Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model (i)

The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Madal ID		Nun	nber o	f viola	ations	5	Maan (Å)	Mor (Å)	SD^6 (Å)	Madian (Å)
Model ID	IR^1	SQ^2	MR^3	LR^4	$ IC^5 $	Total	Mean (Å)	Max (Å)	$SD^{*}(A)$	Median (Å)
1	0	2	2	0	0	4	0.26	0.32	0.09	0.3
2	0	2	2	0	0	4	0.25	0.33	0.08	0.28
3	0	2	2	0	0	4	0.26	0.39	0.09	0.24
4	0	2	2	0	0	4	0.26	0.34	0.08	0.28
5	0	2	2	0	0	4	0.25	0.4	0.1	0.24
6	0	2	2	0	0	4	0.24	0.38	0.09	0.24
7	0	1	2	0	0	3	0.3	0.46	0.12	0.27
8	0	2	2	0	0	4	0.25	0.37	0.09	0.26
9	0	2	2	0	0	4	0.26	0.39	0.09	0.24
10	0	2	2	0	0	4	0.26	0.45	0.12	0.22
11	0	2	2	0	0	4	0.25	0.37	0.09	0.26

Continued on next page...




Model ID	Number of violations						Mean (Å)	Max (Å)	SD^6 (Å)	Median (Å)
model ID	IR^1	SQ^2	MR^3	LR^4	IC^5	Total		Max (II)		
12	0	2	2	0	0	4	0.26	0.39	0.09	0.24
13	0	2	2	0	0	4	0.25	0.41	0.1	0.23
14	0	2	2	0	0	4	0.25	0.35	0.08	0.26
15	0	2	2	0	0	4	0.24	0.35	0.09	0.24

Continued from previous page...

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints, ⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model (i)



The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble (i)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints




Nu	mber	of vio	lated	Fraction of the ensemble			
IR^1	SQ^2	MR^3	LR^4	IC ⁵	Total	Count ⁶	%
0	0	0	0	0	0	1	6.7
0	0	0	0	0	0	2	13.3
0	0	0	0	0	0	3	20.0
0	0	0	0	0	0	4	26.7
0	0	0	0	0	0	5	33.3
0	0	0	0	0	0	6	40.0
0	0	0	0	0	0	7	46.7
0	0	0	0	0	0	8	53.3
0	0	0	0	0	0	9	60.0
0	0	0	0	0	0	10	66.7
0	0	0	0	0	0	11	73.3
0	0	0	0	0	0	12	80.0
0	0	0	0	0	0	13	86.7
0	1	0	0	0	1	14	93.3
0	1	2	0	0	3	15	100.0

for a given fraction of the ensemble. In total, 18(IR:0, SQ:13, MR:5, LR:0, IC:0) restraints are not violated in the ensemble.

 1 Intra-residue restraints, 2 Sequential restraints, 3 Medium range restraints, 4 Long range restraints, 5 Inter-chain restraints, 6 Number of models with violations






9.3.1 Bar graph : Distance violation statistics for the ensemble (i)

9.4 Most violated distance restraints in the ensemble (i)

9.4.1 Histogram : Distribution of mean distance violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





9.4.2 Table: Most violated distance restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	\mathbf{Models}^1	Mean (Å)	SD^1 (Å)	Median (Å)
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	15	0.38	0.04	0.38
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	15	0.27	0.01	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	15	0.27	0.01	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	15	0.27	0.01	0.27
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	15	0.13	0.02	0.13
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	14	0.23	0.04	0.22


¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints (i)

9.5.1 Histogram : Distribution of distance violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





9.5.2 Table : All distance violations (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	7	0.46
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	10	0.45
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	13	0.41
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	5	0.4
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	3	0.39
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	9	0.39
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	12	0.39
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	6	0.38
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	8	0.37
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	11	0.37
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	14	0.35
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	15	0.35
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	4	0.34
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	2	0.33
(1,6)	1:5:A:LYS:H	1:4:A:CYS:HA	1	0.32
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	1	0.3

Continued on next page...



Contin	ued from previous p	page		
Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	1	0.3
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	1	0.3
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	1	0.3
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	2	0.28
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	4	0.28
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	2	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	2	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	2	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	3	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	3	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	3	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	4	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	4	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	4	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	5	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	5	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	5	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	7	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	7	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	7	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	8	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	8	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	8	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	9	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	9	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	9	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	10	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	10	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	10	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	11	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	11	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	11	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	12	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	12	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	12	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	13	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	13	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	13	0.27
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	14	0.27
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	14	0.27
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	14	0.27
(1 1 0)			-	0.00

Co

Continued on next page...

0.26



6

1:4:A:CYS:H

1:1:A:DBB:HG1

(1,18)

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	6	0.26
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	6	0.26
(1,18)	1:1:A:DBB:HG1	1:4:A:CYS:H	15	0.26
(1,18)	1:1:A:DBB:HG2	1:4:A:CYS:H	15	0.26
(1,18)	1:1:A:DBB:HG3	1:4:A:CYS:H	15	0.26
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	14	0.26
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	8	0.24
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	11	0.24
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	15	0.23
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	3	0.22
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	6	0.22
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	9	0.22
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	12	0.22
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	5	0.2
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	13	0.19
(1,5)	1:2:A:PRO:HA	1:3:A:GLY:H	10	0.17
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	7	0.16
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	10	0.16
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	3	0.14
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	9	0.14
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	12	0.14
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	4	0.13
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	5	0.13
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	8	0.13
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	11	0.13
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	13	0.13
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	6	0.12
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	14	0.12
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	1	0.11
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	2	0.11
(1,19)	1:4:A:CYS:H	1:1:A:DBB:HA	15	0.11

Continued from previous page...




10 Dihedral-angle violation analysis (i)


10.1 Summary of dihedral-angle violations (i)

The following table provides the summary of dihedral-angle violations in different dihedral-angle types. Violations less than 1° are not included in the calculation.

A 1	Count	$\%^1$	${f Violated^3}$			Consistently Violated ⁴		
Angle type			Count	$\%^2$	$\%^1$	Count	$\%^2$	$\%^1$
PHI	3	100.0	1	33.3	33.3	1	33.3	33.3
Total	3	100.0	1	33.3	33.3	1	33.3	33.3

 1 percentage calculated with respect to total number of dihedral-angle restraints, 2 percentage calculated with respect to number of restraints in a particular dihedral-angle type, 3 violated in at least one model, 4 violated in all the models






Violated and consistently violated restraints are shown using different hatch patterns in their respective categories




10.2 Dihedral-angle violation statistics for each model (i)

The following table provides the dihedral-angle violation statistics for each model in the ensemble. Violations less than 1° are not included in the statistics.

Model ID	Number of violations		Mean (°)	Max (°)	SD (°)	Median (°)	
	PHI	Total	()	()	~ ()		
1	1	1	2.93	2.93	0.0	2.93	
2	1	1	3.25	3.25	0.0	3.25	
3	1	1	4.55	4.55	0.0	4.55	
4	1	1	4.03	4.03	0.0	4.03	
5	1	1	4.01	4.01	0.0	4.01	
6	1	1	3.73	3.73	0.0	3.73	
7	1	1	3.51	3.51	0.0	3.51	
8	1	1	4.06	4.06	0.0	4.06	
9	1	1	4.07	4.07	0.0	4.07	
10	1	1	4.54	4.54	0.0	4.54	
11	1	1	4.13	4.13	0.0	4.13	
12	1	1	4.53	4.53	0.0	4.53	
13	1	1	3.74	3.74	0.0	3.74	
14	1	1	3.74	3.74	0.0	3.74	
15	1	1	3.44	3.44	0.0	3.44	







The mean(dot), median(x) and the standard deviation are shown in blue with respect to the y axis on the right

10.3 Dihedral-angle violation statistics for the ensemble (i)


Violation analysis may find that some restraints are violated in very few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of ensemble.

Num	ber of violated restraints	Fraction of the ensemble		
PHI	Total	Count^1	%	
0	0	1	6.7	
0	0	2	13.3	
0	0	3	20.0	
0	0	4	26.7	
0	0	5	33.3	
0	0	6	40.0	
0	0	7	46.7	
0	0	8	53.3	
0	0	9	60.0	
0	0	10	66.7	
0	0	11	73.3	
0	0	12	80.0	
0	0	13	86.7	
0	0	14	93.3	
1	1	15	100.0	

¹ Number of models with violations



10.3.1 Bar graph : Dihedral-angle Violation statistics for the ensemble (i)




10.4 Most violated dihedral-angle restraints in the ensemble (i)

10.4.1 Histogram : Distribution of mean dihedral-angle violations (i)

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble





10.4.2 Table: Most violated dihedral-angle restraints (i)

The following table provides the mean and the standard deviation of the violation for each restraint sorted by number of violated models and the mean value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	\mathbf{Models}^1	Mean	\mathbf{SD}^2	Median
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	15	3.88	0.46	4.01


¹ Number of violated models, ²Standard deviation, All angle values are in degree (°)

10.5 All violated dihedral-angle restraints (i)

10.5.1 Histogram : Distribution of violations (i)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.





10.5.2 Table: All violated dihedral-angle restraints (i)

The following table lists the absolute value of the violation for each restraint in the ensemble sorted by its value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint.

Key	Atom-1	Atom-2	Atom-3	Atom-4	Model ID	Violation ($^{\circ}$)
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	3	4.55
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	10	4.54
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	12	4.53
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	11	4.13
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	9	4.07
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	8	4.06
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	4	4.03
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	5	4.01
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	13	3.74
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	14	3.74
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	6	3.73
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	7	3.51
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	15	3.44
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	2	3.25
(1,3)	1:3:A:GLY:C	1:4:A:CYS:N	1:4:A:CYS:CA	1:4:A:CYS:C	1	2.93

