



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 9, 2020 – 08:45 AM BST

PDB ID : 5QZO  
Title : PanDDA analysis group deposition – Auto-refined data of Aar2/RNaseH for ground state model 39  
Authors : Weiss, M.S.; Wollenhaupt, J.; Metz, A.; Barthel, T.; Lima, G.M.A.; Heine, A.; Mueller, U.; Klebe, G.  
Deposited on : 2020-02-12  
Resolution : 1.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1



## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4728 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Pre-mRNA-splicing factor 8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	237	2002	1283	335	372	12	0	12	0

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1833	GLY	-	expression tag	UNP P33334
A	1834	ALA	-	expression tag	UNP P33334
A	1835	MET	-	expression tag	UNP P33334

- Molecule 2 is a protein called A1 cistron-splicing factor AAR2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	300	2580	1654	421	485	20	0	9	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-3	GLY	-	expression tag	UNP P32357
B	-2	ALA	-	expression tag	UNP P32357
B	-1	MET	-	expression tag	UNP P32357
B	0	ALA	-	expression tag	UNP P32357
B	166	SER	-	linker	UNP P32357
B	167	SER	-	linker	UNP P32357
B	168	SER	-	linker	UNP P32357
B	169	SER	-	linker	UNP P32357
B	170	SER	-	linker	UNP P32357


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	76	Total 76	O 76	0	0
3	B	70	Total 70	O 70	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Pre-mRNA-splicing factor 8



- Molecule 2: A1 cistron-splicing factor AAR2



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	88.33Å 81.94Å 93.58Å 90.00° 108.41° 90.00°	Depositor
Resolution (Å)	44.44 – 1.39 44.40 – 1.39	Depositor EDS
% Data completeness (in resolution range)	95.6 (44.44-1.39) 95.7 (44.40-1.39)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.96 (at 1.39Å)	Xtrriage
Refinement program	REFMAC 5.8.0238, PHENIX 1.16.3549	Depositor
R, $R_{free}$	0.220 , 0.221 0.225 , 0.222	Depositor DCC
$R_{free}$ test set	5938 reflections (4.88%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	29.9	Xtrriage
Anisotropy	0.244	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 40.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4728	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.76% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.74	0/2049	0.88	2/2775 (0.1%)
2	B	0.72	0/2651	0.84	1/3581 (0.0%)
All	All	0.73	0/4700	0.86	3/6356 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	1859	ARG	NE-CZ-NH1	-7.52	116.54	120.30
1	A	1859	ARG	NE-CZ-NH2	7.46	124.03	120.30
2	B	123	MET	CG-SD-CE	5.63	109.21	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2002	0	2029	10	0
2	B	2580	0	2450	15	0
3	A	76	0	0	1	0
3	B	70	0	0	4	0
All	All	4728	0	4479	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:280:SER:HB3	2:B:313:TYR:CE1	2.17	0.79
1:A:1976:ASP:HB3	3:A:2137:HOH:O	1.94	0.66
2:B:1:MET:HB3	2:B:35:ASP:HA	1.78	0.65
2:B:70:GLN:HB3	2:B:81:MET:HE2	1.82	0.60
1:A:1962:ARG:O	1:A:2013:ARG:NH1	2.35	0.60
1:A:1848:ILE:H	1:A:1931[A]:LYS:HZ2	1.52	0.58
2:B:70:GLN:HB3	2:B:81:MET:CE	2.35	0.56
2:B:280:SER:CB	2:B:313:TYR:CE1	2.87	0.56
2:B:287:ARG:O	2:B:291:ILE:HD13	2.07	0.55
1:A:2062:GLU:HB3	1:A:2066:LYS:HE3	1.89	0.55
2:B:54[A]:MET:HE1	3:B:461:HOH:O	2.06	0.55
2:B:258:LYS:HD2	2:B:258:LYS:H	1.71	0.55
2:B:2:ASN:HB2	3:B:433:HOH:O	2.08	0.54
1:A:1836:ASN:OD1	1:A:1838:SER:HB2	2.09	0.52
1:A:2062:GLU:O	1:A:2066:LYS:HG2	2.10	0.52
2:B:54[A]:MET:CE	3:B:461:HOH:O	2.59	0.51
2:B:280:SER:HB3	2:B:313:TYR:CZ	2.49	0.48
1:A:2023:LYS:O	1:A:2027:LEU:HG	2.16	0.46
2:B:16:GLY:HA3	2:B:45:HIS:CE1	2.51	0.45
1:A:2058:LEU:HD23	1:A:2058:LEU:C	2.37	0.45
2:B:51:ASN:C	2:B:53:SER:H	2.20	0.45
2:B:51:ASN:ND2	3:B:404:HOH:O	2.53	0.42
1:A:1843:LEU:HA	1:A:1849:LYS:HD2	2.02	0.42
1:A:1848:ILE:H	1:A:1931[A]:LYS:NZ	2.19	0.40
2:B:77:LEU:HD21	2:B:79:LYS:HE3	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	248/258 (96%)	242 (98%)	6 (2%)	0	100	100
2	B	306/308 (99%)	295 (96%)	11 (4%)	0	100	100
All	All	554/566 (98%)	537 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	226/233 (97%)	222 (98%)	4 (2%)	59	28
2	B	287/284 (101%)	285 (99%)	2 (1%)	84	66
All	All	513/517 (99%)	507 (99%)	6 (1%)	81	47

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1962	ARG
1	A	1979[A]	MET
1	A	1979[B]	MET
1	A	1979[C]	MET
2	B	258	LYS
2	B	315	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	150	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

#### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

#### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

#### 5.7 Other polymers [i](#)

There are no such residues in this entry.

#### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	237/258 (91%)	0.84	24 (10%) 7 6	21, 30, 59, 88	0
2	B	300/308 (97%)	0.86	33 (11%) 5 5	22, 33, 64, 81	0
All	All	537/566 (94%)	0.85	57 (10%) 6 5	21, 32, 63, 88	0

All (57) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	1	MET	8.3
2	B	53	SER	7.2
2	B	172	PRO	6.5
2	B	279	TYR	6.4
1	A	1833	GLY	6.3
1	A	2063	TYR	6.0
1	A	2027	LEU	5.7
1	A	2069	VAL	5.4
2	B	170	SER	5.3
1	A	2066	LYS	5.2
2	B	173	ALA	5.1
2	B	52	SER	5.1
1	A	2064	GLY	5.0
1	A	2028	SER	4.5
1	A	2068	ASN	4.3
1	A	2060	LEU	4.2
2	B	54[A]	MET	4.1
1	A	1866	PHE	4.0
1	A	1836	ASN	3.9
2	B	22	PHE	3.8
2	B	316	LEU	3.8
2	B	109	ASP	3.7
2	B	50	ASP	3.6
1	A	1962	ARG	3.5

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	171	ASP	3.5
1	A	1838	SER	3.5
2	B	108	ILE	3.3
1	A	1837	SER	3.2
2	B	87	ALA	3.2
2	B	84	ARG	3.1
2	B	280	SER	2.9
2	B	278	GLN	2.9
2	B	110	GLU	2.9
1	A	2065	ARG	2.8
1	A	1835	MET	2.8
1	A	1979[A]	MET	2.8
2	B	313	TYR	2.7
1	A	1953	ASN	2.7
2	B	101	MET	2.7
2	B	174	HIS	2.7
1	A	2061	THR	2.7
2	B	91	ASN	2.6
1	A	2057[A]	ASP	2.6
2	B	96	PHE	2.5
1	A	1991	ILE	2.5
1	A	2046	GLU	2.4
2	B	117	LEU	2.4
1	A	2067	TYR	2.4
2	B	224	LEU	2.3
2	B	62	CYS	2.3
2	B	292	CYS	2.3
1	A	1904	ARG	2.2
2	B	203[A]	TYR	2.1
2	B	150	ASN	2.1
2	B	86	GLY	2.1
2	B	293	LEU	2.1
2	B	317	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.