



wwPDB X-ray Structure Validation Summary Report

Oct 16, 2021 – 10:15 PM EDT

PDB ID : 1R0A
Title : Crystal structure of HIV-1 reverse transcriptase covalently tethered to DNA template-primer solved to 2.8 angstroms
Authors : Tuske, S.; Ding, J.; Arnold, E.
Deposited on : 2003-09-19
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the  symbol.

The following versions of software and data (see [references](#) ) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	T	27	
2	P	21	
3	A	558	
4	B	429	
5	L	211	
6	H	225	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
7	GOL	B	3001	-	X	-	-
7	GOL	P	3002	-	X	-	-

2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 12292 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called 5'-D(*A*TP*GP*CP*AP*TP*CP*GP*GP*CP*GP*CP*TP*CP*GP*AP*AP*CP*AP*GP*GP*GP*AP*CP*GP*GP*T)-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
1	T	24	493	233	97	140	23	0	0	0

- Molecule 2 is a DNA chain called 5'-D(*C*CP*GP*TP*CP*CP*CP*TP*GP*TP*TP*CP*GP*AP*GP*CP*GP*CP*CP*GP*(2DA))-3'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	P	20	402	192	72	119	19	0	0	0

- Molecule 3 is a protein called Reverse transcriptase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	A	558	4482	2901	741	832	8	15	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	258	CYS	GLN	engineered mutation	UNP P03366
A	280	SER	CYS	engineered mutation	UNP P03366

- Molecule 4 is a protein called Reverse transcriptase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	B	429	3534	2304	586	637	7	4	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	280	SER	CYS	engineered mutation	UNP P03366


- Molecule 5 is a protein called monoclonal antibody (light chain).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
5	L	211	1643	1025	270	342	6	0	0	0

- Molecule 6 is a protein called monoclonal antibody (heavy chain).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
6	H	225	1685	1060	276	340	9	0	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
7	P	1	6	3	3	0	0
7	B	1	6	3	3	0	0

- Molecule 8 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Mg	0	0
			1	1		

- Molecule 9 is alpha-D-glucopyranose (three-letter code: GLC) (formula: C₆H₁₂O₆).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	H	1	Total C O 12 6 6	0	0



- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	T	1	Total O 1 1	0	0
10	A	7	Total O 7 7	0	0
10	B	15	Total O 15 15	0	0
10	L	2	Total O 2 2	0	0
10	H	3	Total O 3 3	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 5'-D(*A*TP*GP*CP*AP*TP*CP*GP*GP*CP*GP*CP*TP*CP*GP*AP*AP*CP*AP*GP*GP*GP*AP*CP*GP*GP*T)-3'




- Molecule 2: 5'-D(*C*CP*GP*TP*CP*CP*CP*TP*GP*TP*TP*CP*GP*AP*GP*CP*GP*CP*CP*GP*(2DA))-3'




- Molecule 3: Reverse transcriptase








• Molecule 4: Reverse transcriptase



• Molecule 5: monoclonal antibody (light chain)



• Molecule 6: monoclonal antibody (heavy chain)



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 1 2	Depositor
Cell constants a, b, c, α , β , γ	165.82Å 165.82Å 220.72Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.96 – 2.80 39.20 – 2.80	Depositor EDS
% Data completeness (in resolution range)	90.0 (19.96-2.80) 94.6 (39.20-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.19 (at 2.81Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.239 , 0.272 0.237 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	60.6	Xtrriage
Anisotropy	0.010	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 57.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	12292	wwPDB-VP
Average B, all atoms (Å ²)	66.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.11% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, 2DA, MG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	T	0.77	1/554 (0.2%)	1.12	3/854 (0.4%)
2	P	0.69	0/426	1.02	3/655 (0.5%)
3	A	0.51	1/4600 (0.0%)	0.76	1/6259 (0.0%)
4	B	0.59	0/3639	0.79	1/4949 (0.0%)
5	L	0.52	0/1681	0.78	1/2283 (0.0%)
6	H	0.53	0/1729	0.80	1/2372 (0.0%)
All	All	0.56	2/12629 (0.0%)	0.81	10/17372 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	B	0	2
5	L	0	1
All	All	0	3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	248	GLU	CD-OE2	6.89	1.33	1.25
1	T	703	DG	O3'-P	5.73	1.68	1.61

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	T	704	DC	O5'-P-OP1	-14.16	92.96	105.70
2	P	816	DG	N9-C1'-C2'	7.49	126.83	112.60
2	P	815	DA	N9-C1'-C2'	7.13	126.15	112.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	H	187	LEU	CA-CB-CG	6.72	130.75	115.30
3	A	289	LEU	N-CA-C	5.74	126.50	111.00

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	B	188	TYR	Sidechain
4	B	405	TYR	Sidechain
5	L	49	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	T	493	0	269	38	0
2	P	402	0	226	23	0
3	A	4482	0	4485	400	0
4	B	3534	0	3568	270	2
5	L	1643	0	1565	136	0
6	H	1685	0	1640	92	0
7	B	6	0	4	0	0
7	P	6	0	4	0	0
8	A	1	0	0	0	0
9	H	12	0	12	2	0
10	A	7	0	0	1	0
10	B	15	0	0	0	0
10	H	3	0	0	0	0
10	L	2	0	0	0	0
10	T	1	0	0	0	0
All	All	12292	0	11773	918	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 38.

The worst 5 of 918 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:T:703:DG:H1'	1:T:704:DC:OP1	1.50	1.11
3:A:498:ASP:HB2	3:A:538:ALA:HB2	1.17	1.10
3:A:22:LYS:H	3:A:22:LYS:HD3	1.12	1.09
1:T:721:DG:H2''	1:T:722:DG:H5''	1.10	1.08
4:B:296:THR:HG22	4:B:298:GLU:H	1.09	1.07

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:B:2:ILE:O	4:B:2:ILE:O[6_565]	1.96	0.24
4:B:3:SER:OG	4:B:3:SER:OG[6_565]	2.02	0.18

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	556/558 (100%)	456 (82%)	69 (12%)	31 (6%)	2	5
4	B	427/429 (100%)	361 (84%)	52 (12%)	14 (3%)	4	13
5	L	209/211 (99%)	184 (88%)	18 (9%)	7 (3%)	4	13
6	H	223/225 (99%)	198 (89%)	22 (10%)	3 (1%)	12	36
All	All	1415/1423 (99%)	1199 (85%)	161 (11%)	55 (4%)	3	10

5 of 55 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	A	273	GLY
3	A	278	GLN
3	A	286	THR
3	A	345	PRO
3	A	466	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	485/498 (97%)	437 (90%)	48 (10%)	8	23
4	B	388/391 (99%)	351 (90%)	37 (10%)	8	25
5	L	190/190 (100%)	169 (89%)	21 (11%)	6	19
6	H	196/196 (100%)	180 (92%)	16 (8%)	11	33
All	All	1259/1275 (99%)	1137 (90%)	122 (10%)	8	24

5 of 122 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
4	B	109	LEU
6	H	72	SER
4	B	295	LEU
6	H	64	SER
6	H	192	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
5	L	90	GLN
6	H	62	ASN
5	L	137	ASN
5	L	210	ASN
6	H	112	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	2DA	P	822	1,2	17,22,23	0.74	0	13,31,34	0.96	1 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	2DA	P	822	1,2	-	0/3/18/19	0/3/3/3

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	822	2DA	C5-C6-N6	2.30	123.85	120.35

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	P	822	2DA	1	0

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 1 is monoatomic - leaving 3 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
7	GOL	P	3002	-	5,5,5	4.50	5 (100%)	5,5,5	5.78	3 (60%)
9	GLC	H	1725	-	12,12,12	0.61	0	17,17,17	0.67	0
7	GOL	B	3001	-	5,5,5	4.42	5 (100%)	5,5,5	5.77	3 (60%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
7	GOL	P	3002	-	-	3/4/4/4	-
9	GLC	H	1725	-	-	0/2/22/22	0/1/1/1
7	GOL	B	3001	-	-	2/4/4/4	-

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
7	P	3002	GOL	C3-C2	-7.44	1.21	1.51
7	B	3001	GOL	C3-C2	-7.18	1.22	1.51
7	B	3001	GOL	O1-C1	4.76	1.62	1.42
7	P	3002	GOL	O1-C1	4.76	1.62	1.42
7	B	3001	GOL	O3-C3	3.41	1.56	1.42

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	P	3002	GOL	O3-C3-C2	10.56	160.82	110.20
7	B	3001	GOL	O3-C3-C2	10.49	160.49	110.20
7	B	3001	GOL	O2-C2-C3	6.77	138.92	109.12
7	P	3002	GOL	O2-C2-C3	6.66	138.45	109.12
7	P	3002	GOL	O1-C1-C2	3.23	125.71	110.20

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
7	P	3002	GOL	C1-C2-C3-O3
7	B	3001	GOL	O1-C1-C2-C3
7	B	3001	GOL	C1-C2-C3-O3
7	P	3002	GOL	O1-C1-C2-O2
7	P	3002	GOL	O1-C1-C2-C3

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	H	1725	GLC	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	T	24/27 (88%)	0.30	3 (12%) 3 2	43, 87, 120, 126	0
2	P	19/21 (90%)	-0.12	1 (5%) 26 17	60, 75, 109, 117	0
3	A	556/558 (99%)	0.17	23 (4%) 37 27	31, 73, 101, 109	1 (0%)
4	B	429/429 (100%)	-0.05	11 (2%) 56 46	26, 54, 101, 112	1 (0%)
5	L	211/211 (100%)	-0.07	1 (0%) 91 88	39, 64, 98, 104	0
6	H	225/225 (100%)	-0.18	6 (2%) 54 44	35, 55, 87, 109	0
All	All	1464/1471 (99%)	0.01	45 (3%) 49 39	26, 64, 101, 126	2 (0%)

The worst 5 of 45 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
6	H	141	GLN	5.9
6	H	138	SER	5.4
3	A	297	GLU	5.1
3	A	252	TRP	4.8
1	T	702	DT	3.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	2DA	P	822	20/21	0.97	0.20	51,54,57,58	0

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
7	GOL	B	3001	6/6	0.87	0.18	67,70,71,72	0
9	GLC	H	1725	12/12	0.90	0.16	61,63,65,67	0
7	GOL	P	3002	6/6	0.92	0.13	62,65,66,68	0
8	MG	A	559	1/1	0.97	0.27	37,37,37,37	0

6.5 Other polymers [i](#)

There are no such residues in this entry.