

Full wwPDB X-ray Structure Validation Report (i)

Aug 14, 2023 – 02:10 PM EDT

PDB ID	:	1R0E
Title	:	Glycogen synthase kinase-3 beta in complex with 3-indolyl-4-arylmaleimide
		inhibitor
Authors	:	Allard, J.; Nikolcheva, T.; Gong, L.; Wang, J.; Dunten, P.; Avnur, Z.; Waters,
		R.; Sun, Q.; Skinner, B.
Deposited on	:	2003-09-20
Resolution	:	2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
	$(\# {\it Entries})$	$(\# { m Entries}, { m resolution} { m range}({ m \AA}))$
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length	Quality of chain			
1	А	391	76%	11%	•	12%
1	В	391	75%	11%	••	12%



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5791 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	Δ	344	Total	С	Ν	0	\mathbf{S}	0	9	0
1	A	044	2760	1777	474	498	11	0	Δ	0
1	р	344	Total	С	Ν	0	S	0	2	0
1	D	044	2760	1777	474	498	11	0	2	0


• Molecule 1 is a protein called Glycogen synthase kinase-3 beta.

Chain	Residue	Modelled	Actual	Comment	Reference
А	30	GLY	-	cloning artifact	UNP P49841
А	31	ALA	-	cloning artifact	UNP P49841
А	32	MET	-	cloning artifact	UNP P49841
А	33	SER	-	cloning artifact	UNP P49841
А	34	LEU	-	cloning artifact	UNP P49841
В	30	GLY	-	cloning artifact	UNP P49841
В	31	ALA	-	cloning artifact	UNP P49841
В	32	MET	-	cloning artifact	UNP P49841
В	33	SER	-	cloning artifact	UNP P49841
В	34	LEU	-	cloning artifact	UNP P49841

There are 10 discrepancies between the modelled and reference sequences:


• Molecule 2 is CITRATE ANION (three-letter code: FLC) (formula: $C_6H_5O_7$).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total C O 13 6 7	0	0
2	В	1	Total C O 13 6 7	0	0

• Molecule 3 is 3-[3-(2,3-DIHYDROXY-PROPYLAMINO)-PHENYL]-4-(5-FLUORO-1-METHYL-1H-INDOL-3-YL)-PYRROLE-2,5-DIONE (three-letter code: DFN) (formula: C₂₂H₂₀FN₃O₄).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	Δ	1	Total	С	F	Ν	Ο	0	0
0	A	1	30	22	1	3	4	0	0



Continued from previous page...

Mol	Chain	Residues		Atc	oms			ZeroOcc	AltConf
2	В	1	Total	С	F	Ν	0	0	0
3	D		30	22	1	3	4		0

• Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	90	Total O 90 90	0	0
4	В	95	$\begin{array}{cc} \text{Total} & \text{O} \\ 95 & 95 \end{array}$	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.



• Molecule 1: Glycogen synthase kinase-3 beta



4 Data and refinement statistics (i)

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	83.96Å 84.01Å 180.20Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	91.29 - 2.25	Depositor
% Data completeness	98.4 (91.29-2.25)	Depositor
(in resolution range)	30.4 (31.23-2.20)	Depositor
R_{merge}	(Not available)	Depositor
R _{sym}	0.05	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.223 , 0.252	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	5791	wwPDB-VP
Average B, all atoms $(Å^2)$	44.0	wwPDB-VP



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: FLC, DFN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bo	ond angles
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.57	1/2839~(0.0%)	0.75	8/3865~(0.2%)
1	В	0.57	1/2839~(0.0%)	0.75	7/3865~(0.2%)
All	All	0.57	2/5678~(0.0%)	0.75	15/7730~(0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	218	CYS	CB-SG	-5.33	1.73	1.81
1	В	218	CYS	CB-SG	-5.02	1.73	1.81

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
1	В	105	ASP	CB-CG-OD2	6.72	124.34	118.30
1	А	260	ASP	CB-CG-OD2	6.52	124.17	118.30
1	В	233	ASP	CB-CG-OD2	6.45	124.11	118.30
1	А	264	ASP	CB-CG-OD2	6.02	123.71	118.30
1	А	49	ASP	CB-CG-OD2	5.91	123.62	118.30
1	А	233	ASP	CB-CG-OD2	5.66	123.39	118.30
1	А	341	ASP	CB-CG-OD2	5.51	123.26	118.30
1	В	341	ASP	CB-CG-OD2	5.42	123.18	118.30
1	А	90	ASP	CB-CG-OD2	5.37	123.14	118.30
1	В	49	ASP	CB-CG-OD2	5.34	123.10	118.30
1	А	105	ASP	CB-CG-OD2	5.26	123.03	118.30
1	А	192	ASP	CB-CG-OD2	5.15	122.93	118.30
1	В	190	ASP	CB-CG-OD2	5.11	122.90	118.30
1	В	192	ASP	CB-CG-OD2	5.09	122.89	118.30
1	В	264	ASP	CB-CG-OD2	5.06	122.85	118.30

All (15) bond angle outliers are listed below:

There are no chirality outliers.



There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2760	0	2772	16	0
1	В	2760	0	2772	17	0
2	А	13	0	5	0	0
2	В	13	0	5	0	0
3	А	30	0	20	1	0
3	В	30	0	20	0	0
4	А	90	0	0	0	0
4	В	95	0	0	0	0
All	All	5791	0	5594	32	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (32) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
		distance (Å)	overlap (Å)
1:B:217:ILE:O	1:B:218:CYS:HB2	1.84	0.78
1:A:218:CYS:HB3	1:A:223:ARG:HG2	1.69	0.72
1:B:218:CYS:HB3	1:B:223:ARG:HG2	1.73	0.70
1:A:217:ILE:O	1:A:218:CYS:HB2	1.90	0.69
1:A:67:PHE:O	1:A:85:LYS:HE3	1.96	0.65
1:A:263:VAL:HG21	1:B:217:ILE:HG22	1.82	0.62
1:B:144:ARG:O	1:B:148:ARG:HG3	2.03	0.58
1:B:217:ILE:O	1:B:218:CYS:CB	2.53	0.55
1:B:97:GLU:O	1:B:101:MET:HG2	2.08	0.54
1:B:370:ASN:HD22	1:B:372:PRO:HD2	1.72	0.53
1:A:155:VAL:HG12	1:A:159:LYS:HE3	1.92	0.52
1:A:370:ASN:HD22	1:A:372:PRO:HD2	1.76	0.51
1:B:60:LYS:HE2	1:B:72:GLN:NE2	2.27	0.50
1:A:217:ILE:O	1:A:218:CYS:CB	2.60	0.49
1:A:238:ILE:HG13	1:A:239:ASP:N	2.28	0.48
1:A:319:ARG:HB3	1:A:329:LEU:HG	1.95	0.48



Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:97:GLU:O	1:A:101:MET:HG2	2.14	0.47
1:A:85:LYS:HD2	3:A:702:DFN:H12	1.97	0.47
1:B:67:PHE:O	1:B:85:LYS:HE3	2.16	0.45
1:A:40:VAL:HG11	1:A:116:PHE:CE1	2.51	0.45
1:B:40:VAL:HG11	1:B:116:PHE:CE1	2.51	0.45
1:B:238:ILE:HG13	1:B:239:ASP:N	2.31	0.44
1:B:313:ALA:HB2	1:B:339:PHE:CE1	2.53	0.44
1:A:294:PRO:HB2	1:A:296:ILE:CD1	2.47	0.43
1:A:84:ILE:HA	1:A:130:LEU:O	2.18	0.43
1:B:84:ILE:HA	1:B:130:LEU:O	2.19	0.42
1:B:233:ASP:OD1	1:B:233:ASP:N	2.53	0.41
1:B:110:VAL:HG13	1:B:133:ASP:HB2	2.02	0.41
1:A:305:PHE:CD2	1:A:314:ILE:HG12	2.55	0.41
1:A:370:ASN:HD22	1:A:370:ASN:C	2.24	0.41
1:B:305:PHE:CD2	1:B:314:ILE:HG12	2.56	0.41
1:B:162:MET:HE1	1:B:243:ALA:O	2.21	0.41

Continued from previous page...

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	А	342/391~(88%)	329~(96%)	12~(4%)	1 (0%)	41	46
1	В	342/391~(88%)	328 (96%)	12~(4%)	2(1%)	25	25
All	All	684/782~(88%)	657~(96%)	24~(4%)	3~(0%)	34	37

All (3) Ramachandran outliers are listed below:

Mol	Chain	\mathbf{Res}	\mathbf{Type}
1	А	221	TYR
1	В	221	TYR



Continued from previous page...

Mol	Chain	\mathbf{Res}	Type
1	В	49	ASP

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	А	307/338~(91%)	290~(94%)	17 (6%)	21 21
1	В	307/338~(91%)	288 (94%)	19 (6%)	18 17
All	All	614/676~(91%)	578 (94%)	36~(6%)	19 19

All (36) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	36	LYS
1	А	46	GLN
1	А	49	ASP
1	А	60	LYS
1	А	88	LEU
1	А	89	GLN
1	А	90	ASP
1	А	144	ARG
1	А	209	ARG
1	А	218	CYS
1	А	232	THR
1	А	278	ARG
1	А	290	GLU
1	А	303	LYS
1	А	306	ARG
1	А	369	SER
1	А	370	ASN
1	В	36	LYS
1	В	46	GLN
1	В	49	ASP
1	В	72	GLN
1	В	88	LEU



Mol	Chain	Res	Type
1	В	89	GLN
1	В	94	LYS
1	В	144	ARG
1	В	151	GLN
1	В	209	ARG
1	В	218	CYS
1	В	232	THR
1	В	278	ARG
1	В	279	GLU
1	В	295	GLN
1	В	303	LYS
1	В	369	SER
1	В	370	ASN
1	В	383	ARG

Continued from previous page...

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	46	GLN
1	А	72	GLN
1	А	370	ASN
1	В	72	GLN
1	В	370	ASN
1	В	381	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.



5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2
2	FLC	В	601	-	12,12,12	1.02	0	17,17,17	1.44	2 (11%)
3	DFN	В	501	-	31,33,33	1.15	3 (9%)	38,48,48	0.94	1 (2%)
2	FLC	А	701	-	12,12,12	0.97	0	17,17,17	1.57	2 (11%)
3	DFN	А	702	-	31,33,33	1.10	2 (6%)	38,48,48	1.02	1 (2%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLC	В	601	-	-	0/16/16/16	-
3	DFN	В	501	-	-	0/11/31/31	0/4/4/4
2	FLC	А	701	-	-	0/16/16/16	-
3	DFN	А	702	-	-	0/11/31/31	0/4/4/4

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	В	501	DFN	C19-C01	-2.63	1.44	1.50
3	А	702	DFN	C19-C01	-2.60	1.44	1.50
3	А	702	DFN	C23-C29	2.54	1.40	1.36
3	В	501	DFN	C23-C29	2.46	1.40	1.36
3	В	501	DFN	C19-C20	2.37	1.44	1.42

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^{o})$	$Ideal(^{o})$
2	А	701	FLC	OB2-CBC-CB	4.34	120.58	113.05
2	В	601	FLC	OB2-CBC-CB	4.01	120.01	113.05



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
2	А	701	FLC	OB1-CBC-CB	-3.32	117.56	122.25
2	В	601	FLC	OB1-CBC-CB	-3.04	117.95	122.25
3	А	702	DFN	C24-C21-N22	-2.80	129.74	132.14
3	В	501	DFN	C24-C21-N22	-2.23	130.22	132.14

Continued from previous page...

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.

1 monomer is involved in 1 short contact:

Μ	[ol	Chain	Res	Type	Clashes	Symm-Clashes
,	3	А	702	DFN	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates (i)

EDS was not executed - this section is therefore empty.

6.4 Ligands (i)

EDS was not executed - this section is therefore empty.

6.5 Other polymers (i)

EDS was not executed - this section is therefore empty.

