



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 14, 2024 – 04:48 am GMT

PDB ID : 6R2D
Title : Crystal structure of the SucA domain of Mycobacterium smegmatis KGD after soaking with succinylphosphonate phosphonoethyl ester, followed by temperature increase
Authors : Wagner, T.; Alzari, P.M.; Bellinzoni, M.
Deposited on : 2019-03-15
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	868	 5% 91% 6%
1	B	868	 3% 91% 6%
1	C	868	 6% 90% 7%
1	D	868	 3% 89% 7%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 26221 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Multifunctional 2-oxoglutarate metabolism enzyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	814	6306	3974	1111	1197	24	0	1	0
1	B	812	6243	3938	1108	1172	25	0	1	0
1	C	808	6271	3949	1109	1191	22	0	0	0
1	D	811	6240	3934	1096	1186	24	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	360	GLY	-	expression tag	UNP A0R2B1
B	360	GLY	-	expression tag	UNP A0R2B1
C	360	GLY	-	expression tag	UNP A0R2B1
D	360	GLY	-	expression tag	UNP A0R2B1

- Molecule 2 is (4 {S})-4-[(2 {R})-3-[(4-azanyl-2-methyl-pyrimidin-5-yl)methyl]-4-methyl-5-[2-[oxidanyl(phosphonoxy)phosphoryl]oxyethyl]-2 {H}-1,3-thiazol-2-yl]-4-[ethoxy(oxidanyl)phosphoryl]-4-oxidanyl-butanoic acid (three-letter code: ZP1) (formula: C₁₈H₃₁N₄O₁₃P₃S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	N	O	P			S
2	A	1	Total	C	N	O	P	S	0	0
			39	18	4	13	3	1		
2	B	1	Total	C	N	O	P	S	0	0
			39	18	4	13	3	1		
2	C	1	Total	C	N	O	P	S	0	0
			39	18	4	13	3	1		
2	D	1	Total	C	N	O	P	S	0	0
			39	18	4	13	3	1		

- Molecule 3 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mg		
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		
3	C	1	Total	Mg	0	0
			1	1		
3	D	1	Total	Mg	0	0
			1	1		

- Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Ca		
4	A	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total Ca 1 1	0	0
4	C	1	Total Ca 1 1	0	0
4	D	1	Total Ca 1 1	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	256	Total O 256 256	0	0
5	B	260	Total O 260 260	0	0
5	C	251	Total O 251 251	0	0
5	D	230	Total O 230 230	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme




- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme




- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme





- Molecule 1: Multifunctional 2-oxoglutarate metabolism enzyme

Chain D: 3% 89% 7%



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	80.84Å 83.73Å 160.57Å 99.52° 98.94° 100.81°	Depositor
Resolution (Å)	33.44 – 2.30 77.92 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.2 (33.44-2.30) 97.2 (77.92-2.30)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.29Å)	Xtrriage
Refinement program	BUSTER 2.10.3	Depositor
R, R_{free}	0.217 , 0.242 0.220 , 0.246	Depositor DCC
R_{free} test set	8614 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	47.0	Xtrriage
Anisotropy	0.308	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 61.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.000 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	26221	wwPDB-VP
Average B, all atoms (Å ²)	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.73 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.3690e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZP1, CA, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/6436	0.63	0/8727
1	B	0.50	0/6373	0.63	0/8649
1	C	0.51	0/6397	0.64	0/8673
1	D	0.50	0/6369	0.63	0/8643
All	All	0.50	0/25575	0.63	0/34692

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6306	0	6095	10	0
1	B	6243	0	6013	10	0
1	C	6271	0	6076	14	0
1	D	6240	0	6002	15	0
2	A	39	0	0	0	0
2	B	39	0	0	0	0
2	C	39	0	0	0	0
2	D	39	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	256	0	0	0	0
5	B	260	0	0	2	0
5	C	251	0	0	2	0
5	D	230	0	0	0	0
All	All	26221	0	24186	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:778:LEU:HB3	1:A:784:ILE:HG12	1.85	0.59
1:C:705:PRO:HG2	1:C:735:VAL:HG13	1.88	0.56
1:D:1093:VAL:HB	1:D:1150:ARG:HG2	1.89	0.55
1:B:1183:THR:O	1:B:1186:GLU:HG2	2.06	0.55
1:C:981:VAL:HG22	1:C:988:ILE:HD11	1.93	0.51
1:C:754:SER:HB2	1:D:857:HIS:HE2	1.75	0.51
1:D:778:LEU:HB3	1:D:784:ILE:HG12	1.94	0.49
1:B:1041[B]:MET:HE1	5:B:2345:HOH:O	2.12	0.48
1:C:505:ARG:HB2	5:C:2156:HOH:O	2.14	0.48
1:C:505:ARG:HG2	5:C:2168:HOH:O	2.13	0.48
1:D:771:ARG:HH12	1:D:793:LEU:HD23	1.80	0.47
1:D:981:VAL:HG22	1:D:988:ILE:HD11	1.96	0.47
1:B:981:VAL:HG22	1:B:988:ILE:HD11	1.95	0.46
1:C:1174:GLN:OE1	1:C:1206:PRO:HA	2.16	0.46
1:B:1174:GLN:OE1	1:B:1206:PRO:HA	2.16	0.46
1:C:753:PRO:HB2	1:C:761:TYR:CD2	2.51	0.46
1:A:1174:GLN:OE1	1:A:1206:PRO:HA	2.16	0.46
1:B:396:ARG:HD3	5:B:2356:HOH:O	2.15	0.46
1:B:896:VAL:HG22	1:B:970:MET:HB3	1.98	0.45
1:A:1177:TRP:CD1	1:A:1197:ARG:HD3	2.52	0.45
1:B:1177:TRP:CD1	1:B:1197:ARG:HD3	2.52	0.45
1:D:546:LEU:HD21	1:D:599:LEU:HD23	1.97	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:1177:TRP:CD1	1:D:1197:ARG:HD3	2.52	0.45
1:C:1177:TRP:CD1	1:C:1197:ARG:HD3	2.52	0.45
1:D:1130:GLU:HB2	1:D:1132:ARG:HG2	1.98	0.45
1:C:1055:HIS:HE1	1:C:1062:ARG:O	1.98	0.45
1:C:1112:LEU:HD22	1:C:1136:ALA:HB3	1.99	0.44
1:A:1130:GLU:HB2	1:A:1132:ARG:HG2	2.01	0.43
1:A:903:THR:O	1:A:911:ARG:HD2	2.19	0.43
1:D:505:ARG:HA	1:D:747:HIS:O	2.18	0.43
1:A:981:VAL:HG22	1:A:988:ILE:HD11	2.00	0.43
1:D:1055:HIS:HE1	1:D:1062:ARG:O	2.03	0.42
1:C:555:SER:O	1:C:559:SER:HB2	2.20	0.42
1:B:447:VAL:HG22	1:B:709:VAL:HG12	2.01	0.42
1:B:633:ASN:O	1:B:732:LYS:HE2	2.20	0.42
1:A:1055:HIS:HE1	1:A:1062:ARG:O	2.03	0.41
1:D:896:VAL:HG22	1:D:970:MET:HB3	2.02	0.41
1:A:898:LEU:O	1:A:945:VAL:HA	2.21	0.41
1:D:952:GLU:HG2	1:D:980:PHE:O	2.20	0.41
1:C:753:PRO:HB2	1:C:761:TYR:CE2	2.56	0.41
1:C:952:GLU:HG2	1:C:980:PHE:O	2.20	0.41
1:C:1130:GLU:HB2	1:C:1132:ARG:HG2	2.02	0.41
1:D:536:ALA:HB3	1:D:613:LEU:HD22	2.03	0.41
1:D:1174:GLN:OE1	1:D:1206:PRO:HA	2.20	0.41
1:A:952:GLU:HG2	1:A:980:PHE:O	2.21	0.41
1:B:1171:PRO:HB3	1:B:1201:ARG:O	2.20	0.41
1:A:555:SER:O	1:A:559:SER:HB2	2.22	0.40
1:D:447:VAL:HG22	1:D:709:VAL:HG23	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	805/868 (93%)	782 (97%)	22 (3%)	1 (0%)	51	64
1	B	803/868 (92%)	784 (98%)	19 (2%)	0	100	100
1	C	798/868 (92%)	777 (97%)	20 (2%)	1 (0%)	51	64
1	D	802/868 (92%)	783 (98%)	18 (2%)	1 (0%)	51	64
All	All	3208/3472 (92%)	3126 (97%)	79 (2%)	3 (0%)	51	64

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	1034	GLU
1	C	682	PHE
1	A	682	PHE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	650/726 (90%)	642 (99%)	8 (1%)	71	84
1	B	635/726 (88%)	627 (99%)	8 (1%)	69	82
1	C	648/726 (89%)	639 (99%)	9 (1%)	67	81
1	D	638/726 (88%)	631 (99%)	7 (1%)	73	86
All	All	2571/2904 (88%)	2539 (99%)	32 (1%)	71	84

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	539	HIS
1	A	682	PHE
1	A	744	ARG
1	A	810	LEU
1	A	831	LEU
1	A	953	PHE
1	A	976	GLN
1	A	1112	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	490	GLU
1	B	539	HIS
1	B	682	PHE
1	B	909	THR
1	B	953	PHE
1	B	976	GLN
1	B	1112	LEU
1	B	1220	GLU
1	C	466	ARG
1	C	490	GLU
1	C	539	HIS
1	C	682	PHE
1	C	855	THR
1	C	953	PHE
1	C	976	GLN
1	C	1112	LEU
1	C	1114	LEU
1	D	539	HIS
1	D	682	PHE
1	D	808	ARG
1	D	909	THR
1	D	953	PHE
1	D	976	GLN
1	D	1112	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	539	HIS
1	B	539	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ZP1	D	2001	3	34,40,40	2.52	8 (23%)	45,61,61	1.75	9 (20%)
2	ZP1	A	2001	3	34,40,40	2.48	10 (29%)	45,61,61	1.85	11 (24%)
2	ZP1	C	2001	3	34,40,40	2.36	8 (23%)	45,61,61	1.87	10 (22%)
2	ZP1	B	2001	3	34,40,40	2.54	11 (32%)	45,61,61	1.73	11 (24%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ZP1	D	2001	3	-	4/33/58/58	0/2/2/2
2	ZP1	A	2001	3	-	6/33/58/58	0/2/2/2
2	ZP1	C	2001	3	-	4/33/58/58	0/2/2/2
2	ZP1	B	2001	3	-	4/33/58/58	0/2/2/2

All (37) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	ZP1	P4-O8	10.07	1.72	1.57
2	D	2001	ZP1	P4-O8	10.07	1.72	1.57
2	B	2001	ZP1	P4-O8	9.37	1.71	1.57
2	C	2001	ZP1	P4-O8	9.32	1.71	1.57
2	B	2001	ZP1	P4-O5	-5.09	1.46	1.56
2	D	2001	ZP1	P4-O5	-4.60	1.47	1.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	2001	ZP1	C4-N3	4.33	1.48	1.36
2	B	2001	ZP1	C4-N3	4.17	1.47	1.36
2	D	2001	ZP1	C4-N3	4.10	1.47	1.36
2	D	2001	ZP1	PA-O7	4.07	1.75	1.59
2	C	2001	ZP1	PA-O7	4.04	1.75	1.59
2	B	2001	ZP1	PA-O7	3.86	1.74	1.59
2	A	2001	ZP1	PA-O7	3.76	1.74	1.59
2	C	2001	ZP1	C4-N3	3.63	1.46	1.36
2	C	2001	ZP1	P4-O5	-3.31	1.49	1.56
2	A	2001	ZP1	C2-N3	3.25	1.50	1.46
2	A	2001	ZP1	P4-O5	-3.15	1.50	1.56
2	C	2001	ZP1	C2-N3	3.10	1.49	1.46
2	B	2001	ZP1	C6-C7	2.95	1.59	1.51
2	B	2001	ZP1	C2-N3	2.94	1.49	1.46
2	C	2001	ZP1	C7'-N3	2.78	1.51	1.46
2	D	2001	ZP1	C6-C7	2.76	1.58	1.51
2	A	2001	ZP1	C6-C7	2.75	1.58	1.51
2	C	2001	ZP1	C4'-N3'	2.60	1.38	1.35
2	B	2001	ZP1	C7'-N3	2.54	1.50	1.46
2	D	2001	ZP1	C2-N3	2.45	1.49	1.46
2	B	2001	ZP1	O11-C11	-2.45	1.41	1.43
2	A	2001	ZP1	O11-C11	-2.41	1.41	1.43
2	C	2001	ZP1	C6-C7	2.40	1.57	1.51
2	A	2001	ZP1	C4'-N3'	2.39	1.38	1.35
2	D	2001	ZP1	C7'-N3	2.28	1.50	1.46
2	A	2001	ZP1	CM4-C4	2.24	1.53	1.49
2	B	2001	ZP1	O16-C14	-2.24	1.23	1.30
2	B	2001	ZP1	C4'-N3'	2.15	1.38	1.35
2	A	2001	ZP1	C7'-N3	2.13	1.50	1.46
2	B	2001	ZP1	O7-C7	-2.11	1.36	1.44
2	D	2001	ZP1	C4'-N3'	2.06	1.38	1.35

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	2001	ZP1	C5'-C7'-N3	5.59	121.35	113.34
2	A	2001	ZP1	C5'-C7'-N3	5.39	121.05	113.34
2	D	2001	ZP1	C5'-C7'-N3	5.16	120.73	113.34
2	D	2001	ZP1	O8-P4-O6	-4.84	106.89	115.48
2	B	2001	ZP1	C5'-C7'-N3	4.73	120.11	113.34
2	C	2001	ZP1	O8-P4-O6	-4.68	107.19	115.48
2	A	2001	ZP1	O8-P4-O6	-4.63	107.27	115.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	2001	ZP1	O8-P4-O6	-4.44	107.61	115.48
2	C	2001	ZP1	S1-C2-N3	-3.51	104.01	106.91
2	A	2001	ZP1	S1-C2-N3	-3.36	104.14	106.91
2	B	2001	ZP1	C13-C12-C11	3.27	122.19	114.33
2	D	2001	ZP1	C13-C12-C11	3.23	122.10	114.33
2	C	2001	ZP1	CM4-C4-N3	3.18	122.92	119.89
2	C	2001	ZP1	C13-C12-C11	3.17	121.95	114.33
2	A	2001	ZP1	C13-C12-C11	3.08	121.74	114.33
2	A	2001	ZP1	P4-C11-O11	2.93	109.95	104.90
2	B	2001	ZP1	CM4-C4-N3	2.92	122.67	119.89
2	A	2001	ZP1	CM4-C4-N3	2.87	122.63	119.89
2	C	2001	ZP1	P4-C11-O11	2.75	109.64	104.90
2	A	2001	ZP1	O16-C14-C13	2.60	122.38	114.03
2	C	2001	ZP1	O16-C14-C13	2.60	122.37	114.03
2	B	2001	ZP1	O16-C14-C13	2.59	122.36	114.03
2	D	2001	ZP1	S1-C2-N3	-2.58	104.78	106.91
2	B	2001	ZP1	S1-C2-N3	-2.57	104.79	106.91
2	D	2001	ZP1	CM4-C4-N3	2.55	122.32	119.89
2	A	2001	ZP1	O5-P4-O6	2.40	117.12	111.34
2	B	2001	ZP1	O5-P4-O6	2.35	117.02	111.34
2	D	2001	ZP1	P4-C11-O11	2.34	108.94	104.90
2	D	2001	ZP1	O16-C14-C13	2.33	121.53	114.03
2	B	2001	ZP1	P4-C11-O11	2.33	108.92	104.90
2	D	2001	ZP1	O5-P4-O6	2.31	116.91	111.34
2	B	2001	ZP1	PA-O3A-PB	-2.20	125.29	132.83
2	A	2001	ZP1	N4'-C4'-N3'	-2.18	113.96	117.03
2	B	2001	ZP1	O7-PA-O2A	-2.16	100.64	109.07
2	A	2001	ZP1	O3B-PB-O2B	2.16	115.88	107.64
2	C	2001	ZP1	N4'-C4'-N3'	-2.14	114.01	117.03
2	C	2001	ZP1	O5-P4-O6	2.09	116.38	111.34
2	C	2001	ZP1	O15-C14-C13	-2.08	116.40	123.08
2	D	2001	ZP1	O7-PA-O2A	-2.05	101.07	109.07
2	B	2001	ZP1	N4'-C4'-N3'	-2.04	114.15	117.03
2	A	2001	ZP1	PA-O3A-PB	-2.01	125.93	132.83

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	2001	ZP1	C8-O8-P4-O5
2	B	2001	ZP1	C8-O8-P4-O5
2	C	2001	ZP1	C8-O8-P4-O5

Continued on next page...


Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	D	2001	ZP1	C8-O8-P4-O5
2	A	2001	ZP1	P4-C11-C12-C13
2	B	2001	ZP1	P4-C11-C12-C13
2	C	2001	ZP1	P4-C11-C12-C13
2	D	2001	ZP1	P4-C11-C12-C13
2	A	2001	ZP1	O11-C11-C12-C13
2	B	2001	ZP1	O11-C11-C12-C13
2	C	2001	ZP1	O11-C11-C12-C13
2	D	2001	ZP1	O11-C11-C12-C13
2	A	2001	ZP1	C2-C11-C12-C13
2	B	2001	ZP1	C2-C11-C12-C13
2	C	2001	ZP1	C2-C11-C12-C13
2	D	2001	ZP1	C2-C11-C12-C13
2	A	2001	ZP1	PA-O3A-PB-O3B
2	A	2001	ZP1	C9-C8-O8-P4


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	814/868 (93%)	0.45	44 (5%) 25 32	34, 54, 87, 121	0
1	B	812/868 (93%)	0.35	23 (2%) 53 60	35, 55, 90, 123	0
1	C	808/868 (93%)	0.51	49 (6%) 21 27	36, 56, 90, 121	0
1	D	811/868 (93%)	0.45	23 (2%) 53 60	35, 56, 88, 116	0
All	All	3245/3472 (93%)	0.44	139 (4%) 35 42	34, 56, 89, 123	0

All (139) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	574	GLY	7.0
1	D	810	LEU	6.5
1	C	775	THR	6.1
1	C	500	TYR	6.1
1	B	810	LEU	6.0
1	A	1083	ARG	5.8
1	A	792	ALA	5.6
1	C	810	LEU	5.4
1	C	1083	ARG	5.0
1	A	561	PHE	4.9
1	B	501	VAL	4.8
1	C	769	GLY	4.8
1	A	796	TYR	4.7
1	D	413	TRP	4.7
1	C	779	ILE	4.7
1	C	805	ASN	4.6
1	B	371	ILE	4.5
1	A	506	PHE	4.3
1	B	1102	GLY	4.3
1	A	810	LEU	4.3
1	C	506	PHE	4.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	805	ASN	4.2
1	A	807	VAL	4.2
1	D	501	VAL	4.2
1	D	797	GLN	4.2
1	D	777	ALA	4.2
1	D	793	LEU	4.1
1	C	764	ILE	4.0
1	C	792	ALA	4.0
1	D	368	ALA	4.0
1	C	481	TYR	3.9
1	A	793	LEU	3.8
1	A	784	ILE	3.7
1	C	807	VAL	3.7
1	A	500	TYR	3.7
1	C	1210	SER	3.7
1	A	750	GLY	3.6
1	C	804	PHE	3.6
1	B	394	ASN	3.6
1	C	812	LYS	3.6
1	B	581	GLY	3.6
1	D	807	VAL	3.6
1	C	796	TYR	3.5
1	A	855	THR	3.5
1	A	800	LEU	3.5
1	A	803	VAL	3.5
1	C	394	ASN	3.4
1	C	800	LEU	3.3
1	C	489	ALA	3.3
1	D	397	PHE	3.3
1	A	775	THR	3.2
1	D	800	LEU	3.2
1	B	368	ALA	3.1
1	B	777	ALA	3.1
1	A	804	PHE	3.1
1	C	799	GLN	3.1
1	C	793	LEU	3.1
1	C	785	SER	3.1
1	A	812	LYS	3.0
1	B	807	VAL	3.0
1	C	495	PHE	3.0
1	C	803	VAL	2.9
1	A	909	THR	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	769	GLY	2.9
1	C	371	ILE	2.9
1	C	576	VAL	2.9
1	B	783	ASP	2.9
1	C	750	GLY	2.9
1	B	397	PHE	2.8
1	B	432	LEU	2.8
1	C	551	GLY	2.7
1	C	1213	VAL	2.7
1	A	765	ASP	2.6
1	D	471	HIS	2.6
1	C	774	TYR	2.6
1	D	626	THR	2.6
1	C	784	ILE	2.6
1	B	801	GLU	2.5
1	B	796	TYR	2.5
1	C	580	LEU	2.5
1	A	575	ASP	2.4
1	A	410	LEU	2.4
1	C	633	ASN	2.4
1	D	775	THR	2.4
1	D	472	ASP	2.4
1	D	559	SER	2.4
1	C	778	LEU	2.4
1	B	502	GLY	2.4
1	D	805	ASN	2.4
1	A	392	LEU	2.4
1	C	547	ALA	2.4
1	A	491	ALA	2.4
1	C	501	VAL	2.4
1	A	1210	SER	2.4
1	B	804	PHE	2.3
1	A	840	LEU	2.3
1	A	688	ASP	2.3
1	A	495	PHE	2.3
1	A	581	GLY	2.3
1	D	581	GLY	2.3
1	D	630	GLY	2.3
1	A	547	ALA	2.3
1	A	580	LEU	2.3
1	B	809	GLU	2.2
1	A	488	ALA	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	844	GLY	2.2
1	B	413	TRP	2.2
1	A	371	ILE	2.2
1	C	1034	GLU	2.2
1	B	1101	ASP	2.2
1	A	1082	ILE	2.2
1	A	790	GLU	2.2
1	C	1035	GLY	2.2
1	B	395	THR	2.2
1	A	785	SER	2.2
1	B	472	ASP	2.2
1	D	367	ASN	2.1
1	B	840	LEU	2.1
1	C	392	LEU	2.1
1	A	799	GLN	2.1
1	C	595	ILE	2.1
1	C	546	LEU	2.1
1	B	782	GLY	2.1
1	A	774	TYR	2.1
1	C	368	ALA	2.1
1	C	749	GLU	2.1
1	D	667	ARG	2.1
1	C	474	PRO	2.1
1	C	419	PHE	2.1
1	C	578	TYR	2.1
1	D	1210	SER	2.1
1	A	802	ARG	2.0
1	C	702	ILE	2.0
1	A	399	SER	2.0
1	A	539	HIS	2.0
1	C	808	ARG	2.0
1	A	702	ILE	2.0
1	D	858	PRO	2.0
1	A	499	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	ZP1	C	2001	39/39	0.95	0.17	36,44,61,63	0
2	ZP1	B	2001	39/39	0.97	0.18	34,42,54,57	0
2	ZP1	A	2001	39/39	0.97	0.16	34,44,59,60	0
2	ZP1	D	2001	39/39	0.97	0.17	34,44,58,62	0
3	MG	A	2002	1/1	0.97	0.14	37,37,37,37	0
3	MG	B	2002	1/1	0.98	0.10	36,36,36,36	0
3	MG	C	2002	1/1	0.98	0.13	35,35,35,35	0
3	MG	D	2002	1/1	0.98	0.10	40,40,40,40	0
4	CA	B	2003	1/1	0.98	0.10	60,60,60,60	0
4	CA	C	2003	1/1	0.98	0.05	51,51,51,51	0
4	CA	D	2003	1/1	0.98	0.11	58,58,58,58	0
4	CA	A	2003	1/1	0.99	0.08	54,54,54,54	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around ZP1 C 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ZP1 B 2001:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around ZP1 A 2001:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.