



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 21, 2024 – 11:42 PM EST

PDB ID : 4R39
Title : Histidine kinase domain from Erythrobacter litoralis EL346 blue-light activated histidine kinase
Authors : Tomchick, D.R.; Rivera-Cancel, G.; Gardner, K.H.
Deposited on : 2014-08-14
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3163 (2.60-2.60)
Clashscore	141614	3518 (2.60-2.60)
Ramachandran outliers	138981	3455 (2.60-2.60)
Sidechain outliers	138945	3455 (2.60-2.60)
RSRZ outliers	127900	3104 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	 88% 5% 6%
1	B	232	 84% 12%
1	C	232	 82% 6% 12%
1	D	232	 89% 6%

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 13273 atoms, of which 6525 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blue-light-activated histidine kinase 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	217	3296	1003	1662	305	321	5	0	0	0
1	B	205	3120	949	1577	286	303	5	0	0	0
1	C	204	3108	945	1573	285	300	5	0	0	0
1	D	217	3295	1003	1661	305	321	5	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	115	GLY	-	expression tag	UNP Q2NB77
A	116	GLU	-	expression tag	UNP Q2NB77
A	117	PHE	-	expression tag	UNP Q2NB77
A	118	LYS	-	expression tag	UNP Q2NB77
A	119	GLY	-	expression tag	UNP Q2NB77
A	120	LEU	-	expression tag	UNP Q2NB77
B	115	GLY	-	expression tag	UNP Q2NB77
B	116	GLU	-	expression tag	UNP Q2NB77
B	117	PHE	-	expression tag	UNP Q2NB77
B	118	LYS	-	expression tag	UNP Q2NB77
B	119	GLY	-	expression tag	UNP Q2NB77
B	120	LEU	-	expression tag	UNP Q2NB77
C	115	GLY	-	expression tag	UNP Q2NB77
C	116	GLU	-	expression tag	UNP Q2NB77
C	117	PHE	-	expression tag	UNP Q2NB77
C	118	LYS	-	expression tag	UNP Q2NB77
C	119	GLY	-	expression tag	UNP Q2NB77
C	120	LEU	-	expression tag	UNP Q2NB77
D	115	GLY	-	expression tag	UNP Q2NB77
D	116	GLU	-	expression tag	UNP Q2NB77
D	117	PHE	-	expression tag	UNP Q2NB77

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	118	LYS	-	expression tag	UNP Q2NB77
D	119	GLY	-	expression tag	UNP Q2NB77
D	120	LEU	-	expression tag	UNP Q2NB77

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Mg 1 1	0	0
2	B	1	Total Mg 1 1	0	0
2	C	1	Total Mg 1 1	0	0
2	D	1	Total Mg 1 1	0	0

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C₁₀H₁₇N₆O₁₂P₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C H N O P 44 10 13 6 12 3	0	0
3	B	1	Total C H N O P 44 10 13 6 12 3	0	0
3	C	1	Total C H N O P 44 10 13 6 12 3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
3	D	1	44	10	13	6	12	3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	82	Total	O	0	0
			82	82		
4	B	68	Total	O	0	0
			68	68		
4	C	54	Total	O	0	0
			54	54		
4	D	70	Total	O	0	0
			70	70		

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Blue-light-activated histidine kinase 2

Chain A: 





- Molecule 1: Blue-light-activated histidine kinase 2

Chain B: 




- Molecule 1: Blue-light-activated histidine kinase 2


Chain C: 



THR
LYS
ASP

- Molecule 1: Blue-light-activated histidine kinase 2

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	120.50Å 120.50Å 192.30Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.25 – 2.60 41.25 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.7 (41.25-2.60) 98.7 (41.25-2.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.07 (at 2.61Å)	Xtrriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, R_{free}	0.183 , 0.221 0.185 , 0.223	Depositor DCC
R_{free} test set	2180 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	47.3	Xtrriage
Anisotropy	0.035	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 43.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13273	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.91% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ANP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/1649	0.52	0/2230
1	B	0.23	0/1555	0.47	0/2104
1	C	0.24	0/1547	0.49	0/2093
1	D	0.24	0/1649	0.48	0/2230
All	All	0.24	0/6400	0.49	0/8657

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1634	1662	1658	6	0
1	B	1543	1577	1573	5	0
1	C	1535	1573	1569	7	0
1	D	1634	1661	1658	5	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	31	13	13	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	B	31	13	13	0	0
3	C	31	13	13	0	0
3	D	31	13	13	0	0
4	A	82	0	0	2	0
4	B	68	0	0	0	0
4	C	54	0	0	1	0
4	D	70	0	0	0	0
All	All	6748	6525	6510	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (22) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:258:ASP:OD2	1:D:285:ARG:NH1	2.09	0.85
1:C:316:ASN:ND2	4:C:534:HOH:O	2.27	0.68
1:C:218:ARG:NH2	1:C:259:ARG:O	2.29	0.66
1:D:218:ARG:NH2	1:D:259:ARG:O	2.34	0.61
1:C:289:GLU:OE1	1:C:289:GLU:N	2.34	0.60
1:A:305:ARG:NH1	4:A:538:HOH:O	2.38	0.56
1:B:132:GLU:OE2	1:B:135:ARG:NH1	2.42	0.48
1:B:336:LYS:HG3	1:B:337:THR:N	2.30	0.47
1:A:258:ASP:OD1	1:A:258:ASP:N	2.47	0.46
1:A:282:ASP:HB2	1:A:324:THR:H	1.81	0.46
1:A:174:ARG:HD2	1:A:249:THR:OG1	2.16	0.45
1:A:302:ARG:NH1	4:A:537:HOH:O	2.45	0.44
1:D:212:ILE:HG22	1:D:252:LEU:HD12	1.99	0.44
1:C:174:ARG:HD3	1:C:212:ILE:HD12	2.00	0.44
1:B:212:ILE:HG22	1:B:252:LEU:HD12	2.01	0.43
1:D:143:ARG:HG3	1:D:186:MET:CE	2.49	0.42
1:C:134:ASP:OD2	1:C:156:ARG:NH2	2.48	0.42
1:B:283:ASP:OD2	1:D:272:SER:OG	2.35	0.42
1:C:243:LEU:HD11	1:C:304:VAL:HG13	2.03	0.41
1:A:182:LEU:HG	1:A:186:MET:CE	2.51	0.41
1:B:174:ARG:HA	1:B:174:ARG:NE	2.36	0.41
1:C:144:VAL:HG12	1:C:145:LYS:N	2.35	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	215/232 (93%)	206 (96%)	8 (4%)	1 (0%)	29	52
1	B	201/232 (87%)	193 (96%)	8 (4%)	0	100	100
1	C	200/232 (86%)	193 (96%)	7 (4%)	0	100	100
1	D	215/232 (93%)	206 (96%)	9 (4%)	0	100	100
All	All	831/928 (90%)	798 (96%)	32 (4%)	1 (0%)	51	75

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	165	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	176/188 (94%)	174 (99%)	2 (1%)	73	88
1	B	167/188 (89%)	165 (99%)	2 (1%)	71	87
1	C	166/188 (88%)	165 (99%)	1 (1%)	86	95
1	D	176/188 (94%)	175 (99%)	1 (1%)	86	95
All	All	685/752 (91%)	679 (99%)	6 (1%)	78	91

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	159	SER
1	A	173	SER
1	B	171	SER
1	B	187	ASP
1	C	171	SER
1	D	303	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	316	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ANP	D	402	2	29,33,33	1.06	4 (13%)	31,52,52	1.09	3 (9%)
3	ANP	B	402	2	29,33,33	1.08	4 (13%)	31,52,52	1.10	4 (12%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ANP	C	402	2	29,33,33	1.07	4 (13%)	31,52,52	1.05	2 (6%)
3	ANP	A	402	2	29,33,33	1.04	3 (10%)	31,52,52	1.08	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	D	402	2	-	5/14/38/38	0/3/3/3
3	ANP	B	402	2	-	4/14/38/38	0/3/3/3
3	ANP	C	402	2	-	7/14/38/38	0/3/3/3
3	ANP	A	402	2	-	4/14/38/38	0/3/3/3

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	402	ANP	PG-O1G	2.51	1.50	1.46
3	A	402	ANP	PG-N3B	2.50	1.69	1.63
3	C	402	ANP	PB-O1B	2.47	1.50	1.46
3	C	402	ANP	PG-O1G	2.46	1.50	1.46
3	B	402	ANP	PG-N3B	2.41	1.69	1.63
3	C	402	ANP	PG-N3B	2.41	1.69	1.63
3	D	402	ANP	PG-N3B	2.41	1.69	1.63
3	B	402	ANP	PB-O1B	2.32	1.49	1.46
3	B	402	ANP	PG-O1G	2.28	1.49	1.46
3	B	402	ANP	PB-O3A	-2.28	1.56	1.59
3	D	402	ANP	PB-O1B	2.21	1.49	1.46
3	A	402	ANP	PG-O1G	2.21	1.49	1.46
3	A	402	ANP	PB-O1B	2.17	1.49	1.46
3	C	402	ANP	PB-O3A	-2.07	1.56	1.59
3	D	402	ANP	PB-O3A	-2.02	1.56	1.59

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	402	ANP	PB-O3A-PA	-3.21	121.31	132.62
3	A	402	ANP	PB-O3A-PA	-3.09	121.72	132.62
3	D	402	ANP	PB-O3A-PA	-2.97	122.16	132.62
3	B	402	ANP	PB-O3A-PA	-2.56	123.60	132.62
3	D	402	ANP	O2B-PB-O3A	2.35	112.49	104.64

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	ANP	C5-C6-N6	2.35	123.92	120.35
3	A	402	ANP	C5-C6-N6	2.30	123.85	120.35
3	D	402	ANP	C5-C6-N6	2.29	123.84	120.35
3	C	402	ANP	C5-C6-N6	2.29	123.83	120.35
3	B	402	ANP	O1B-PB-N3B	-2.26	108.45	111.77
3	B	402	ANP	O2B-PB-O3A	2.22	112.06	104.64

There are no chirality outliers.

All (20) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	402	ANP	PB-N3B-PG-O1G
3	A	402	ANP	PA-O3A-PB-O2B
3	B	402	ANP	PB-N3B-PG-O1G
3	B	402	ANP	PG-N3B-PB-O3A
3	B	402	ANP	PA-O3A-PB-O1B
3	B	402	ANP	PA-O3A-PB-O2B
3	C	402	ANP	PB-N3B-PG-O1G
3	C	402	ANP	PB-O3A-PA-O5'
3	C	402	ANP	C5'-O5'-PA-O2A
3	D	402	ANP	PB-N3B-PG-O1G
3	D	402	ANP	PG-N3B-PB-O1B
3	D	402	ANP	PA-O3A-PB-O1B
3	D	402	ANP	PA-O3A-PB-O2B
3	C	402	ANP	C3'-C4'-C5'-O5'
3	C	402	ANP	O4'-C4'-C5'-O5'
3	C	402	ANP	C5'-O5'-PA-O3A
3	A	402	ANP	O4'-C4'-C5'-O5'
3	C	402	ANP	C5'-O5'-PA-O1A
3	A	402	ANP	C3'-C4'-C5'-O5'
3	D	402	ANP	PG-N3B-PB-O3A


There are no ring outliers.


No monomer is involved in short contacts.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring

in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	217/232 (93%)	-0.14	0 100 100	11, 27, 56, 82	0
1	B	205/232 (88%)	0.06	6 (2%) 51 45	12, 29, 68, 112	0
1	C	204/232 (87%)	0.11	4 (1%) 65 60	17, 36, 72, 95	0
1	D	217/232 (93%)	-0.02	5 (2%) 60 54	13, 25, 63, 90	0
All	All	843/928 (90%)	0.00	15 (1%) 68 64	11, 29, 67, 112	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	340	ALA	4.4
1	B	146	ASN	3.9
1	B	193	LYS	3.1
1	D	167	SER	3.1
1	B	290	ASP	3.0
1	C	146	ASN	2.5
1	C	339	ILE	2.4
1	B	194	ALA	2.4
1	C	147	HIS	2.3
1	D	128	ASP	2.1
1	C	226	GLN	2.1
1	B	338	LEU	2.1
1	D	191	ALA	2.1
1	D	193	LYS	2.1
1	D	168	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates

There are no monosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	MG	C	401	1/1	0.92	0.06	28,28,28,28	0
2	MG	B	401	1/1	0.94	0.04	20,20,20,20	0
3	ANP	C	402	31/31	0.96	0.11	17,34,44,55	0
2	MG	D	401	1/1	0.97	0.08	16,16,16,16	0
3	ANP	B	402	31/31	0.98	0.11	5,22,33,40	0
2	MG	A	401	1/1	0.98	0.09	3,3,3,3	0
3	ANP	D	402	31/31	0.98	0.12	7,16,23,25	0
3	ANP	A	402	31/31	0.99	0.14	3,16,27,32	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ANP C 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around ANP B 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around ANP D 402:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around ANP A 402:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.