



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 13, 2023 – 06:44 AM EDT

PDB ID : 4R3A  
Title : Erythrobacter litoralis EL346 blue-light activated histidine kinase  
Authors : Tomchick, D.R.; Rivera-Cancel, G.; Gardner, K.H.  
Deposited on : 2014-08-14  
Resolution : 2.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2307 (2.94-2.90)
Clashscore	141614	2531 (2.94-2.90)
Ramachandran outliers	138981	2462 (2.94-2.90)
Sidechain outliers	138945	2464 (2.94-2.90)
RSRZ outliers	127900	2248 (2.94-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	
1	B	352	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	RBF	B	402	-	-	-	X

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 9372 atoms, of which 4639 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Blue-light-activated histidine kinase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	318	4903	1525	2437	449	481	11	0	0	0
1	B	278	4285	1336	2136	386	416	11	0	0	0


There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP Q2NB77
A	-4	GLU	-	expression tag	UNP Q2NB77
A	-3	PHE	-	expression tag	UNP Q2NB77
A	-2	LYS	-	expression tag	UNP Q2NB77
A	-1	GLY	-	expression tag	UNP Q2NB77
A	0	LEU	-	expression tag	UNP Q2NB77
B	-5	GLY	-	expression tag	UNP Q2NB77
B	-4	GLU	-	expression tag	UNP Q2NB77
B	-3	PHE	-	expression tag	UNP Q2NB77
B	-2	LYS	-	expression tag	UNP Q2NB77
B	-1	GLY	-	expression tag	UNP Q2NB77
B	0	LEU	-	expression tag	UNP Q2NB77

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>6</sub>O<sub>12</sub>P<sub>3</sub>).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	H	N	O			P
3	A	1	Total	C	H	N	O	P	0	0
			44	10	13	6	12	3		
3	B	1	Total	C	H	N	O	P	0	0
			44	10	13	6	12	3		

- Molecule 4 is RIBOFLAVIN (three-letter code: RBF) (formula: C<sub>17</sub>H<sub>20</sub>N<sub>4</sub>O<sub>6</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
4	A	1	Total	C	H	N	O	0	0
			47	17	20	4	6		
4	B	1	Total	C	H	N	O	0	0
			47	17	20	4	6		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Blue-light-activated histidine kinase 2



- Molecule 1: Blue-light-activated histidine kinase 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	98.88Å 98.88Å 422.25Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	46.60 – 2.92 46.65 – 2.92	Depositor EDS
% Data completeness (in resolution range)	94.7 (46.60-2.92) 94.7 (46.65-2.92)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.42 (at 2.91Å)	Xtrriage
Refinement program	PHENIX	Depositor
R, $R_{free}$	0.290 , 0.330 0.293 , 0.332	Depositor DCC
$R_{free}$ test set	1693 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	53.1	Xtrriage
Anisotropy	0.073	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 61.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.85	EDS
Total number of atoms	9372	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	98.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, RBF, ANP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.29	0/2497	0.57	0/3370
1	B	0.24	0/2172	0.49	0/2931
All	All	0.27	0/4669	0.54	0/6301

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2466	2437	2437	19	0
1	B	2149	2136	2134	10	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	31	13	13	1	0
3	B	31	13	13	1	0
4	A	27	20	20	0	0
4	B	27	20	20	2	0
All	All	4733	4639	4637	26	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.



All (26) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:ALA:HB3	4:B:402:RBF:HC82	1.87	0.56
1:A:123:GLN:O	1:B:123:GLN:NE2	2.40	0.55
1:A:135:ARG:NH1	1:A:139:GLU:OE2	2.44	0.51
1:A:13:TRP:HB3	1:A:37:ARG:HG3	1.94	0.50
1:A:218:ARG:N	1:A:218:ARG:HD3	2.27	0.49
1:A:104:LEU:HG	1:A:152:VAL:HG11	1.94	0.48
1:A:37:ARG:CD	1:A:37:ARG:H	2.27	0.48
1:A:255:ALA:O	1:A:285:ARG:N	2.45	0.48
1:A:207:ARG:HG2	1:B:203:ALA:HB1	1.97	0.46
1:B:276:LEU:N	1:B:330:ILE:O	2.44	0.46
1:A:290:ASP:OD1	1:A:291:CYS:N	2.49	0.46
1:A:152:VAL:O	1:A:155:ILE:HG13	2.16	0.45
1:B:290:ASP:OD1	1:B:291:CYS:N	2.49	0.45
1:B:59:GLN:OE1	4:B:402:RBF:O4'	2.35	0.45
3:A:401:ANP:O2G	3:A:401:ANP:O3A	2.33	0.45
1:A:203:ALA:HB1	1:B:207:ARG:HG2	1.98	0.45
1:B:175:ARG:NH2	3:B:401:ANP:O1G	2.47	0.44
1:A:223:VAL:HB	1:B:229:THR:CG2	2.47	0.44
1:A:104:LEU:N	1:A:113:TYR:O	2.45	0.44
1:A:240:ILE:HG12	1:A:330:ILE:HD13	1.99	0.43
1:A:37:ARG:H	1:A:37:ARG:HD3	1.84	0.43
1:A:37:ARG:HH12	1:A:173:SER:HB3	1.83	0.42
1:A:104:LEU:HD12	1:A:104:LEU:HA	1.95	0.42
1:A:19:SER:HB3	1:A:118:GLN:HB2	2.02	0.42
1:B:198:ILE:HD11	1:B:234:VAL:CG2	2.51	0.41
1:A:176:VAL:HG12	1:A:176:VAL:O	2.21	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	310/352 (88%)	295 (95%)	11 (4%)	4 (1%)	12	36
1	B	264/352 (75%)	257 (97%)	7 (3%)	0	100	100
All	All	574/704 (82%)	552 (96%)	18 (3%)	4 (1%)	22	53

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	106	ASP
1	A	219	GLY
1	A	108	ASP
1	A	220	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/289 (92%)	263 (99%)	3 (1%)	73	91
1	B	234/289 (81%)	233 (100%)	1 (0%)	91	97
All	All	500/578 (86%)	496 (99%)	4 (1%)	81	93

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	37	ARG
1	A	111	CYS
1	A	218	ARG
1	B	196	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	RBF	A	402	-	29,29,29	1.27	2 (6%)	41,43,43	1.31	8 (19%)
3	ANP	B	401	2	29,33,33	1.10	4 (13%)	31,52,52	1.16	3 (9%)
4	RBF	B	402	-	29,29,29	1.30	2 (6%)	41,43,43	1.44	8 (19%)
3	ANP	A	401	2	29,33,33	1.07	4 (13%)	31,52,52	1.11	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	RBF	A	402	-	-	3/14/14/14	0/3/3/3
3	ANP	B	401	2	-	5/14/38/38	0/3/3/3
4	RBF	B	402	-	-	0/14/14/14	0/3/3/3
3	ANP	A	401	2	-	6/14/38/38	0/3/3/3

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	402	RBF	C5A-N5	-3.02	1.33	1.39

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	402	RBF	C5A-N5	-3.00	1.33	1.39
3	A	401	ANP	PB-O1B	2.85	1.50	1.46
3	B	401	ANP	PG-N3B	2.63	1.70	1.63
3	B	401	ANP	PG-O1G	2.52	1.50	1.46
3	B	401	ANP	PB-O1B	2.48	1.50	1.46
3	A	401	ANP	PG-N3B	2.24	1.69	1.63
3	A	401	ANP	PG-O1G	2.12	1.49	1.46
3	B	401	ANP	PB-O3A	-2.09	1.56	1.59
3	A	401	ANP	PB-O3A	-2.07	1.56	1.59
4	B	402	RBF	C4A-N5	2.06	1.34	1.30
4	A	402	RBF	O2'-C2'	-2.03	1.39	1.43

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	401	ANP	PB-O3A-PA	-3.45	120.47	132.62
3	A	401	ANP	PB-O3A-PA	-3.06	121.83	132.62
4	B	402	RBF	C4A-C4-N3	2.88	120.49	113.19
4	B	402	RBF	C4-N3-C2	-2.87	120.33	125.64
4	B	402	RBF	C4A-C10-N1	-2.75	118.36	124.73
3	B	401	ANP	C5-C6-N6	2.73	124.51	120.35
4	A	402	RBF	C4A-C4-N3	2.73	120.11	113.19
4	A	402	RBF	C4A-C10-N1	-2.71	118.44	124.73
4	A	402	RBF	C4-N3-C2	-2.69	120.67	125.64
4	B	402	RBF	O4-C4-C4A	-2.61	119.68	126.60
3	A	401	ANP	O1G-PG-N3B	-2.53	108.04	111.77
4	A	402	RBF	O4-C4-C4A	-2.52	119.90	126.60
3	A	401	ANP	C5-C6-N6	2.45	124.08	120.35
4	B	402	RBF	C10-N1-C2	2.40	121.71	116.90
4	A	402	RBF	C10-N1-C2	2.31	121.53	116.90
4	B	402	RBF	C4-C4A-N5	2.24	121.41	118.23
4	B	402	RBF	C10-C4A-N5	-2.20	120.19	124.86
4	B	402	RBF	C4A-C10-N10	2.18	119.66	116.48
4	A	402	RBF	C4A-C10-N10	2.17	119.65	116.48
4	A	402	RBF	C9A-N10-C10	-2.14	117.44	120.77
4	A	402	RBF	C5A-C9A-N10	2.11	120.14	117.95
3	B	401	ANP	O1G-PG-N3B	-2.05	108.75	111.77

There are no chirality outliers.

All (14) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	401	ANP	PB-N3B-PG-O1G
3	A	401	ANP	PB-O3A-PA-O5'
3	A	401	ANP	C5'-O5'-PA-O2A
3	B	401	ANP	PB-N3B-PG-O1G
3	B	401	ANP	PG-N3B-PB-O1B
3	B	401	ANP	PA-O3A-PB-O1B
3	B	401	ANP	PA-O3A-PB-O2B
4	A	402	RBF	N10-C1'-C2'-O2'
4	A	402	RBF	O4'-C4'-C5'-O5'
3	A	401	ANP	C5'-O5'-PA-O3A
3	A	401	ANP	C5'-O5'-PA-O1A
3	A	401	ANP	PG-N3B-PB-O1B
4	A	402	RBF	C3'-C4'-C5'-O5'
3	B	401	ANP	PG-N3B-PB-O3A


There are no ring outliers.


3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	401	ANP	1	0
4	B	402	RBF	2	0
3	A	401	ANP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	318/352 (90%)	0.68	34 (10%) <b>6</b> <b>4</b>	20, 54, 125, 170	0
1	B	278/352 (78%)	1.99	116 (41%) <b>0</b> <b>0</b>	69, 119, 168, 195	0
All	All	596/704 (84%)	1.29	150 (25%) <b>0</b> <b>0</b>	20, 87, 161, 195	0

All (150) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	111	CYS	7.4
1	B	104	LEU	7.4
1	B	50	VAL	6.9
1	B	287	ILE	6.4
1	B	186	MET	6.3
1	B	83	GLU	6.2
1	B	48	SER	6.1
1	A	165	VAL	6.1
1	B	34	TYR	6.1
1	B	53	ARG	6.0
1	B	45	TYR	5.9
1	A	261	SER	5.9
1	B	39	PHE	5.8
1	B	63	THR	5.4
1	A	319	ARG	5.3
1	A	320	GLY	5.3
1	A	8	HIS	5.2
1	B	28	ASP	5.0
1	B	47	ARG	5.0
1	A	337	THR	4.9
1	B	44	GLY	4.9
1	B	275	GLN	4.8
1	B	141	GLN	4.8
1	A	124	SER	4.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	29	ASP	4.5
1	B	123	GLN	4.5
1	B	79	GLU	4.5
1	B	30	GLU	4.4
1	B	155	ILE	4.2
1	B	51	VAL	4.1
1	B	220	ALA	4.1
1	B	116	GLY	4.1
1	B	156	ARG	4.1
1	B	84	THR	4.1
1	B	114	PHE	4.1
1	B	26	SER	4.1
1	B	322	THR	4.0
1	B	25	ILE	4.0
1	A	334	GLN	3.9
1	B	293	TRP	3.9
1	B	112	ARG	3.9
1	B	40	GLU	3.9
1	B	33	ILE	3.8
1	B	62	LYS	3.8
1	B	291	CYS	3.8
1	B	37	ARG	3.8
1	B	81	VAL	3.7
1	A	321	GLY	3.7
1	A	336	LYS	3.7
1	B	138	ALA	3.6
1	B	135	ARG	3.6
1	A	164	GLY	3.6
1	B	76	ARG	3.6
1	B	103	PRO	3.5
1	B	100	LEU	3.5
1	B	252	LEU	3.5
1	A	125	GLU	3.4
1	B	115	VAL	3.4
1	A	333	SER	3.4
1	B	35	VAL	3.4
1	B	65	PRO	3.4
1	B	154	MET	3.4
1	B	96	TRP	3.3
1	A	289	GLU	3.3
1	B	91	ASP	3.3
1	B	68	VAL	3.3

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	274	GLU	3.3
1	B	290	ASP	3.3
1	A	10	LYS	3.2
1	B	21	THR	3.2
1	B	147	HIS	3.2
1	B	38	ALA	3.2
1	A	338	LEU	3.2
1	B	189	ALA	3.1
1	B	97	ASN	3.1
1	B	73	LYS	3.1
1	B	316	ASN	3.1
1	B	324	THR	3.0
1	B	49	SER	3.0
1	B	292	ASP	3.0
1	B	57	PHE	2.9
1	A	275	GLN	2.9
1	B	77	ASN	2.9
1	A	162	ALA	2.9
1	A	260	ALA	2.9
1	A	322	THR	2.9
1	A	290	ASP	2.8
1	B	289	GLU	2.8
1	B	32	LEU	2.8
1	B	179	LEU	2.8
1	A	291	CYS	2.8
1	B	52	GLY	2.8
1	B	267	ARG	2.8
1	A	130	ALA	2.8
1	B	46	SER	2.7
1	B	36	ASN	2.7
1	B	22	ILE	2.7
1	B	82	GLU	2.7
1	B	86	TYR	2.7
1	B	176	VAL	2.7
1	B	71	LEU	2.7
1	A	166	GLY	2.7
1	B	198	ILE	2.6
1	B	325	ILE	2.6
1	B	137	LEU	2.6
1	B	288	PRO	2.6
1	B	248	LEU	2.5
1	A	220	ALA	2.5

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	307	LEU	2.5
1	B	72	ALA	2.5
1	B	172	LEU	2.5
1	B	218	ARG	2.5
1	B	69	GLU	2.4
1	B	85	ILE	2.4
1	B	80	GLU	2.4
1	B	247	VAL	2.4
1	A	335	GLN	2.4
1	B	202	GLY	2.4
1	B	334	GLN	2.4
1	A	129	ARG	2.4
1	B	263	VAL	2.4
1	B	24	ASP	2.4
1	B	323	GLY	2.4
1	A	121	MET	2.4
1	B	139	GLU	2.4
1	B	101	MET	2.4
1	B	75	ILE	2.4
1	B	261	SER	2.3
1	B	23	ALA	2.3
1	B	235	GLU	2.3
1	A	46	SER	2.3
1	B	136	GLN	2.3
1	B	317	VAL	2.3
1	B	271	MET	2.3
1	B	195	THR	2.3
1	A	136	GLN	2.3
1	B	318	THR	2.3
1	B	64	ASP	2.2
1	B	142	HIS	2.2
1	B	311	LEU	2.2
1	A	222	LYS	2.2
1	A	163	GLY	2.2
1	B	183	TYR	2.1
1	B	41	GLN	2.1
1	A	167	SER	2.1
1	B	19	SER	2.1
1	B	140	VAL	2.1
1	A	135	ARG	2.0
1	B	214	HIS	2.0
1	B	336	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
4	RBF	B	402	27/27	0.76	0.41	121,148,228,236	0
3	ANP	A	401	31/31	0.80	0.29	23,52,185,217	0
2	MG	A	400	1/1	0.83	0.26	24,24,24,24	0
3	ANP	B	401	31/31	0.84	0.18	42,106,139,160	0
4	RBF	A	402	27/27	0.93	0.18	12,37,74,86	0
2	MG	B	400	1/1	0.96	0.11	79,79,79,79	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around RBF B 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around ANP A 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around ANP B 401:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around RBF A 402:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.