

# wwPDB X-ray Structure Validation Summary Report (i)

#### Jan 14, 2024 - 07:46 am GMT

PDB ID	:	6R4V
Title	:	Crystal structure of human geranylgeranyl diphosphate synthase bound to
		ibandronate
Authors	:	Lisnyansky, M.; Giladi, M.; Haitin, Y.
Deposited on		
Resolution	:	2.20  Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Mogul	:	1.8.4, CSD as541be (2020)
Xtriage (Phenix)	:	1.13
EDS	:	2.36
buster-report	:	1.1.7 (2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:  $X\text{-}RAY\;DIFFRACTION$ 

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1	А	307	.% 84%	9%	• 6%
1	В	307	.% 87%	7%	6%
1	С	307	86%	9%	••
1	D	307	.% 87%	6%	7%
1	Е	307	87%	7%	6%



Continued from previous page...

Mol	Chain	Length	Quality of chain							
	-		% •							
1	F	307	85%	8%	7%					



# 2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 15063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace	
1	1 Λ	288	Total	С	Ν	0	S	0	0	0	
	А	200	2279	1469	377	426	7	0	0	0	
1	С	294	Total	С	Ν	Ο	S	0	0	0	
1	U	294	2322	1493	391	431	7	0	0	0	
1	F	286	Total	С	Ν	Ο	S	0	0	0	
1	Ľ	280	2252	1457	371	417	7	0	0	0	
1	Е	000	288	Total	С	Ν	Ο	S	0	0	0
1	Ľ	280	2281	1476	378	420	7	0	0	0	
1	В	290	Total	С	Ν	Ο	S	0	0	0	
1		D 290	2282	1477	378	420	7	0	0	0	
1	1 D	287	Total	С	Ν	Ο	S	0	0	0	
		201	2262	1463	375	417	7		0	0	

• Molecule 1 is a protein called Geranylgeranyl pyrophosphate synthase.

There are 48 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference						
A	-6	GLY	-	expression tag	UNP O95749						
А	-5	SER	-	expression tag	UNP O95749						
А	-4	GLY	-	expression tag	UNP 095749						
А	-3	SER	-	expression tag	UNP O95749						
А	-2	GLY	-	expression tag	UNP O95749						
А	-1	SER	-	expression tag	UNP 095749						
А	0	GLY	-	expression tag	UNP O95749						
А	109	GLN	PRO	variant	UNP 095749						
С	-6	GLY	-	expression tag	UNP O95749						
С	-5	SER	-	expression tag	UNP O95749						
С	-4	GLY	-	expression tag	UNP 095749						
С	-3	SER	-	expression tag	UNP O95749						
С	-2	GLY	-	expression tag	UNP 095749						
С	-1	SER	-	expression tag	UNP O95749						
С	0	GLY	-	expression tag	UNP O95749						
С	109	GLN	PRO	variant	UNP O95749						
F	-6	GLY	-	expression tag	UNP O95749						
	Continued on next nage										



Chain	Residue	Modelled	Actual	Comment	Reference
F	-5	SER	-	expression tag	UNP 095749
F	-4	GLY	-	expression tag	UNP 095749
F	-3	SER	-	expression tag	UNP 095749
F	-2	GLY	-	expression tag	UNP 095749
F	-1	SER	-	expression tag	UNP 095749
F	0	GLY	-	expression tag	UNP 095749
F	109	GLN	PRO	variant	UNP 095749
Е	-6	GLY	-	expression tag	UNP 095749
Е	-5	SER	-	expression tag	UNP 095749
Е	-4	GLY	-	expression tag	UNP 095749
Е	-3	SER	-	expression tag	UNP 095749
Е	-2	GLY	-	expression tag	UNP 095749
Е	-1	SER	-	expression tag	UNP 095749
Е	0	GLY	-	expression tag	UNP 095749
Е	109	GLN	PRO	variant	UNP 095749
В	-6	GLY	-	expression tag	UNP 095749
В	-5	SER	-	expression tag	UNP 095749
В	-4	GLY	-	expression tag	UNP 095749
В	-3	SER	-	expression tag	UNP 095749
В	-2	GLY	-	expression tag	UNP 095749
В	-1	SER	-	expression tag	UNP 095749
В	0	GLY	-	expression tag	UNP 095749
В	109	GLN	PRO	variant	UNP 095749
D	-6	GLY	-	expression tag	UNP 095749
D	-5	SER	-	expression tag	UNP 095749
D	-4	GLY	-	expression tag	UNP 095749
D	-3	SER	-	expression tag	UNP 095749
D	-2	GLY	-	expression tag	UNP 095749
D	-1	SER	-	expression tag	UNP 095749
D	0	GLY	-	expression tag	UNP 095749
D	109	GLN	PRO	variant	UNP 095749

Continued from previous page...

• Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	3	Total Mg 3 3	0	0
2	С	2	Total Mg 2 2	0	0
2	F	3	Total Mg 3 3	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	Е	3	Total Mg 3 3	0	0
2	В	3	Total Mg 3 3	0	0
2	D	3	Total Mg 3 3	0	0


• Molecule 3 is IBANDRONATE (three-letter code: BFQ) (formula:  $C_9H_{23}NO_7P_2$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	oms			ZeroOcc	AltConf	
3	А	٨	1	Total	С	Ν	0	Р	0	0
5	Л	I	19	9	1	$\overline{7}$	2	0	0	
3	С	1	Total	С	Ν	Ο	Р	0	0	
5	U	T	19	9	1	7	2	0	0	
3	F	1	Total	С	Ν	Ο	Р	0	0	
5	Ľ	1	19	9	1	7	2	0	0	
3	Е	1	Total	С	Ν	Ο	Р	0	0	
5	Ľ	1	19	9	1	7	2	0	0	
3	В	1	Total	С	Ν	Ο	Р	0	0	
5	D	T	19	9	1	7	2	0	0	
3	р	1	Total	С	Ν	Ō	Р	0	0	
5	D	L	19	9	1	7	2	0	0	

• Molecule 4 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 6  3  3 \end{array}$	0	0
4	С	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 6  3  3 \end{array}$	0	0


• Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	215	Total O 215 215	0	0
5	С	238	Total         O           238         238	0	0
5	F	167	Total O 167 167	0	0
5	Ε	229	Total O 229 229	0	0
5	В	205	Total O 205 205	0	0
5	D	188	Total O 188 188	0	0




# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.




• Molecule 1: Geranyl geranyl pyrophosphate synthase

• Molecule 1: Geranylgeranyl pyrophosphate synthase





#### GLU ASN GLU





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants	68.45Å 153.01Å 198.59Å	Depositor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $90.00^{\circ}$ $90.00^{\circ}$	Depositor
Resolution (Å)	49.65 - 2.20	Depositor
Resolution (A)	49.65 - 2.20	EDS
% Data completeness	97.6 (49.65-2.20)	Depositor
(in resolution range)	97.6(49.65-2.20)	EDS
R <sub>merge</sub>	(Not available)	Depositor
R <sub>sym</sub>	(Not available)	Depositor
$< I/\sigma(I) > 1$	$1.85 (at 2.20 \text{\AA})$	Xtriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
D D.	0.179 , $0.225$	Depositor
$R, R_{free}$	0.182 , $0.182$	DCC
$R_{free}$ test set	5187 reflections $(5.00\%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	30.3	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.32 , $47.4$	EDS
L-test for twinning <sup>2</sup>	$ \langle L  \rangle = 0.50, \langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	15063	wwPDB-VP
Average B, all atoms $(Å^2)$	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The analyses of the Patterson function reveals a significant off-origin peak that is 22.21 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.9728e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

# 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, GOL, BFQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bo	ond angles
	Unam	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.36	1/2328~(0.0%)	0.50	3/3160~(0.1%)
1	В	0.29	0/2332	0.48	2/3170~(0.1%)
1	С	0.31	0/2371	0.49	0/3217
1	D	0.30	0/2312	0.46	1/3144~(0.0%)
1	Е	0.31	0/2331	0.50	1/3164~(0.0%)
1	F	0.31	0/2302	0.47	2/3131~(0.1%)
All	All	0.31	1/13976~(0.0%)	0.48	9/18986~(0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	102	LYS	CE-NZ	7.33	1.67	1.49

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	Е	188	ASP	CB-CG-OD1	-7.95	111.14	118.30
1	F	112	VAL	CG1-CB-CG2	6.21	120.83	110.90
1	А	102	LYS	CA-CB-CG	5.80	126.15	113.40
1	В	112	VAL	CG1-CB-CG2	-5.74	101.71	110.90
1	D	121	GLU	CA-CB-CG	-5.63	101.02	113.40

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	2279	0	2199	26	0
1	В	2282	0	2183	15	0
1	С	2322	0	2236	23	0
1	D	2262	0	2171	12	0
1	Е	2281	0	2216	18	0
1	F	2252	0	2161	17	0
2	А	3	0	0	0	0
2	В	3	0	0	0	0
2	С	2	0	0	0	0
2	D	3	0	0	0	0
2	Е	3	0	0	0	0
2	F	3	0	0	0	0
3	А	19	0	19	2	0
3	В	19	0	19	2	0
3	С	19	0	19	1	0
3	D	19	0	19	3	0
3	Е	19	0	19	2	0
3	F	19	0	19	3	0
4	А	6	0	8	0	0
4	С	6	0	8	1	0
5	А	215	0	0	4	0
5	В	205	0	0	2	0
5	С	238	0	0	6	0
5	D	188	0	0	1	0
5	Е	229	0	0	5	0
5	F	167	0	0	4	0
All	All	15063	0	13296	109	0

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 109 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:24:GLY:HA2	5:F:1496:HOH:O	1.52	1.07
1:C:115:PHE:O	1:C:119:LEU:HD12	1.74	0.87
1:C:4:THR:O	1:C:7:THR:HG23	1.75	0.86
1:C:229:GLN:OE1	1:C:246:TYR:OH	1.96	0.82
1:E:229:GLN:OE1	1:E:246:TYR:OH	1.98	0.81



There are no symmetry-related clashes.

## 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percer	ntiles
1	А	284/307~(92%)	278~(98%)	6(2%)	0	100	100
1	В	286/307~(93%)	280 (98%)	6 (2%)	0	100	100
1	С	290/307~(94%)	284 (98%)	6 (2%)	0	100	100
1	D	283/307~(92%)	278 (98%)	5 (2%)	0	100	100
1	Ε	284/307~(92%)	281 (99%)	3 (1%)	0	100	100
1	F	282/307~(92%)	279~(99%)	3 (1%)	0	100	100
All	All	1709/1842~(93%)	1680 (98%)	29 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the side chain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	А	237/275~(86%)	236 (100%)	1 (0%)	91	96
1	В	231/275~(84%)	230 (100%)	1 (0%)	91	96
1	С	239/275~(87%)	236~(99%)	3 (1%)	69	81
1	D	232/275~(84%)	231 (100%)	1 (0%)	91	96
1	Е	237/275~(86%)	237 (100%)	0	100	100
1	F	232/275~(84%)	232 (100%)	0	100	100



Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1408/1650~(85%)	1402 (100%)	6~(0%)	91 96

5 of 6 residues with a non-rotameric side chain are listed below:

Mol	Chain	Res	Type
1	С	119	LEU
1	В	156	PHE
1	D	156	PHE
1	С	6	GLU
1	А	147	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	21	GLN
1	F	249	HIS
1	Е	262	ASN
1	D	150	GLN
1	D	262	ASN

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

# 5.6 Ligand geometry (i)

Of 25 ligands modelled in this entry, 17 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The



Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond length (or angles).

Mol	Type Chain Res Link		Link	Bo	ond leng	ths	Bond angles			
MOI	туре	Ullalli	nes	LIIIK	Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
3	BFQ	В	1304	2	18,18,18	1.29	2 (11%)	24,27,27	1.15	2 (8%)
3	BFQ	F	1304	2	18,18,18	1.33	2 (11%)	24,27,27	1.09	1 (4%)
3	BFQ	D	1304	2	18,18,18	1.40	5 (27%)	24,27,27	1.30	2 (8%)
3	BFQ	С	1303	1,2	18,18,18	1.79	5 (27%)	24,27,27	1.23	3 (12%)
4	GOL	А	1305	-	$5,\!5,\!5$	0.98	0	$5,\!5,\!5$	1.44	0
4	GOL	С	1304	-	$5,\!5,\!5$	0.85	0	$5,\!5,\!5$	1.16	0
3	BFQ	А	1304	2	18,18,18	1.41	3 (16%)	$24,\!27,\!27$	1.00	0
3	BFQ	Е	1304	2	18,18,18	1.29	4 (22%)	$24,\!27,\!27$	1.58	7 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BFQ	В	1304	2	-	6/29/29/29	-
3	BFQ	F	1304	2	-	9/29/29/29	-
3	BFQ	D	1304	2	-	5/29/29/29	-
3	BFQ	С	1303	1,2	-	9/29/29/29	-
4	GOL	А	1305	-	-	4/4/4/4	-
4	GOL	С	1304	-	-	4/4/4/4	-
3	BFQ	А	1304	2	-	13/29/29/29	-
3	BFQ	Е	1304	2	-	11/29/29/29	-

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	С	1303	BFQ	P1-C7	5.02	1.88	1.85
3	F	1304	BFQ	O14-C7	-3.23	1.40	1.44
3	В	1304	BFQ	O14-C7	-3.23	1.40	1.44
3	Е	1304	BFQ	O14-C7	-3.19	1.40	1.44
3	D	1304	BFQ	O14-C7	-3.12	1.40	1.44

The worst 5 of 15 bond angle outliers are listed below:



6R4V
------

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	Ε	1304	BFQ	P8-C7-P1	-3.25	107.00	112.81
3	D	1304	BFQ	P8-C7-P1	-3.19	107.11	112.81
3	С	1303	BFQ	P8-C7-P1	-3.10	107.27	112.81
3	Е	1304	BFQ	O10-P8-O9	-2.91	106.53	113.06
3	В	1304	BFQ	P8-C7-P1	-2.88	107.66	112.81

There are no chirality outliers.

5 of 61 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	А	1304	BFQ	C19-C16-C7-P1
3	А	1304	BFQ	C19-C16-C7-P8
3	А	1304	BFQ	C19-C16-C7-O14
3	А	1304	BFQ	C16-C19-N22-C23
3	С	1303	BFQ	C19-C16-C7-P1

There are no ring outliers.


7 monomers are involved in 14 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	В	1304	BFQ	2	0
3	F	1304	BFQ	3	0
3	D	1304	BFQ	3	0
3	С	1303	BFQ	1	0
4	С	1304	GOL	1	0
3	А	1304	BFQ	2	0
3	Е	1304	BFQ	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and sufficient the outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.










































## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

# 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle RSRZ \rangle$	#RSI	RZ>2	$OWAB(A^2)$	$\mathbf{Q}{<}0.9$
1	А	288/307~(93%)	-0.44	2(0%)	87 86	23, 34, 55, 102	0
1	В	290/307~(94%)	-0.46	3 (1%)	82 81	20, 36, 56, 85	0
1	С	294/307~(95%)	-0.36	1 (0%)	94 93	21, 30, 56, 94	0
1	D	287/307~(93%)	-0.46	2(0%)	87 86	19, 34, 54, 95	0
1	Ε	288/307~(93%)	-0.48	1 (0%)	94 93	21, 32, 52, 85	0
1	F	286/307~(93%)	-0.21	3 (1%)	82 81	21, 38, 59, 102	0
All	All	1733/1842~(94%)	-0.40	12 (0%)	87 86	19, 34, 56, 102	0

The worst 5 of 12 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	23	PRO	8.0
1	F	24	GLY	4.8
1	Е	24	GLY	4.7
1	В	201	ASN	3.8
1	D	194	HIS	2.5

# 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

# 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.




### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B$ -factors( $Å^2$ )	Q<0.9
2	MG	F	1302	1/1	0.60	0.13	69,69,69,69	0
2	MG	Е	1302	1/1	0.64	0.21	58,58,58,58	0
2	MG	А	1301	1/1	0.71	0.33	45,45,45,45	0
3	BFQ	А	1304	19/19	0.73	0.33	31,62,82,96	0
3	BFQ	С	1303	19/19	0.82	0.28	24,61,75,76	0
2	MG	А	1302	1/1	0.83	0.27	67,67,67,67	0
4	GOL	А	1305	6/6	0.86	0.25	31,50,50,58	0
4	GOL	С	1304	6/6	0.90	0.29	20,39,47,67	0
2	MG	С	1301	1/1	0.94	0.16	37,37,37,37	0
2	MG	С	1302	1/1	0.94	0.06	25,25,25,25	0
2	MG	В	1301	1/1	0.95	0.07	27,27,27,27	0
3	BFQ	В	1304	19/19	0.95	0.16	19,40,75,77	0
3	BFQ	Е	1304	19/19	0.96	0.16	31,50,64,66	0
2	MG	В	1302	1/1	0.96	0.07	35,35,35,35	0
2	MG	D	1301	1/1	0.96	0.05	23,23,23,23	0
3	BFQ	F	1304	19/19	0.96	0.18	30,50,74,74	0
2	MG	F	1301	1/1	0.97	0.13	26,26,26,26	0
2	MG	Е	1303	1/1	0.97	0.05	24,24,24,24	0
2	MG	F	1303	1/1	0.98	0.11	30,30,30,30	0
2	MG	D	1302	1/1	0.98	0.04	41,41,41,41	0
2	MG	D	1303	1/1	0.98	0.08	18,18,18,18	0
3	BFQ	D	1304	19/19	0.98	0.16	26,44,65,68	0
2	MG	Е	1301	1/1	0.98	0.07	25,25,25,25	0
2	MG	А	1303	1/1	0.98	0.07	26,26,26,26	0
2	MG	В	1303	1/1	0.99	0.07	23,23,23,23	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.








































































































# 6.5 Other polymers (i)

There are no such residues in this entry.

