

Full wwPDB X-ray Structure Validation Report (i)

Sep 2, 2023 – 10:17 AM EDT

PDB ID : 3R9J

Title: 4.3A resolution structure of a MinD-MinE(I24N) protein complex

Authors: Lovell, S.; Battaile, K.P.; Park, K.-T.; Wu, W.; Holyoak, T.; Lutkenhaus, J.

Deposited on : 2011-03-25

Resolution : 4.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity : 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.35

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 4.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive	Similar resolution
TVIOUTE	$(\# \mathrm{Entries})$	$(\# ext{Entries}, ext{ resolution range}(ext{Å}))$
R_{free}	130704	1014 (4.80-3.80)
Clashscore	141614	1077 (4.80-3.80)
Ramachandran outliers	138981	1029 (4.80-3.80)
Sidechain outliers	138945	1012 (4.80-3.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain					
1	A	260		88%	8% •			
1	В	260	-	87%	8% • •			
2	С	77	39%	39%	12% • 9%			
2	D	77	47%	36%	8% 9%			



2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 4973 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Septum site-determining protein minD.

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	A	250	Total 1893	C 1189	7.	O 372	S 7	0	0	0
1	В	250		C 1191		O 369	S 7	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	40	ALA	ASP	engineered mutation	UNP P0AEZ3
В	40	ALA	ASP	engineered mutation	UNP P0AEZ3

• Molecule 2 is a protein called Cell division topological specificity factor.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	С	70	Total	С	N	О	S	0	0	0
	70	566	354	100	110	2	0	0		
9	D	70	Total	С	N	О	S	0	0	0
	D	10	566	354	100	110	2	0	0	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	24	ASN	ILE	engineered mutation	UNP P0A734
D	24	ASN	ILE	engineered mutation	UNP P0A734

• Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).






Mol	Chain	Residues	Atoms				ZeroOcc	AltConf		
9	Λ	1	Total	С	N	О	Р	0	0	
$\begin{array}{ c c c c c } \hline 3 & \mathbf{A} & \\ \hline \end{array}$	1	27	10	5	10	2	U	0		
9	D	1	Total	С	N	О	Р	0	0	
3	3 B	1	27	10	5	10	2	U	U	



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Septum site-determining protein minD





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants	94.39Å 94.39Å 284.98Å	Donositor
a, b, c, α , β , γ	90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.71 - 4.30	Depositor
rtesolution (A)	48.71 - 4.30	EDS
% Data completeness	(Not available) (48.71-4.30)	Depositor
(in resolution range)	100.0 (48.71-4.30)	EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	5.23 (at 4.29Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.8.0	Depositor
D D.	0.294 , 0.311	Depositor
R, R_{free}	0.310 , 0.320	DCC
R_{free} test set	455 reflections $(4.85%)$	wwPDB-VP
Wilson B-factor (Å ²)	167.7	Xtriage
Anisotropy	0.496	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.29 , 97.6	EDS
L-test for twinning ²	$ < L > = 0.45, < L^2> = 0.28$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	4973	wwPDB-VP
Average B, all atoms (Å ²)	134.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.61% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bo	nd angles
MIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.38	0/1912	0.55	0/2586
1	В	0.39	0/1913	0.56	0/2586
2	С	0.50	0/572	0.75	0/775
2	D	0.51	0/572	0.79	2/775~(0.3%)
All	All	0.42	0/4969	0.61	$2/6722 \ (0.0\%)$

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
2	D	59	GLU	N-CA-CB	-5.69	100.36	110.60
2	D	58	PRO	CB-CA-C	-5.06	99.36	112.00

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1893	0	1924	12	0
1	В	1894	0	1938	15	0
2	С	566	0	582	48	0
2	D	566	0	582	38	0
3	A	27	0	12	0	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	В	27	0	12	0	0
All	All	4973	0	5050	95	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (95) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic	Clash
Atom-1	Atom-2	${f distance}({ m \AA})$	overlap (Å)
2:D:57:ASP:O	2:D:59:GLU:HB3	1.53	1.06
2:C:57:ASP:O	2:C:59:GLU:HB3	1.58	1.03
2:D:59:GLU:O	2:D:59:GLU:CD	2.08	0.92
2:D:59:GLU:O	2:D:59:GLU:OE1	1.89	0.91
2:C:45:ASP:O	2:C:49:VAL:HG23	1.73	0.89
2:D:45:ASP:O	2:D:49:VAL:HG23	1.74	0.87
2:C:58:PRO:HA	2:C:59:GLU:C	1.97	0.83
2:C:65:LEU:HD12	2:C:75:LEU:HD22	1.61	0.81
2:C:60:MET:O	2:C:60:MET:HG2	1.78	0.81
2:C:57:ASP:O	2:C:59:GLU:CB	2.29	0.79
2:D:78:ASN:OD1	2:D:80:THR:HG23	1.84	0.76
2:C:49:VAL:O	2:C:51:CYS:N	2.20	0.75
2:C:81:LEU:CD1	2:D:75:LEU:HD12	2.18	0.74
2:D:49:VAL:O	2:D:51:CYS:N	2.21	0.73
2:C:65:LEU:HD12	2:C:75:LEU:CD2	2.19	0.72
2:C:45:ASP:O	2:C:49:VAL:CG2	2.37	0.72
1:A:92:ARG:HG2	1:B:125:ILE:HD12	1.71	0.71
2:D:45:ASP:O	2:D:49:VAL:CG2	2.38	0.71
2:C:60:MET:O	2:C:60:MET:CG	2.40	0.69
2:C:58:PRO:HA	2:C:59:GLU:O	1.93	0.68
2:C:81:LEU:HD12	2:D:75:LEU:HD12	1.77	0.66
2:C:81:LEU:CD1	2:D:75:LEU:CD1	2.73	0.66
2:C:49:VAL:C	2:C:51:CYS:H	1.98	0.66
1:B:44:ARG:HD2	1:B:88:ALA:O	1.96	0.65
2:D:49:VAL:C	2:D:51:CYS:H	1.98	0.65
1:B:95:ASP:O	1:B:97:LEU:N	2.30	0.64
2:C:51:CYS:O	2:C:55:GLN:HG3	1.96	0.64
1:A:44:ARG:HD2	1:A:88:ALA:O	1.97	0.64
1:A:151:ARG:HD2	1:B:48:LEU:HD22	1.80	0.64
2:C:58:PRO:HA	2:C:59:GLU:HB3	1.81	0.62
2:C:81:LEU:HD13	2:D:75:LEU:CD1	2.29	0.62
2:C:59:GLU:O	2:C:60:MET:C	2.38	0.62



 $Continued\ from\ previous\ page...$

Continuea from pre		Interatomic	Clash	
Atom-1	Atom-2	${\rm distance}(\mathring{\rm A})$	overlap (Å)	
2:D:25:ILE:O	2:D:28:GLU:HB3	2.00	0.61	
2:D:57:ASP:O	2:D:59:GLU:CB	2.39	0.61	
2:C:48:GLU:O	2:C:51:CYS:HB2	2.00	0.61	
2:D:58:PRO:CA	2:D:59:GLU:HB3	2.31	0.61	
2:D:48:GLU:O	2:D:51:CYS:HB2	2.01	0.60	
2:C:58:PRO:CA	2:C:59:GLU:C	2.69	0.59	
1:B:53:GLU:HG3	2:C:25:ILE:HD11	1.85	0.58	
1:B:94:LYS:O	1:B:127:THR:HG21	2.04	0.57	
1:A:177:HIS:HB3	1:A:208:VAL:HB	1.88	0.55	
1:B:177:HIS:HB3	1:B:208:VAL:HB	1.89	0.55	
2:C:16:ASN:O	2:C:20:GLU:HG3	2.08	0.54	
2:C:74:ILE:HG12	2:D:80:THR:HG22	1.91	0.53	
2:D:58:PRO:HA	2:D:59:GLU:HB3	1.90	0.53	
2:D:58:PRO:HA	2:D:59:GLU:C	2.29	0.53	
2:D:57:ASP:C	2:D:59:GLU:HB3	2.28	0.53	
2:C:65:LEU:HD21	2:C:67:GLN:HE21	1.74	0.52	
2:D:16:ASN:O	2:D:20:GLU:HG3	2.09	0.52	
1:B:220:ALA:HB1	1:B:225:GLU:O	2.10	0.52	
2:C:63:VAL:HG13	2:C:77:LEU:HG	1.91	0.52	
2:C:59:GLU:O	2:C:61:VAL:N	2.44	0.51	
2:C:58:PRO:CA	2:C:59:GLU:HB3	2.41	0.50	
2:C:58:PRO:HA	2:C:59:GLU:CB	2.37	0.50	
1:A:220:ALA:HB1	1:A:225:GLU:O	2.11	0.50	
1:A:201:GLU:OE2	2:C:23:GLN:NE2	2.45	0.49	
2:D:49:VAL:C	2:D:51:CYS:N	2.65	0.49	
2:C:49:VAL:C	2:C:51:CYS:N	2.65	0.49	
2:C:63:VAL:HG22	2:C:77:LEU:CD2	2.42	0.49	
2:C:58:PRO:HG3	2:C:61:VAL:HG23	1.93	0.49	
2:C:79:VAL:HG23	2:D:75:LEU:O	2.13	0.49	
2:C:50:ILE:HD11	2:D:42:LEU:HB3	1.94	0.48	
1:B:144:ASN:ND2	1:B:181:THR:HG23	2.29	0.48	
1:A:144:ASN:ND2	1:A:181:THR:HG23	2.29	0.47	
1:B:14:VAL:O	1:B:181:THR:HG21	2.13	0.47	
1:B:81:GLU:O	1:B:82:ASN:HB2	2.14	0.47	
2:D:25:ILE:HA	2:D:28:GLU:HB3	1.97	0.47	
2:D:65:LEU:HD12	2:D:75:LEU:HD21	1.95	0.47	
2:D:68:LYS:HE2	2:D:74:ILE:HD11	1.95	0.47	
1:A:14:VAL:O	1:A:181:THR:HG21	2.14	0.47	
1:B:57:VAL:O	2:C:44:LYS:NZ	2.39	0.47	
2:C:80:THR:HG22	2:D:74:ILE:HG12	1.97	0.46	
2:C:40:PRO:HA	2:C:43:ARG:HG2	1.98	0.45	



Continued from previous page...

A 4 a sa 1	A4 a 2	Interatomic	Clash
Atom-1	Atom-2	${\rm distance}({\rm \AA})$	overlap (Å)
2:D:43:ARG:HG3	2:D:44:LYS:N	2.31	0.45
2:C:43:ARG:HG3	2:C:44:LYS:N	2.31	0.45
1:A:81:GLU:O	1:A:82:ASN:HB2	2.16	0.45
2:C:76:GLU:OE1	2:D:78:ASN:ND2	2.50	0.45
2:D:40:PRO:HA	2:D:43:ARG:HG2	1.98	0.45
2:C:53:TYR:HB3	2:D:42:LEU:HD13	1.97	0.45
2:C:75:LEU:O	2:D:79:VAL:HG23	2.17	0.45
2:D:62:THR:HB	2:D:78:ASN:HB3	1.99	0.44
2:C:63:VAL:HG22	2:C:77:LEU:HD23	1.99	0.44
2:C:25:ILE:O	2:C:28:GLU:HB2	2.17	0.44
1:A:53:GLU:HG2	1:A:54:ARG:HG3	1.99	0.43
2:C:61:VAL:O	2:C:62:THR:O	2.35	0.43
1:B:53:GLU:HG2	1:B:54:ARG:HG3	2.01	0.42
2:D:36:PRO:HG2	2:D:39:LEU:HB2	2.01	0.42
2:D:25:ILE:O	2:D:29:ARG:N	2.47	0.42
1:A:147:VAL:CG1	1:B:48:LEU:HG	2.49	0.42
1:A:48:LEU:HD22	1:B:151:ARG:HD2	2.02	0.42
2:D:65:LEU:HD12	2:D:75:LEU:CD2	2.51	0.41
2:C:68:LYS:HE3	2:C:69:ASP:H	1.86	0.41
2:C:26:VAL:O	2:C:30:ARG:HG3	2.22	0.40
2:D:78:ASN:OD1	2:D:80:THR:CG2	2.62	0.40
2:C:36:PRO:O	2:C:40:PRO:CD	2.70	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed Favoured Allowed		Outliers	Perc	entiles	
1	A	$246/260 \ (95\%)$	239 (97%)	4 (2%)	3 (1%)	13	50
1	В	$246/260 \ (95\%)$	238 (97%)	4 (2%)	4 (2%)	9	45
2	С	68/77 (88%)	56 (82%)	9 (13%)	3 (4%)	2	25



Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perc	entiles
2	D	68/77 (88%)	57 (84%)	8 (12%)	3 (4%)	2	25
All	All	628/674 (93%)	590 (94%)	25 (4%)	13 (2%)	7	39

All (13) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	95	ASP
2	С	50	ILE
2	D	50	ILE
2	D	60	MET
1	A	9	SER
1	A	96	ALA
1	A	126	GLU
1	В	9	SER
1	В	96	ALA
1	В	126	GLU
2	С	55	GLN
2	С	60	MET
2	D	55	GLN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percer	ntiles
1	A	203/216 (94%)	202 (100%)	1 (0%)	88	93
1	В	204/216 (94%)	201 (98%)	3 (2%)	65	80
2	С	$65/71 \; (92\%)$	53 (82%)	12 (18%)	1	11
2	D	$65/71 \ (92\%)$	56 (86%)	9 (14%)	3	20
All	All	537/574 (94%)	512 (95%)	25 (5%)	26	53

All (25) residues with a non-rotameric sidechain are listed below:

		<i>0</i> 1
1 A	60	PHE



Continued from previous page...

Mol	Chain	Res	Type
1	В	60	PHE
1	В	94	LYS
1	В	97	LEU
2 2	С	22	LEU
	С	37	HIS
2	C C C C C C C D	39	LEU
2	С	46	ILE
2	С	53	TYR
2 2 2 2	С	57	ASP
2	С	60	MET
2	С	62	THR
2	С	65	LEU
2	С	68	LYS
2	С	76	GLU
2	С	81	LEU
2	D	37	HIS
2	D	39	LEU
2	D	46	ILE
2	D D	53	TYR
2	D	57	ASP
2	D	60	MET
2	D	62	THR
2	D	77	LEU
2	D	81	LEU

Sometimes side chains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such side chains are listed below:

Mol	Chain	Res	Type
1	A	45	ASN
1	A	144	ASN
1	A	177	HIS
1	A	215	GLN
1	В	45	ASN
1	В	144	ASN
1	В	215	GLN
2	С	67	GLN
2	С	78	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Mol Type Chain Res		Link	Bond lengths			В	ond ang	cles	
IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	A	261	-	24,29,29	1.15	2 (8%)	29,45,45	0.93	0
3	ADP	В	261	-	24,29,29	0.97	2 (8%)	29,45,45	1.03	2 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	261	-	-	5/12/32/32	0/3/3/3
3	ADP	В	261	-	-	3/12/32/32	0/3/3/3

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	Ideal(Å)
3	A	261	ADP	O4'-C1'	3.58	1.46	1.41
3	A	261	ADP	C2-N3	3.30	1.37	1.32
3	В	261	ADP	C2-N3	3.00	1.36	1.32
3	В	261	ADP	O4'-C1'	2.05	1.43	1.41

All (2) bond angle outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	В	261	ADP	C5-C6-N6	2.24	123.76	120.35
3	В	261	ADP	O2B-PB-O3A	2.13	111.76	104.64

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	261	ADP	PA-O3A-PB-O2B
3	В	261	ADP	C5'-O5'-PA-O1A
3	В	261	ADP	PA-O3A-PB-O1B
3	A	261	ADP	C5'-O5'-PA-O3A
3	В	261	ADP	C5'-O5'-PA-O3A
3	A	261	ADP	PA-O3A-PB-O1B
3	A	261	ADP	PA-O3A-PB-O3B
3	A	261	ADP	C5'-O5'-PA-O1A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.


6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

