



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 8, 2023 – 12:28 PM EDT

PDB ID : 4R9L
Title : Structure of a thermostable elevenfold mutant of limonene epoxide hydrolase from *Rhodococcus erythropolis*, containing two stabilizing disulfide bonds
Authors : Floor, R.J.; Wijma, H.J.; Jekel, P.A.; Terwisscha van Scheltinga, A.C.; Dijkstra, B.W.; Janssen, D.B.
Deposited on : 2014-09-05
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5950 (1.80-1.80)
Clashscore	141614	6793 (1.80-1.80)
Ramachandran outliers	138981	6697 (1.80-1.80)
Sidechain outliers	138945	6696 (1.80-1.80)
RSRZ outliers	127900	5850 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	174	 2% 80% 16%
1	B	174	 3% 83% 5% 12%
1	C	174	 8% 83% 8% 8%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 3854 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Limonene-1,2-epoxide hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	147	1162	744	189	219	10	0	3	0
1	B	153	1202	768	196	228	10	0	2	0
1	C	160	1254	800	204	242	8	0	0	0

There are 114 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	expression tag	UNP Q9ZAG3
A	2	ALA	-	expression tag	UNP Q9ZAG3
A	5	CYS	ILE	engineered mutation	UNP Q9ZAG3
A	15	PRO	SER	engineered mutation	UNP Q9ZAG3
A	19	LYS	ALA	engineered mutation	UNP Q9ZAG3
A	76	LYS	THR	engineered mutation	UNP Q9ZAG3
A	84	CYS	GLU	engineered mutation	UNP Q9ZAG3
A	85	VAL	THR	engineered mutation	UNP Q9ZAG3
A	89	CYS	GLY	engineered mutation	UNP Q9ZAG3
A	91	CYS	SER	engineered mutation	UNP Q9ZAG3
A	92	LYS	ASN	engineered mutation	UNP Q9ZAG3
A	96	PHE	TYR	engineered mutation	UNP Q9ZAG3
A	124	ASP	GLU	engineered mutation	UNP Q9ZAG3
A	150	LYS	-	expression tag	UNP Q9ZAG3
A	151	LEU	-	expression tag	UNP Q9ZAG3
A	152	GLY	-	expression tag	UNP Q9ZAG3
A	153	PRO	-	expression tag	UNP Q9ZAG3
A	154	GLU	-	expression tag	UNP Q9ZAG3
A	155	GLN	-	expression tag	UNP Q9ZAG3
A	156	LYS	-	expression tag	UNP Q9ZAG3
A	157	LEU	-	expression tag	UNP Q9ZAG3
A	158	ILE	-	expression tag	UNP Q9ZAG3
A	159	SER	-	expression tag	UNP Q9ZAG3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	160	GLU	-	expression tag	UNP Q9ZAG3
A	161	GLU	-	expression tag	UNP Q9ZAG3
A	162	ASP	-	expression tag	UNP Q9ZAG3
A	163	LEU	-	expression tag	UNP Q9ZAG3
A	164	ASN	-	expression tag	UNP Q9ZAG3
A	165	SER	-	expression tag	UNP Q9ZAG3
A	166	ALA	-	expression tag	UNP Q9ZAG3
A	167	VAL	-	expression tag	UNP Q9ZAG3
A	168	ASP	-	expression tag	UNP Q9ZAG3
A	169	HIS	-	expression tag	UNP Q9ZAG3
A	170	HIS	-	expression tag	UNP Q9ZAG3
A	171	HIS	-	expression tag	UNP Q9ZAG3
A	172	HIS	-	expression tag	UNP Q9ZAG3
A	173	HIS	-	expression tag	UNP Q9ZAG3
A	174	HIS	-	expression tag	UNP Q9ZAG3
B	1	MET	-	expression tag	UNP Q9ZAG3
B	2	ALA	-	expression tag	UNP Q9ZAG3
B	5	CYS	ILE	engineered mutation	UNP Q9ZAG3
B	15	PRO	SER	engineered mutation	UNP Q9ZAG3
B	19	LYS	ALA	engineered mutation	UNP Q9ZAG3
B	76	LYS	THR	engineered mutation	UNP Q9ZAG3
B	84	CYS	GLU	engineered mutation	UNP Q9ZAG3
B	85	VAL	THR	engineered mutation	UNP Q9ZAG3
B	89	CYS	GLY	engineered mutation	UNP Q9ZAG3
B	91	CYS	SER	engineered mutation	UNP Q9ZAG3
B	92	LYS	ASN	engineered mutation	UNP Q9ZAG3
B	96	PHE	TYR	engineered mutation	UNP Q9ZAG3
B	124	ASP	GLU	engineered mutation	UNP Q9ZAG3
B	150	LYS	-	expression tag	UNP Q9ZAG3
B	151	LEU	-	expression tag	UNP Q9ZAG3
B	152	GLY	-	expression tag	UNP Q9ZAG3
B	153	PRO	-	expression tag	UNP Q9ZAG3
B	154	GLU	-	expression tag	UNP Q9ZAG3
B	155	GLN	-	expression tag	UNP Q9ZAG3
B	156	LYS	-	expression tag	UNP Q9ZAG3
B	157	LEU	-	expression tag	UNP Q9ZAG3
B	158	ILE	-	expression tag	UNP Q9ZAG3
B	159	SER	-	expression tag	UNP Q9ZAG3
B	160	GLU	-	expression tag	UNP Q9ZAG3
B	161	GLU	-	expression tag	UNP Q9ZAG3
B	162	ASP	-	expression tag	UNP Q9ZAG3
B	163	LEU	-	expression tag	UNP Q9ZAG3

Continued on next page...

Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
B	164	ASN	-	expression tag	UNP Q9ZAG3
B	165	SER	-	expression tag	UNP Q9ZAG3
B	166	ALA	-	expression tag	UNP Q9ZAG3
B	167	VAL	-	expression tag	UNP Q9ZAG3
B	168	ASP	-	expression tag	UNP Q9ZAG3
B	169	HIS	-	expression tag	UNP Q9ZAG3
B	170	HIS	-	expression tag	UNP Q9ZAG3
B	171	HIS	-	expression tag	UNP Q9ZAG3
B	172	HIS	-	expression tag	UNP Q9ZAG3
B	173	HIS	-	expression tag	UNP Q9ZAG3
B	174	HIS	-	expression tag	UNP Q9ZAG3
C	1	MET	-	expression tag	UNP Q9ZAG3
C	2	ALA	-	expression tag	UNP Q9ZAG3
C	5	CYS	ILE	engineered mutation	UNP Q9ZAG3
C	15	PRO	SER	engineered mutation	UNP Q9ZAG3
C	19	LYS	ALA	engineered mutation	UNP Q9ZAG3
C	76	LYS	THR	engineered mutation	UNP Q9ZAG3
C	84	CYS	GLU	engineered mutation	UNP Q9ZAG3
C	85	VAL	THR	engineered mutation	UNP Q9ZAG3
C	89	CYS	GLY	engineered mutation	UNP Q9ZAG3
C	91	CYS	SER	engineered mutation	UNP Q9ZAG3
C	92	LYS	ASN	engineered mutation	UNP Q9ZAG3
C	96	PHE	TYR	engineered mutation	UNP Q9ZAG3
C	124	ASP	GLU	engineered mutation	UNP Q9ZAG3
C	150	LYS	-	expression tag	UNP Q9ZAG3
C	151	LEU	-	expression tag	UNP Q9ZAG3
C	152	GLY	-	expression tag	UNP Q9ZAG3
C	153	PRO	-	expression tag	UNP Q9ZAG3
C	154	GLU	-	expression tag	UNP Q9ZAG3
C	155	GLN	-	expression tag	UNP Q9ZAG3
C	156	LYS	-	expression tag	UNP Q9ZAG3
C	157	LEU	-	expression tag	UNP Q9ZAG3
C	158	ILE	-	expression tag	UNP Q9ZAG3
C	159	SER	-	expression tag	UNP Q9ZAG3
C	160	GLU	-	expression tag	UNP Q9ZAG3
C	161	GLU	-	expression tag	UNP Q9ZAG3
C	162	ASP	-	expression tag	UNP Q9ZAG3
C	163	LEU	-	expression tag	UNP Q9ZAG3
C	164	ASN	-	expression tag	UNP Q9ZAG3
C	165	SER	-	expression tag	UNP Q9ZAG3
C	166	ALA	-	expression tag	UNP Q9ZAG3
C	167	VAL	-	expression tag	UNP Q9ZAG3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	168	ASP	-	expression tag	UNP Q9ZAG3
C	169	HIS	-	expression tag	UNP Q9ZAG3
C	170	HIS	-	expression tag	UNP Q9ZAG3
C	171	HIS	-	expression tag	UNP Q9ZAG3
C	172	HIS	-	expression tag	UNP Q9ZAG3
C	173	HIS	-	expression tag	UNP Q9ZAG3
C	174	HIS	-	expression tag	UNP Q9ZAG3

- Molecule 2 is (2R)-2-hydroxyhexanamide (three-letter code: HYH) (formula: C₆H₁₃NO₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
2	A	1	9	6	1	2	0	0
2	B	1	9	6	1	2	0	0
2	C	1	9	6	1	2	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	54	54	54	0	0
3	B	66	66	66	0	0
3	C	89	89	89	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Limonene-1,2-epoxide hydrolase



- Molecule 1: Limonene-1,2-epoxide hydrolase



- Molecule 1: Limonene-1,2-epoxide hydrolase



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	87.69Å 87.69Å 108.91Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	44.25 – 1.80 44.25 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.4 (44.25-1.80) 99.4 (44.25-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 1.79Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.160 , 0.207 0.173 , 0.214	Depositor DCC
R_{free} test set	2280 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	23.1	Xtrriage
Anisotropy	0.023	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 43.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	3854	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HYH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.00	0/1196	1.03	2/1619 (0.1%)
1	B	1.00	0/1234	1.01	5/1671 (0.3%)
1	C	1.07	1/1280 (0.1%)	1.00	3/1734 (0.2%)
All	All	1.02	1/3710 (0.0%)	1.01	10/5024 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	49	GLU	CD-OE1	5.27	1.31	1.25

The worst 5 of 10 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	ARG	NE-CZ-NH1	9.33	124.97	120.30
1	C	91	CYS	CB-CA-C	-6.86	96.69	110.40
1	C	24	ASP	CB-CG-OD1	6.45	124.10	118.30
1	A	64	ARG	NE-CZ-NH2	-5.91	117.34	120.30
1	B	91[A]	CYS	CB-CA-C	-5.88	98.64	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1162	0	1152	5	0
1	B	1202	0	1187	9	0
1	C	1254	0	1236	7	0
2	A	9	0	13	2	0
2	B	9	0	13	2	0
2	C	9	0	13	0	0
3	A	54	0	0	2	0
3	B	66	0	0	3	0
3	C	89	0	0	0	0
All	All	3854	0	3614	22	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 22 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:201:HYH:H3	3:A:353:HOH:O	1.67	0.94
1:A:6:GLU:O	3:A:347:HOH:O	2.09	0.69
1:B:155:GLN:OE1	3:B:363:HOH:O	2.10	0.69
1:C:135:ASP:OD2	1:C:138:GLU:HG3	2.03	0.58
1:C:123:THR:O	1:C:123:THR:HG23	2.04	0.57

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	148/174 (85%)	145 (98%)	2 (1%)	1 (1%)	22	10
1	B	153/174 (88%)	147 (96%)	6 (4%)	0	100	100
1	C	158/174 (91%)	157 (99%)	0	1 (1%)	25	12

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	459/522 (88%)	449 (98%)	8 (2%)	2 (0%)	34 21

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	92	LYS
1	C	92	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	127/148 (86%)	127 (100%)	0	100 100
1	B	131/148 (88%)	131 (100%)	0	100 100
1	C	136/148 (92%)	135 (99%)	1 (1%)	84 81
All	All	394/444 (89%)	393 (100%)	1 (0%)	92 91

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	64	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	HYH	C	201	-	8,8,8	1.35	1 (12%)	7,9,9	3.69	3 (42%)
2	HYH	A	201	-	8,8,8	0.64	0	7,9,9	2.21	2 (28%)
2	HYH	B	201	-	8,8,8	1.38	1 (12%)	7,9,9	3.16	3 (42%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HYH	C	201	-	-	3/8/8/8	-
2	HYH	A	201	-	-	1/8/8/8	-
2	HYH	B	201	-	-	3/8/8/8	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	HYH	C4-C5	-3.40	1.49	1.52
2	C	201	HYH	C4-C5	-2.30	1.50	1.52

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	201	HYH	C4-C5-N	8.13	123.89	117.25
2	B	201	HYH	C4-C5-N	6.88	122.87	117.25

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	201	HYH	O1-C5-C4	4.53	123.10	118.94
2	C	201	HYH	O1-C5-C4	-4.13	115.16	118.94
2	B	201	HYH	O-C4-C5	-3.24	103.67	111.10

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	HYH	O-C4-C5-N
2	B	201	HYH	O-C4-C5-N
2	C	201	HYH	O-C4-C5-N
2	B	201	HYH	C3-C4-C5-N
2	B	201	HYH	C3-C4-C5-O1

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	201	HYH	2	0
2	B	201	HYH	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	147/174 (84%)	-0.50	3 (2%) 65 61	16, 26, 52, 79	0
1	B	153/174 (87%)	-0.46	5 (3%) 46 40	16, 25, 64, 87	0
1	C	160/174 (91%)	-0.48	1 (0%) 89 87	13, 22, 42, 72	0
All	All	460/522 (88%)	-0.48	9 (1%) 65 61	13, 25, 52, 87	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	3	SER	5.5
1	B	5	CYS	3.2
1	A	6	GLU	3.0
1	C	164	ASN	2.9
1	B	155	GLN	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
2	HYH	A	201	9/9	0.88	0.12	38,40,47,48	0
2	HYH	C	201	9/9	0.91	0.12	26,30,37,49	0
2	HYH	B	201	9/9	0.94	0.10	33,38,42,52	0

6.5 Other polymers [i](#)

There are no such residues in this entry.