



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 8, 2020 – 10:13 PM BST

PDB ID : 6RB3
Title : Structural basis for recognition and ring-cleavage of the Pseudomonas quinolone signal (PQS) by AqdC variant in complex with its substrate
Authors : Wullich, S.; Kobus, S.; Smits, S.H.; Fetzner, S.
Deposited on : 2019-04-09
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.13.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	5042 (2.30-2.30)
Clashscore	141614	5643 (2.30-2.30)
Ramachandran outliers	138981	5575 (2.30-2.30)
Sidechain outliers	138945	5575 (2.30-2.30)
RSRZ outliers	127900	4938 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	269	
1	B	269	
1	E	269	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	JWW	A	301	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 12739 atoms, of which 6012 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Putative dioxygenase (1H-3-hydroxy-4-oxoquinoline 2,4-dioxygenase).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	E	267	4058	1312	1984	377	376	9	0	0	0
1	A	267	4058	1312	1984	377	376	9	0	0	0
1	B	267	4058	1312	1984	377	376	9	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	94	SER	ALA	engineered mutation	UNP B1MFK2
E	244	ALA	HIS	engineered mutation	UNP B1MFK2
A	94	SER	ALA	engineered mutation	UNP B1MFK2
A	244	ALA	HIS	engineered mutation	UNP B1MFK2
B	94	SER	ALA	engineered mutation	UNP B1MFK2
B	244	ALA	HIS	engineered mutation	UNP B1MFK2

- Molecule 2 is 2-heptylquinoline-3,4-diol (three-letter code: JWW) (formula: C₁₆H₂₁NO₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
2	E	1	Total	C	H	N	O	0	0
			39	16	20	1	2		
2	A	1	Total	C	H	N	O	0	0
			39	16	20	1	2		
2	B	1	Total	C	H	N	O	0	0
			39	16	20	1	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	134	Total	O	0	0
			134	134		
3	A	161	Total	O	0	0
			161	161		
3	B	153	Total	O	0	0
			153	153		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Putative dioxygenase (1H-3-hydroxy-4-oxoquinaldine 2,4-dioxygenase)



- Molecule 1: Putative dioxygenase (1H-3-hydroxy-4-oxoquinaldine 2,4-dioxygenase)



- Molecule 1: Putative dioxygenase (1H-3-hydroxy-4-oxoquinaldine 2,4-dioxygenase)



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	96.64Å 96.64Å 190.14Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	47.53 – 2.30 47.53 – 2.30	Depositor EDS
% Data completeness (in resolution range)	100.0 (47.53-2.30) 100.0 (47.53-2.30)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.12 (at 2.29Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.195 , 0.246 0.195 , 0.245	Depositor DCC
R_{free} test set	2078 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtrriage
Anisotropy	0.059	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	12739	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: JWW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/2132	0.81	2/2903 (0.1%)
1	B	0.57	0/2132	0.71	2/2903 (0.1%)
1	E	0.60	0/2132	0.75	1/2903 (0.0%)
All	All	0.60	0/6396	0.76	5/8709 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	E	0	1
All	All	0	2

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	32	ASP	CB-CG-OD2	-5.56	113.29	118.30
1	B	146	ARG	NE-CZ-NH2	-5.45	117.57	120.30
1	A	147	ASP	CB-CG-OD2	5.32	123.09	118.30
1	B	213	SER	C-N-CA	-5.31	108.42	121.70
1	E	213	SER	C-N-CA	-5.07	109.02	121.70

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	213	SER	Peptide
1	E	213	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2074	1984	1990	64	0
1	B	2074	1984	1990	49	0
1	E	2074	1984	1990	20	0
2	A	19	20	0	0	0
2	B	19	20	0	1	0
2	E	19	20	0	2	0
3	A	161	0	0	18	0
3	B	153	0	0	7	0
3	E	134	0	0	4	0
All	All	6727	6012	5970	103	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (103) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:-1:MET:N	3:B:401:HOH:O	1.88	1.03
1:A:141:ARG:NH2	3:A:401:HOH:O	1.97	0.98
1:B:230:ARG:NH2	3:B:402:HOH:O	2.01	0.93
1:A:146:ARG:NH2	3:A:404:HOH:O	2.02	0.90
1:E:234:PHE:O	1:A:127:ARG:NH1	2.08	0.86
1:A:151:GLN:N	3:A:403:HOH:O	2.09	0.86
1:A:147:ASP:N	1:A:147:ASP:OD1	2.01	0.84
1:E:265:ALA:O	3:E:401:HOH:O	1.95	0.84
1:A:146:ARG:NH1	3:A:404:HOH:O	2.10	0.82
1:A:147:ASP:OD2	3:A:402:HOH:O	2.02	0.78
1:A:151:GLN:OE1	3:A:403:HOH:O	2.02	0.77
1:A:127:ARG:NH2	3:A:406:HOH:O	2.19	0.75
1:A:146:ARG:CZ	3:A:404:HOH:O	2.29	0.75

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:236:TYR:HA	1:B:127:ARG:HH21	1.52	0.74
1:E:15:GLN:HG3	3:E:435:HOH:O	1.87	0.72
1:B:167:ARG:HD3	3:B:407:HOH:O	1.89	0.71
1:A:237:ARG:NH1	1:B:127:ARG:O	2.25	0.69
1:A:237:ARG:CZ	1:B:127:ARG:HB3	2.24	0.68
1:B:211:VAL:CG1	1:B:239:LEU:HD11	2.24	0.67
1:A:153:TRP:HB3	1:A:245:PHE:HE2	1.60	0.66
1:E:236:TYR:H	1:A:127:ARG:HD3	1.60	0.66
1:A:209:ARG:HD3	1:B:127:ARG:CZ	2.25	0.66
1:A:141:ARG:HA	3:A:494:HOH:O	1.96	0.65
1:B:121:ILE:HD11	3:B:541:HOH:O	1.97	0.64
1:A:150:VAL:O	1:A:153:TRP:N	2.29	0.64
1:B:237:ARG:NH2	3:B:404:HOH:O	2.31	0.63
1:B:211:VAL:HG11	1:B:239:LEU:HD11	1.81	0.62
1:E:214:HIS:O	1:E:215:PRO:C	2.38	0.62
1:A:209:ARG:CG	1:B:127:ARG:CZ	2.78	0.61
1:A:209:ARG:HD3	1:B:127:ARG:NH1	2.16	0.60
1:E:228:ALA:HB1	1:A:127:ARG:NH1	2.15	0.60
1:A:154:LEU:O	1:A:157:THR:HG22	2.03	0.59
1:A:237:ARG:NE	1:A:257:GLU:OE2	2.36	0.58
1:A:236:TYR:O	1:B:127:ARG:HB2	2.05	0.57
1:A:127:ARG:NH1	3:A:406:HOH:O	2.38	0.56
1:E:107:LEU:HD13	1:E:111:ARG:HG2	1.88	0.56
1:E:94:SER:OG	2:E:301:JWW:O18	2.19	0.56
1:B:214:HIS:O	1:B:215:PRO:C	2.45	0.55
1:B:54:TRP:CZ3	1:B:72:MET:HB3	2.41	0.55
1:B:116:MET:HE1	1:B:257:GLU:HB3	1.88	0.55
1:A:127:ARG:CZ	3:A:406:HOH:O	2.54	0.55
1:A:127:ARG:O	3:A:405:HOH:O	2.18	0.54
1:A:154:LEU:O	1:A:155:ALA:HB3	2.08	0.54
1:A:239:LEU:HD22	1:A:253:GLN:OE1	2.08	0.54
1:A:215:PRO:HG2	1:A:220:TYR:CE2	2.43	0.54
1:A:143:LYS:HE3	3:A:404:HOH:O	2.08	0.53
1:E:36:TYR:HA	3:E:460:HOH:O	2.10	0.52
1:A:183:ARG:HG3	3:A:429:HOH:O	2.09	0.52
1:A:209:ARG:HG2	1:B:127:ARG:NH2	2.25	0.52
1:E:137:GLN:OE1	1:E:183:ARG:HG2	2.10	0.51
1:A:210:HIS:O	1:B:127:ARG:NH2	2.44	0.51
1:E:215:PRO:HG2	1:E:220:TYR:CE2	2.46	0.50
1:A:71:GLU:OE1	3:A:407:HOH:O	2.20	0.50
1:A:236:TYR:HA	1:B:127:ARG:NH2	2.24	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:211:VAL:HG22	1:B:127:ARG:HH12	1.76	0.49
1:A:236:TYR:N	1:B:127:ARG:HE	2.10	0.49
1:A:237:ARG:NE	1:B:127:ARG:HB3	2.26	0.49
1:A:222:ALA:HA	1:B:216:LYS:HZ1	1.79	0.48
1:B:94:SER:OG	2:B:301:JWW:O18	2.26	0.48
1:E:235:SER:HA	1:A:127:ARG:NH1	2.28	0.48
1:A:237:ARG:N	1:B:127:ARG:CZ	2.77	0.48
1:E:204:GLU:HG3	1:E:204:GLU:O	2.14	0.48
1:A:150:VAL:HB	3:A:403:HOH:O	2.13	0.47
1:A:209:ARG:CD	1:B:127:ARG:CZ	2.91	0.47
1:A:209:ARG:CG	1:B:127:ARG:NH2	2.78	0.47
1:E:224:HIS:HB3	1:E:236:TYR:CD1	2.49	0.47
1:E:87:SER:HA	1:E:111:ARG:O	2.15	0.47
1:A:204:GLU:HG3	1:A:204:GLU:O	2.14	0.47
1:A:237:ARG:HB2	1:B:127:ARG:NH1	2.30	0.47
1:A:143:LYS:HE3	1:A:146:ARG:HH22	1.80	0.46
1:B:217:ILE:HG12	1:B:238:ARG:HD3	1.96	0.46
1:A:237:ARG:HE	1:A:257:GLU:CD	2.19	0.46
1:B:215:PRO:HG2	1:B:220:TYR:CE2	2.50	0.46
1:E:24:VAL:HA	1:E:50:VAL:O	2.15	0.45
1:A:31:HIS:HB2	3:A:472:HOH:O	2.15	0.45
1:A:237:ARG:CB	1:B:127:ARG:NH1	2.80	0.45
1:B:211:VAL:CG1	1:B:239:LEU:CD1	2.93	0.44
1:B:136:ILE:HA	1:B:142:TRP:HA	2.00	0.44
1:A:236:TYR:C	1:B:127:ARG:NE	2.72	0.44
1:B:204:GLU:O	1:B:204:GLU:HG3	2.17	0.44
1:A:154:LEU:HD21	1:A:166:VAL:HB	2.00	0.43
1:A:149:LEU:O	1:A:150:VAL:C	2.56	0.43
1:B:-1:MET:CB	3:B:401:HOH:O	2.66	0.43
1:A:236:TYR:CA	1:B:127:ARG:HE	2.31	0.42
1:B:87:SER:HA	1:B:111:ARG:O	2.19	0.42
1:B:118:LEU:HB3	1:B:246:PRO:HG2	2.02	0.42
1:A:237:ARG:N	1:B:127:ARG:NH2	2.67	0.42
1:A:209:ARG:HG3	1:B:127:ARG:CZ	2.48	0.42
2:E:301:JWW:C09	3:E:423:HOH:O	2.67	0.42
1:B:217:ILE:HG12	1:B:238:ARG:CD	2.49	0.41
1:E:235:SER:HA	1:A:127:ARG:CZ	2.49	0.41
1:A:209:ARG:NE	1:B:127:ARG:HD3	2.35	0.41
1:B:142:TRP:CH2	1:B:174:GLY:HA2	2.55	0.41
1:B:224:HIS:HB3	1:B:236:TYR:CD1	2.56	0.41
1:A:147:ASP:CG	3:A:402:HOH:O	2.52	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:251:PRO:HD2	1:A:252:GLN:HE22	1.86	0.41
1:E:214:HIS:HE1	1:E:242:GLU:O	2.02	0.41
1:E:252:GLN:H	1:E:252:GLN:CD	2.24	0.40
1:B:119:ASP:HB3	3:B:424:HOH:O	2.21	0.40
1:B:54:TRP:HB2	1:B:57:HIS:ND1	2.37	0.40
1:E:234:PHE:O	1:A:127:ARG:CZ	2.67	0.40
1:A:214:HIS:O	1:A:215:PRO:C	2.59	0.40
1:A:209:ARG:HG3	1:B:127:ARG:NE	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	265/269 (98%)	248 (94%)	14 (5%)	3 (1%)	14	15
1	B	265/269 (98%)	255 (96%)	9 (3%)	1 (0%)	34	42
1	E	265/269 (98%)	252 (95%)	11 (4%)	2 (1%)	19	23
All	All	795/807 (98%)	755 (95%)	34 (4%)	6 (1%)	19	23

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	150	VAL
1	E	86	ASP
1	A	119	ASP
1	A	155	ALA
1	B	86	ASP
1	E	214	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	205/208 (99%)	195 (95%)	10 (5%)	25	35
1	B	205/208 (99%)	199 (97%)	6 (3%)	42	58
1	E	205/208 (99%)	200 (98%)	5 (2%)	49	66
All	All	615/624 (99%)	594 (97%)	21 (3%)	37	51

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	31	HIS
1	E	119	ASP
1	E	128	GLU
1	E	141	ARG
1	E	214	HIS
1	A	31	HIS
1	A	128	GLU
1	A	136	ILE
1	A	140	GLU
1	A	141	ARG
1	A	142	TRP
1	A	147	ASP
1	A	160	GLN
1	A	226	ASP
1	A	252	GLN
1	B	31	HIS
1	B	119	ASP
1	B	141	ARG
1	B	214	HIS
1	B	237	ARG
1	B	252	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	E	252	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	JWW	E	301	-	20,20,20	2.24	5 (25%)	21,26,26	1.91	5 (23%)
2	JWW	B	301	-	20,20,20	2.05	5 (25%)	21,26,26	1.67	4 (19%)
2	JWW	A	301	-	20,20,20	2.07	4 (20%)	21,26,26	1.70	5 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JWW	E	301	-	-	4/7/7/7	0/2/2/2
2	JWW	B	301	-	-	4/7/7/7	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	JWW	A	301	-	-	4/7/7/7	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	301	JWW	C09-N08	7.00	1.41	1.32
2	A	301	JWW	C09-N08	6.63	1.40	1.32
2	B	301	JWW	C09-N08	6.44	1.40	1.32
2	E	301	JWW	C06-C05	-3.58	1.36	1.42
2	A	301	JWW	C06-C05	-3.44	1.36	1.42
2	E	301	JWW	C10-C07	3.26	1.45	1.39
2	B	301	JWW	C10-C07	3.17	1.45	1.39
2	E	301	JWW	C04-C05	-2.96	1.36	1.41
2	B	301	JWW	C04-C05	-2.65	1.37	1.41
2	A	301	JWW	C10-C07	2.58	1.44	1.39
2	B	301	JWW	O19-C07	2.38	1.43	1.35
2	B	301	JWW	C06-C05	-2.15	1.39	1.42
2	E	301	JWW	O19-C07	2.13	1.42	1.35
2	A	301	JWW	C04-C05	-2.07	1.38	1.41

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	301	JWW	C09-N08-C05	4.53	123.25	117.67
2	A	301	JWW	C09-N08-C05	4.37	123.06	117.67
2	E	301	JWW	C04-C05-C06	4.19	123.94	119.13
2	B	301	JWW	C04-C05-C06	3.81	123.50	119.13
2	B	301	JWW	C09-N08-C05	3.59	122.10	117.67
2	B	301	JWW	O19-C07-C06	3.03	123.06	116.64
2	A	301	JWW	C06-C05-N08	-2.91	119.72	122.81
2	A	301	JWW	O19-C07-C06	2.79	122.56	116.64
2	A	301	JWW	C04-C05-C06	2.77	122.31	119.13
2	E	301	JWW	O19-C07-C06	2.73	122.43	116.64
2	E	301	JWW	C06-C05-N08	-2.48	120.18	122.81
2	E	301	JWW	C07-C10-C09	-2.18	119.20	120.56
2	B	301	JWW	C04-C05-N08	-2.18	115.38	118.69
2	A	301	JWW	C07-C10-C09	-2.14	119.23	120.56

There are no chirality outliers.

All (12) torsion outliers are listed below:



Mol	Chain	Res	Type	Atoms
2	A	301	JWW	C12-C13-C14-C15
2	E	301	JWW	C11-C12-C13-C14
2	B	301	JWW	C11-C12-C13-C14
2	E	301	JWW	C12-C13-C14-C15
2	B	301	JWW	C12-C13-C14-C15
2	A	301	JWW	C11-C12-C13-C14
2	A	301	JWW	C14-C15-C16-C17
2	E	301	JWW	C14-C15-C16-C17
2	A	301	JWW	C13-C14-C15-C16
2	B	301	JWW	C14-C15-C16-C17
2	E	301	JWW	C13-C14-C15-C16
2	B	301	JWW	C13-C14-C15-C16


There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	301	JWW	2	0
2	B	301	JWW	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	267/269 (99%)	0.17	20 (7%) 14 19	23, 34, 67, 107	0
1	B	267/269 (99%)	-0.20	3 (1%) 80 85	26, 37, 55, 75	0
1	E	267/269 (99%)	-0.19	3 (1%) 80 85	24, 33, 52, 75	0
All	All	801/807 (99%)	-0.07	26 (3%) 47 54	23, 35, 58, 107	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	SER	10.4
1	A	153	TRP	8.0
1	A	154	LEU	6.4
1	A	214	HIS	6.3
1	A	149	LEU	5.7
1	B	127	ARG	5.3
1	B	17	HIS	4.6
1	E	141	ARG	4.5
1	A	127	ARG	4.5
1	A	141	ARG	4.1
1	A	150	VAL	3.9
1	A	143	LYS	3.8
1	B	203	ALA	3.3
1	A	151	GLN	3.3
1	A	148	GLY	3.1
1	A	140	GLU	3.1
1	A	17	HIS	3.1
1	A	146	ARG	2.9
1	A	156	GLY	2.7
1	E	149	LEU	2.5
1	A	142	TRP	2.4
1	A	216	LYS	2.4
1	E	203	ALA	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	145	GLY	2.1
1	A	31	HIS	2.1
1	A	237	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	JWW	A	301	19/19	0.68	0.50	47,70,96,96	39
2	JWW	B	301	19/19	0.72	0.29	50,63,75,91	0
2	JWW	E	301	19/19	0.78	0.34	45,60,73,85	0



The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around JWW A 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around JWW B 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.