Full wwPDB X-ray Structure Validation Report (i)

Jun 16, 2024 – 02:53 AM EDT

PDB ID	:	1RGF
Title	:	HYDROLASE, GUANYLORIBONUCLEASE
Authors	:	Sevcik, J.; Dauter, Z.; Lamzin, V.S.; Wilson, K.S.
Deposited on	:	1995-06-05
Resolution	:	1.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1286 (1.22-1.18)
Ramachandran outliers	138981	1240 (1.22-1.18)
Sidechain outliers	138945	1239 (1.22-1.18)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	96	77% 20% •
1	B	96	81% 16% •

2 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3269 atoms, of which 1371 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBONUCLEASE.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	96	1435	469	683	122	159	2	0	3	0
1	B	96	1439	468	688	122	159	2	10	2	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	191	191	191	0	0
			Total	O		
3	B	194	194	194	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RIBONUCLEASE

Chain A:



- Molecule 1: RIBONUCLEASE

Chain B:



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	64.82Å 78.56Å 39.05Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 1.20 10.00 – 1.20	Depositor EDS
% Data completeness (in resolution range)	95.0 (10.00-1.20) 91.4 (10.00-1.20)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	3.92 (at 1.20Å)	Xtriage
Refinement program	SHELXL-93	Depositor
R, R_{free}	(Not available) , (Not available) 0.152 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å^2)	10.8	Xtriage
Anisotropy	0.283	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.36 , 70.9	EDS
L-test for twinning[2]	$< \|L\| > = 0.49$, $< L^2 > = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.98	EDS
Total number of atoms	3269	wwPDB-VP
Average B, all atoms (Å^2)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.12% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	1.16	1/786 (0.1%)	1.67	10/1075 (0.9%)
1	B	1.94	2/779 (0.3%)	3.40	25/1065 (2.3%)
All	All	1.59	3/1565 (0.2%)	2.67	35/2140 (1.6%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	41	GLU	CG-CD	44.31	2.18	1.51
1	A	78	GLU	CD-OE1	-6.30	1.18	1.25
1	B	31	SER	CB-OG	-6.22	1.34	1.42

All (35) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	40	ARG	NE-CZ-NH2	-70.48	85.06	120.30
1	B	40	ARG	NE-CZ-NH1	55.56	148.08	120.30
1	B	40	ARG	CD-NE-CZ	-20.30	95.18	123.60
1	B	41	GLU	CG-CD-OE1	14.57	147.45	118.30
1	B	63	ARG	NE-CZ-NH2	-12.79	113.90	120.30
1	B	41	GLU	CG-CD-OE2	-12.55	93.19	118.30
1	A	40	ARG	CD-NE-CZ	12.09	140.53	123.60
1	B	41	GLU	CB-CG-CD	-10.91	84.73	114.20
1	A	68	ARG	NE-CZ-NH2	-10.65	114.97	120.30
1	B	63	ARG	CD-NE-CZ	10.33	138.06	123.60
1	A	65	ARG	NE-CZ-NH1	-9.61	115.50	120.30
1	A	63	ARG	NE-CZ-NH2	-8.49	116.06	120.30
1	B	63	ARG	NE-CZ-NH1	8.28	124.44	120.30
1	A	63	ARG	NE-CZ-NH1	7.73	124.16	120.30
1	B	68	ARG	NE-CZ-NH2	-7.73	116.44	120.30
1	B	30	TYR	CB-CG-CD1	-7.19	116.69	121.00

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	78	GLU	OE1-CD-OE2	6.94	131.63	123.30
1	B	65	ARG	NE-CZ-NH2	-6.87	116.87	120.30
1	B	49	TYR	CB-CG-CD1	-6.73	116.96	121.00
1	B	40	ARG	NH1-CZ-NH2	6.72	126.79	119.40
1	B	86	TYR	CB-CG-CD1	6.59	124.96	121.00
1	A	17	ASP	CB-CG-OD1	-6.42	112.52	118.30
1	B	93	ASP	CB-CG-OD2	-6.27	112.65	118.30
1	A	30	TYR	CB-CG-CD1	-6.26	117.24	121.00
1	B	77	GLN	CA-CB-CG	6.16	126.96	113.40
1	B	86	TYR	CG-CD1-CE1	6.10	126.18	121.30
1	B	54[A]	GLU	CG-CD-OE2	-5.73	106.84	118.30
1	B	54[B]	GLU	CG-CD-OE2	-5.73	106.84	118.30
1	B	30	TYR	CB-CG-CD2	5.67	124.41	121.00
1	B	80	TYR	CB-CG-CD2	-5.42	117.75	121.00
1	A	84	ASP	CB-CG-OD2	-5.41	113.43	118.30
1	B	89	PHE	CD1-CE1-CZ	-5.32	113.72	120.10
1	A	5	THR	CA-CB-CG2	5.27	119.77	112.40
1	A	61	GLY	O-C-N	-5.07	114.59	122.70
1	B	51	TYR	O-C-N	5.02	130.74	122.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	752	683	699	19	0
1	B	751	688	698	2	0
2	A	10	0	0	0	0
3	A	191	0	0	16	0
3	B	194	0	0	1	0
All	All	1898	1371	1397	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:91:LEU:HB3	3:A:193:HOH:O	1.71	0.89
1:A:91:LEU:CB	3:A:193:HOH:O	2.27	0.81
1:A:41:GLU:HG3	3:A:188:HOH:O	1.84	0.77
1:A:91:LEU:CD2	3:A:258:HOH:O	2.32	0.76
1:A:91:LEU:HD22	3:A:258:HOH:O	1.89	0.72
1:A:74:GLU:OE2	3:A:160:HOH:O	2.07	0.70
1:A:91:LEU:HD13	3:A:193:HOH:O	1.96	0.64
1:A:74:GLU:CD	3:A:160:HOH:O	2.35	0.63
1:A:76:THR:HG22	1:A:77:GLN:HG2	1.79	0.63
1:A:74:GLU:HG2	3:A:261:HOH:O	2.01	0.61
1:A:31:SER:HB2	3:A:278:HOH:O	2.04	0.57
1:A:91:LEU:CG	3:A:193:HOH:O	2.50	0.55
1:A:53:HIS:HE1	3:A:181:HOH:O	1.91	0.54
1:A:74:GLU:CG	3:A:261:HOH:O	2.55	0.54
1:A:53:HIS:HD2	3:A:102:HOH:O	1.94	0.51
1:B:53:HIS:HD2	3:B:103:HOH:O	1.92	0.51
1:A:91:LEU:HD23	3:A:258:HOH:O	2.04	0.45
1:A:54[A]:GLU:OE1	1:A:86:TYR:OH	2.31	0.44
1:B:54[B]:GLU:CD	1:B:86:TYR:HH	2.20	0.43
1:A:91:LEU:HD22	3:A:193:HOH:O	2.18	0.42
1:A:80:TYR:CE1	1:A:91:LEU:HG	2.54	0.42

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	97/96 (101%)	97 (100%)	0	0	100	100
1	B	96/96 (100%)	96 (100%)	0	0	100	100
All	All	193/192 (100%)	193 (100%)	0	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	85/82 (104%)	80 (94%)	5 (6%)	19 1
1	B	84/82 (102%)	82 (98%)	2 (2%)	49 12
All	All	169/164 (103%)	162 (96%)	7 (4%)	29 4

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	PRO
1	A	25	ASP
1	A	40	ARG
1	A	74	GLU
1	A	91	LEU
1	B	40	ARG
1	B	42	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	53	HIS
1	B	53	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
2	SO4	A	98	-	4,4,4	0.76	0	6,6,6	0.88	0
2	SO4	A	97	-	4,4,4	0.63	0	6,6,6	0.33	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands (i)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers (i)

Unable to reproduce the depositors R factor - this section is therefore empty.