



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2023 – 02:50 PM EDT

PDB ID : 1RKY  
Title : PPLO + Xe  
Authors : Guss, J.M.; Duff, A.P.  
Deposited on : 2003-11-24  
Resolution : 1.68 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*



The reported resolution of this entry is 1.68 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6780 (1.70-1.66)
Clashscore	141614	7310 (1.70-1.66)
Ramachandran outliers	138981	7173 (1.70-1.66)
Sidechain outliers	138945	7172 (1.70-1.66)
RSRZ outliers	127900	6661 (1.70-1.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	747	 3% 91% 7% ..
2	B	3	 67% 33%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
9	XE	A	901	-	-	X	-
9	XE	A	907	-	-	X	-

## 2 Entry composition [i](#)

There are 10 unique types of molecules in this entry. The entry contains 6853 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called lysyl oxidase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	735	6087	3861	974	1236	16	2	49	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	314	VAL	ILE	conflict	GB 13936870
A	338	ASP	GLU	conflict	GB 13936870
A	417	SER	ALA	conflict	GB 13936870
A	478	TPQ	TYR	modified residue	GB 13936870
A	549	GLN	LYS	conflict	GB 13936870
A	577	PHE	LEU	conflict	GB 13936870
A	579	ASN	LYS	conflict	GB 13936870
A	650	THR	GLU	conflict	GB 13936870
A	758	VAL	LEU	conflict	GB 13936870
A	761	ALA	PRO	conflict	GB 13936870

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	B	3	39	22	2	15	0	0	0

- Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: C<sub>8</sub>H<sub>15</sub>NO<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	1
			28	16	2	10		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		
3	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 4 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cu	0	0
			1	1		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	2	Total	Ca	0	0
			2	2		


- Molecule 6 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	3	Total	Mg	0	0
			3	3		

- Molecule 7 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	6	Total Cl 6 6	0	0

- Molecule 8 is IMIDAZOLE (three-letter code: IMD) (formula: C<sub>3</sub>H<sub>5</sub>N<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total C N 5 3 2	0	0

- Molecule 9 is XENON (three-letter code: XE) (formula: Xe).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	8	Total Xe 8 8	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	632	Total O 632 632	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: lysyl oxidase



- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	140.14Å 66.70Å 108.62Å 90.00° 119.24° 90.00°	Depositor
Resolution (Å)	31.47 – 1.68 31.59 – 1.68	Depositor EDS
% Data completeness (in resolution range)	87.7 (31.47-1.68) 87.7 (31.59-1.68)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.63 (at 1.68Å)	Xtrriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.151 , 0.182 0.161 , 0.188	Depositor DCC
$R_{free}$ test set	3316 reflections (1.82%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	23.9	Xtrriage
Anisotropy	0.076	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 47.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	6853	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, XE, CL, MG, IMD, CA, CU, TPQ, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.59	0/6412	0.83	22/8732 (0.3%)

There are no bond length outliers.

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	642	LEU	CA-CB-CG	7.82	133.29	115.30
1	A	288	ASP	CB-CG-OD2	6.56	124.20	118.30
1	A	521[A]	ASP	CB-CG-OD2	6.43	124.09	118.30
1	A	521[B]	ASP	CB-CG-OD2	6.43	124.09	118.30
1	A	430	ASP	CB-CG-OD2	6.29	123.96	118.30
1	A	310	ASP	CB-CG-OD2	6.21	123.89	118.30
1	A	539	ASP	CB-CG-OD1	6.18	123.86	118.30
1	A	175	ASP	CB-CG-OD2	6.13	123.81	118.30
1	A	254	ASP	CB-CG-OD2	5.88	123.59	118.30
1	A	665	ASP	CB-CG-OD2	5.87	123.58	118.30
1	A	675	ASP	CB-CG-OD2	5.76	123.49	118.30
1	A	388	ASP	CB-CG-OD2	5.69	123.42	118.30
1	A	398	ASP	CB-CG-OD2	5.63	123.36	118.30
1	A	727	ARG	NE-CZ-NH1	5.60	123.10	120.30
1	A	497	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	363	ASP	CB-CG-OD2	5.38	123.14	118.30
1	A	447	ASP	CB-CG-OD2	5.36	123.12	118.30
1	A	728	ASP	CB-CG-OD2	5.31	123.08	118.30
1	A	738	ASP	CB-CG-OD2	5.20	122.98	118.30
1	A	752	ASP	CB-CG-OD2	5.18	122.96	118.30
1	A	117	ASP	CB-CG-OD2	5.16	122.94	118.30
1	A	74	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6087	0	5599	28	0
2	B	39	0	34	0	0
3	A	70	0	65	0	0
4	A	1	0	0	0	0
5	A	2	0	0	0	0
6	A	3	0	0	0	0
7	A	6	0	0	0	0
8	A	5	0	5	0	0
9	A	8	0	0	9	0
10	A	632	0	0	4	0
All	All	6853	0	5703	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:371[A]:ILE:HD12	9:A:901:XE:XE	1.76	1.63
1:A:532:LEU:CD1	9:A:907:XE:XE	2.31	1.56
1:A:371[A]:ILE:CD1	9:A:901:XE:XE	2.30	1.56
1:A:371[A]:ILE:HD11	9:A:901:XE:XE	2.07	1.23
1:A:532:LEU:HD11	9:A:907:XE:XE	2.12	1.19
1:A:532:LEU:HD13	9:A:907:XE:XE	2.09	1.16
1:A:541[A]:GLY:O	10:A:2627:HOH:O	1.66	1.11
1:A:532:LEU:HD12	9:A:907:XE:XE	2.35	1.05
1:A:463:GLN:NE2	10:A:2438:HOH:O	2.31	0.63
1:A:691:GLY:HA2	9:A:907:XE:XE	2.76	0.63
1:A:45:CYS:SG	1:A:758:VAL:HG13	2.39	0.62
1:A:553:LYS:HG2	10:A:2625:HOH:O	2.01	0.61
1:A:380[B]:LEU:HD21	1:A:480:TYR:CZ	2.38	0.58
1:A:380[B]:LEU:HD12	1:A:532:LEU:CD2	2.38	0.53

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:377:LEU:HD21	1:A:380[A]:LEU:HD11	1.92	0.52
1:A:451:LEU:HD12	1:A:451:LEU:C	2.31	0.51
1:A:380[B]:LEU:HD22	1:A:398:ASP:OD2	2.11	0.49
1:A:532:LEU:HD11	9:A:908:XE:XE	2.93	0.47
1:A:650:THR:HG23	10:A:2620:HOH:O	2.17	0.44
1:A:631:GLY:N	1:A:632:PRO:HD3	2.34	0.43
1:A:642:LEU:HD22	1:A:688:VAL:HG22	2.01	0.43
1:A:314:VAL:HG13	1:A:318:ASP:HB3	2.01	0.43
1:A:727:ARG:HA	1:A:727:ARG:CZ	2.49	0.42
1:A:231:GLY:HA3	1:A:251:PHE:CE1	2.54	0.42
1:A:504:ALA:CB	1:A:523:LEU:HD11	2.49	0.42
1:A:449[B]:SER:HB3	1:A:462:THR:O	2.20	0.41
1:A:533:ASN:HB2	1:A:637:TRP:CE3	2.56	0.40
1:A:250:PHE:HB2	1:A:269:ILE:HB	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	776/747 (104%)	754 (97%)	22 (3%)	0	<b>100</b> <b>100</b>

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	686/658 (104%)	675 (98%)	11 (2%)	62 46

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	243[A]	ASN
1	A	243[B]	ASN
1	A	310	ASP
1	A	355	ARG
1	A	388	ASP
1	A	395	PHE
1	A	642	LEU
1	A	703	ASN
1	A	739	ASP
1	A	740	GLU
1	A	769	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	173	HIS
1	A	608	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	TPQ	A	478	4,1	13,14,15	1.74	3 (23%)	15,19,21	1.60	2 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPQ	A	478	4,1	-	2/5/22/24	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	478	TPQ	C3-C4	3.34	1.40	1.35
1	A	478	TPQ	O4-C4	-3.31	1.25	1.34
1	A	478	TPQ	C6-C1	2.26	1.40	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	478	TPQ	CA-CB-C1	-4.87	104.21	113.51
1	A	478	TPQ	CB-C1-C2	2.30	122.69	118.57

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	478	TPQ	C-CA-CB-C1
1	A	478	TPQ	N-CA-CB-C1

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	NAG	B	1	2,1	14,14,15	0.70	0	17,19,21	1.00	1 (5%)
2	NAG	B	2	2	14,14,15	0.51	0	17,19,21	0.94	0
2	BMA	B	3	2	11,11,12	0.57	0	15,15,17	0.69	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2,1	-	0/6/23/26	0/1/1/1
2	NAG	B	2	2	-	0/6/23/26	0/1/1/1
2	BMA	B	3	2	-	2/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	NAG	O5-C1-C2	-2.50	107.35	111.29

There are no chirality outliers.


All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	3	BMA	C4-C5-C6-O6
2	B	3	BMA	O5-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 26 ligands modelled in this entry, 20 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	1081[A]	1	14,14,15	0.48	0	17,19,21	0.86	1 (5%)
3	NAG	A	1309	1	14,14,15	0.65	0	17,19,21	0.78	0
8	IMD	A	821	-	3,5,5	0.50	0	4,5,5	0.69	0
3	NAG	A	1191	1	14,14,15	0.38	0	17,19,21	1.10	0
3	NAG	A	1081[B]	1	14,14,15	0.56	0	17,19,21	0.77	0
3	NAG	A	1434	1	14,14,15	0.59	0	17,19,21	1.01	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1081[A]	1	-	0/6/23/26	0/1/1/1
3	NAG	A	1309	1	-	0/6/23/26	0/1/1/1
8	IMD	A	821	-	-	-	0/1/1/1
3	NAG	A	1191	1	-	1/6/23/26	0/1/1/1
3	NAG	A	1081[B]	1	-	2/6/23/26	0/1/1/1
3	NAG	A	1434	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1434	NAG	O5-C1-C2	-3.30	106.08	111.29
3	A	1081[A]	NAG	C1-O5-C5	2.02	114.93	112.19

There are no chirality outliers.

All (5) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1081[B]	NAG	O5-C5-C6-O6
3	A	1191	NAG	O5-C5-C6-O6
3	A	1434	NAG	C8-C7-N2-C2
3	A	1434	NAG	O7-C7-N2-C2
3	A	1081[B]	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	734/747 (98%)	-0.09	23 (3%) 49 51	21, 30, 52, 105	7 (0%)

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	739	ASP	6.4
1	A	740	GLU	6.2
1	A	776[A]	ILE	6.0
1	A	47	SER	4.5
1	A	737	TYR	4.5
1	A	310	ASP	4.2
1	A	741	THR	4.1
1	A	43	ALA	3.9
1	A	742	GLU	3.7
1	A	149	ASP	3.6
1	A	777[A]	ASN	3.4
1	A	46	VAL	3.3
1	A	44	GLU	3.1
1	A	738	ASP	2.9
1	A	311	SER	2.8
1	A	770[A]	TYR	2.7
1	A	743	GLU	2.7
1	A	288	ASP	2.6
1	A	45	CYS	2.6
1	A	114	GLU	2.6
1	A	289	ASP	2.3
1	A	150	GLU	2.1
1	A	457[A]	SER	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
1	TPQ	A	478	14/15	0.93	0.16	25,40,73,108	9

## 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	NAG	B	2	14/15	0.83	0.16	39,52,78,88	0
2	BMA	B	3	11/12	0.86	0.28	70,85,97,105	0
2	NAG	B	1	14/15	0.93	0.09	31,48,62,63	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAG	A	1309	14/15	0.65	0.36	66,86,102,103	0
3	NAG	A	1081[B]	14/15	0.67	0.27	30,35,50,57	14
3	NAG	A	1081[A]	14/15	0.67	0.27	35,56,70,80	14
6	MG	A	806	1/1	0.78	0.08	59,59,59,59	1
8	IMD	A	821	5/5	0.84	0.16	38,44,48,60	5
3	NAG	A	1434	14/15	0.88	0.22	44,60,78,89	0
3	NAG	A	1191	14/15	0.90	0.27	39,53,71,75	0
9	XE	A	906	1/1	0.94	0.06	43,43,43,43	1
6	MG	A	805	1/1	0.95	0.06	37,37,37,37	1
7	CL	A	814	1/1	0.95	0.08	48,48,48,48	1
9	XE	A	907	1/1	0.95	0.10	39,39,39,39	1
9	XE	A	908	1/1	0.96	0.09	39,39,39,39	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
9	XE	A	902	1/1	0.97	0.06	41,41,41,41	1
7	CL	A	813	1/1	0.98	0.11	43,43,43,43	0
9	XE	A	905	1/1	0.98	0.06	38,38,38,38	1
6	MG	A	804	1/1	0.98	0.04	31,31,31,31	0
7	CL	A	815	1/1	0.98	0.12	46,46,46,46	1
7	CL	A	811	1/1	0.98	0.04	40,40,40,40	0
9	XE	A	903	1/1	0.99	0.04	34,34,34,34	1
9	XE	A	904	1/1	0.99	0.05	38,38,38,38	1
7	CL	A	816	1/1	0.99	0.22	49,49,49,49	1
5	CA	A	803	1/1	0.99	0.04	25,25,25,25	0
9	XE	A	901	1/1	0.99	0.05	27,27,27,27	1
4	CU	A	801	1/1	0.99	0.03	32,32,32,32	0
7	CL	A	812	1/1	1.00	0.08	41,41,41,41	0
5	CA	A	802	1/1	1.00	0.06	23,23,23,23	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.