



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 2, 2023 – 11:22 PM EDT

PDB ID : 3RK2
Title : Truncated SNARE complex
Authors : Kuemmel, D.; Reinisch, K.M.
Deposited on : 2011-04-17
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	37	<div style="display: flex; align-items: center;"> <div style="width: 3%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 65%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 16%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 19%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">3% 65% 16% 19%</p>
1	E	37	<div style="display: flex; align-items: center;"> <div style="width: 62%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 24%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 14%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">62% 24% 14%</p>
2	B	65	<div style="display: flex; align-items: center;"> <div style="width: 74%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 15%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">74% 15% • 9%</p>
2	F	65	<div style="display: flex; align-items: center;"> <div style="width: 78%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 11%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">78% 11% • 9%</p>
3	C	81	<div style="display: flex; align-items: center;"> <div style="width: 0%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 77%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 17%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 5%; height: 10px; background-color: grey;"></div> </div> <p style="margin-left: 20px;">% 77% 17% • 5%</p>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	G	81	 <p>% 78% 17% 5%</p>
4	D	65	 <p>2% 85% 15%</p>
4	H	65	 <p>80% 20%</p>

2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 3825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vesicle-associated membrane protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	30	255	155	48	51	1	12	1	0
1	E	32	268	161	53	53	1	12	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	24	GLY	-	expression tag	UNP P63027
A	25	PRO	-	expression tag	UNP P63027
A	26	LEU	-	expression tag	UNP P63027
A	27	GLY	-	expression tag	UNP P63027
E	24	GLY	-	expression tag	UNP P63027
E	25	PRO	-	expression tag	UNP P63027
E	26	LEU	-	expression tag	UNP P63027
E	27	GLY	-	expression tag	UNP P63027

- Molecule 2 is a protein called Syntaxin-1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	59	481	297	81	98	5	19	0	0
2	F	59	481	297	81	98	5	28	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	189	GLY	-	expression tag	UNP P32851
B	190	SER	-	expression tag	UNP P32851
F	189	GLY	-	expression tag	UNP P32851
F	190	SER	-	expression tag	UNP P32851

- Molecule 3 is a protein called Synaptosomal-associated protein 25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	77	Total	C	N	O	S	13	1	0
			631	375	117	133	6			
3	G	77	Total	C	N	O	S	17	0	0
			630	376	115	133	6			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	3	GLY	-	expression tag	UNP P60880
C	4	SER	-	expression tag	UNP P60880
C	5	HIS	-	expression tag	UNP P60880
C	6	MET	-	expression tag	UNP P60880
C	83	TRP	-	expression tag	UNP P60880
G	3	GLY	-	expression tag	UNP P60880
G	4	SER	-	expression tag	UNP P60880
G	5	HIS	-	expression tag	UNP P60880
G	6	MET	-	expression tag	UNP P60880
G	83	TRP	-	expression tag	UNP P60880

- Molecule 4 is a protein called Synaptosomal-associated protein 25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	65	Total	C	N	O	S	7	1	0
			519	305	99	109	6			
4	H	65	Total	C	N	O	S	3	1	0
			519	305	99	109	6			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	139	GLY	-	expression tag	UNP P60880
D	140	SER	-	expression tag	UNP P60880
H	139	GLY	-	expression tag	UNP P60880
H	140	SER	-	expression tag	UNP P60880

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	2	Total	Ca	0	0
			2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	2	Total Ca 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	3	Total O 3 3	0	0
6	B	6	Total O 6 6	0	0
6	C	11	Total O 11 11	0	0
6	D	6	Total O 6 6	0	0
6	E	1	Total O 1 1	0	0
6	F	2	Total O 2 2	0	0
6	G	5	Total O 5 5	0	0
6	H	3	Total O 3 3	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Vesicle-associated membrane protein 2




- Molecule 1: Vesicle-associated membrane protein 2




- Molecule 2: Syntaxin-1A





- Molecule 2: Syntaxin-1A




- Molecule 3: Synaptosomal-associated protein 25





- Molecule 3: Synaptosomal-associated protein 25



- Molecule 4: Synaptosomal-associated protein 25



- Molecule 4: Synaptosomal-associated protein 25



4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	27.63Å 39.77Å 102.28Å 83.38° 89.94° 89.87°	Depositor
Resolution (Å)	39.50 – 2.20 39.50 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.6 (39.50-2.20) 90.5 (39.50-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.67 (at 2.00Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.227 , 0.268 0.238 , 0.276	Depositor DCC
R_{free} test set	1373 reflections (5.15%)	wwPDB-VP
Wilson B-factor (Å ²)	31.8	Xtrriage
Anisotropy	0.904	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 34.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtrriage
Estimated twinning fraction	0.457 for h,-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3825	wwPDB-VP
Average B, all atoms (Å ²)	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.25% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.92	1/257 (0.4%)	0.86	0/343
1	E	0.93	1/267 (0.4%)	1.06	1/356 (0.3%)
2	B	0.75	0/486	0.80	1/652 (0.2%)
2	F	1.07	1/486 (0.2%)	0.85	0/652
3	C	1.30	2/636 (0.3%)	0.88	3/846 (0.4%)
3	G	0.85	1/632 (0.2%)	0.83	1/840 (0.1%)
4	D	0.94	2/522 (0.4%)	0.88	1/693 (0.1%)
4	H	0.82	0/522	0.92	0/693
All	All	0.97	8/3808 (0.2%)	0.88	7/5075 (0.1%)

The worst 5 of 8 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	62	GLU	CG-CD	-24.95	1.14	1.51
2	F	245	GLU	CG-CD	15.62	1.75	1.51
4	D	197	GLN	CB-CG	-8.90	1.28	1.52
3	C	10	GLU	CB-CG	-8.05	1.36	1.52
3	G	62	GLU	CB-CG	-7.23	1.38	1.52

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	31	ARG	CA-CB-CG	10.82	137.21	113.40
3	C	62	GLU	CB-CG-CD	7.57	134.63	114.20
3	G	7	MET	CB-CG-SD	-7.00	91.41	112.40
3	C	62	GLU	CG-CD-OE1	-6.28	105.74	118.30
2	B	231	ASP	CB-CG-OD2	5.66	123.39	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	255	0	265	5	0
1	E	268	0	276	5	0
2	B	481	0	463	15	0
2	F	481	0	463	12	0
3	C	631	0	604	11	1
3	G	630	0	611	10	0
4	D	519	0	509	7	1
4	H	519	0	509	10	0
5	B	2	0	0	0	0
5	C	2	0	0	0	0
6	A	3	0	0	0	0
6	B	6	0	0	0	0
6	C	11	0	0	0	0
6	D	6	0	0	0	0
6	E	1	0	0	0	0
6	F	2	0	0	0	0
6	G	5	0	0	0	0
6	H	3	0	0	0	0
All	All	3825	0	3700	50	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 50 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:210:ARG:HG3	2:F:210:ARG:HH11	1.04	1.13
2:B:210:ARG:HG3	2:B:210:ARG:HH11	1.15	1.05
2:F:210:ARG:HH11	2:F:210:ARG:CG	1.83	0.91
2:F:210:ARG:HG3	2:F:210:ARG:NH1	1.84	0.83
1:E:39:VAL:HG22	4:H:157:ILE:HD13	1.59	0.82

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:10:GLU:OE1	4:D:202:MET:SD[1_556]	1.55	0.65

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	29/37 (78%)	29 (100%)	0	0	100	100
1	E	30/37 (81%)	30 (100%)	0	0	100	100
2	B	57/65 (88%)	57 (100%)	0	0	100	100
2	F	57/65 (88%)	56 (98%)	1 (2%)	0	100	100
3	C	76/81 (94%)	76 (100%)	0	0	100	100
3	G	75/81 (93%)	75 (100%)	0	0	100	100
4	D	64/65 (98%)	64 (100%)	0	0	100	100
4	H	64/65 (98%)	64 (100%)	0	0	100	100
All	All	452/496 (91%)	451 (100%)	1 (0%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	30/34 (88%)	28 (93%)	2 (7%)	16	18
1	E	31/34 (91%)	29 (94%)	2 (6%)	17	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	54/59 (92%)	53 (98%)	1 (2%)	57	71
2	F	54/59 (92%)	53 (98%)	1 (2%)	57	71
3	C	68/72 (94%)	66 (97%)	2 (3%)	42	54
3	G	69/72 (96%)	66 (96%)	3 (4%)	29	36
4	D	57/56 (102%)	57 (100%)	0	100	100
4	H	57/56 (102%)	55 (96%)	2 (4%)	36	46
All	All	420/442 (95%)	407 (97%)	13 (3%)	42	51

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	210	ARG
3	G	9	ASN
4	H	203	LEU
3	G	66	HIS
4	H	197	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
4	D	196	ASN
3	G	56	GLN
4	H	196	ASN
4	H	149	ASN
4	H	159	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	30/37 (81%)	0.03	1 (3%) 46 44	42, 57, 73, 79	2 (6%)
1	E	32/37 (86%)	-0.04	0 100 100	43, 57, 74, 84	2 (6%)
2	B	59/65 (90%)	-0.06	0 100 100	40, 60, 80, 84	4 (6%)
2	F	59/65 (90%)	-0.18	0 100 100	35, 57, 81, 88	6 (10%)
3	C	77/81 (95%)	-0.09	1 (1%) 77 75	40, 53, 73, 84	3 (3%)
3	G	77/81 (95%)	-0.18	1 (1%) 77 75	38, 52, 78, 83	4 (5%)
4	D	65/65 (100%)	-0.20	1 (1%) 73 72	41, 52, 67, 94	2 (3%)
4	H	65/65 (100%)	-0.29	0 100 100	38, 52, 67, 90	1 (1%)
All	All	464/496 (93%)	-0.14	4 (0%) 84 83	35, 54, 77, 94	24 (5%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	D	139	GLY	2.8
3	G	83	TRP	2.5
3	C	9	ASN	2.2
1	A	38[A]	GLN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	CA	C	2	1/1	0.84	0.15	64,64,64,64	0
5	CA	B	4	1/1	0.92	0.14	73,73,73,73	0
5	CA	B	1	1/1	0.93	0.13	69,69,69,69	0
5	CA	C	84	1/1	0.97	0.18	67,67,67,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.