

wwPDB Geometry-Only Validation Summary Report (i)

Feb 4, 2024 – 10:52 AM EST

PDB ID	:	1RMN
Title	:	A THREE-DIMENSIONAL MODEL FOR THE HAMMERHEAD RI-
		BOZYME BASED ON FLUORESCENCE MEASUREMENTS
Authors	:	Tuschl, T.; Gohlke, C.; Jovin, T.M.; Westhof, E.; Eckstein, F.
Deposited on	:	1994-10-20
Resolution	:	Not provided

This is a wwPDB Geometry-Only Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:


MolProbity	:	4.02b-467
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $FLUORESCENCE\ TRANSFER$

The reported resolution of this entry is unknown.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
Clashscore	141614	-
RNA backbone	3102	-

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Note EDS was not executed.

Mol	Chain	Length		Quality of chain	
1	А	49	33%	45%	22%



2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 1053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a RNA chain called RNA (49-MER).

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace		
1	А	49	Total 1053	C 469	N 198	0 337	Р 49	0	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: RNA (49-MER)





4 Model quality (i)

4.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Boi	nd lengths	Bond angles		
10101	Mol Chain		# Z > 5	RMSZ	# Z > 5	
1	А	0.93	1/1178~(0.1%)	2.27	138/1836~(7.5%)	

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	А	28	А	P-O5'	5.77	1.65	1.59

The worst 5 of 138 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Ζ	$Observed(^{o})$	$Ideal(^{o})$
1	А	43	U	C2-N3-C4	-8.08	122.16	127.00
1	А	43	U	N3-C4-C5	7.97	119.38	114.60
1	А	10	U	C2-N3-C4	-7.92	122.25	127.00
1	А	7	U	C2-N3-C4	-7.78	122.33	127.00
1	А	30	U	C2-N3-C4	-7.71	122.37	127.00

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	1053	0	535	63	0
All	All	1053	0	535	63	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 40.

The worst 5 of 63 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:9:A:H61	1:A:27:A:H3'	1.19	1.03
1:A:23:C:H2'	1:A:24:C:C6	2.07	0.89
1:A:4:C:H2'	1:A:5:U:C6	2.08	0.89
1:A:9:A:N6	1:A:27:A:H3'	1.91	0.83
1:A:9:A:H61	1:A:27:A:C3'	1.92	0.82

There are no symmetry-related clashes.

4.3 Torsion angles (i)

4.3.1 Protein backbone (i)

There are no protein molecules in this entry.

4.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

4.3.3 RNA (i)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	А	48/49~(97%)	13~(27%)	1 (2%)

5 of 13 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
1	А	8	G
1	А	9	А
1	А	10	U
1	А	12	А
1	А	13	G

All (1) RNA pucker outliers are listed below:



Mol	Chain	Res	Type
1	А	12	А

4.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates (i)

There are no monosaccharides in this entry.

4.6 Ligand geometry (i)

There are no ligands in this entry.

4.7 Other polymers (i)

There are no such residues in this entry.

4.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

