



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 1, 2022 – 02:14 PM EDT

PDB ID : 5RMD  
Title : PanDDA analysis group deposition – Crystal Structure of SARS-CoV-2 heli-  
case in complex with Z57614330  
Authors : Newman, J.A.; Yosaatmadja, Y.; Douangamath, A.; Aimon, A.; Powell, A.J.;  
Dias, A.; Fearon, D.; Dunnett, L.; Brandao-Neto, J.; Krojer, T.; Skyner, R.;  
Gorrie-Stone, T.; Thompson, W.; von Delft, F.; Arrowsmith, C.H.; Edwards,  
A.; Bountra, C.; Gileadi, O.  
Deposited on : 2020-09-16  
Resolution : 1.92 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at <http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.29  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.29


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.92 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	7937 (1.94-1.90)
Clashscore	141614	8644 (1.94-1.90)
Ramachandran outliers	138981	8530 (1.94-1.90)
Sidechain outliers	138945	8530 (1.94-1.90)
RSRZ outliers	127900	7793 (1.94-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	601	
1	B	601	

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 9436 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Helicase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	572	Total 4417	C 2816	N 737	O 832	S 32	0	0	0
1	B	585	Total 4508	C 2875	N 750	O 848	S 35	0	1	0

- Molecule 2 is N-ethyl-4-[(methylsulfonyl)amino]benzamide (three-letter code: VWY) (formula:  $C_{10}H_{14}N_2O_3S$ ) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	Total 16	C 10	N 2	O 3	S 1	0	0
2	B	1	Total 16	C 10	N 2	O 3	S 1	0	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total	Zn	0	0
			3	3		
3	B	3	Total	Zn	0	0
			3	3		

- Molecule 4 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	P	0	0
			5	4	1		
4	A	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		
4	B	1	Total	O	P	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	205	Total	O	0	0
			205	205		
5	B	248	Total	O	0	0
			248	248		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Helicase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.48Å 70.42Å 85.58Å 102.71° 95.94° 112.61°	Depositor
Resolution (Å)	81.65 – 1.92 81.65 – 1.92	Depositor EDS
% Data completeness (in resolution range)	94.6 (81.65-1.92) 94.7 (81.65-1.92)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.08 (at 1.92Å)	Xtrriage
Refinement program	REFMAC 5.8.0258	Depositor
R, $R_{free}$	0.217 , 0.261 0.231 , 0.272	Depositor DCC
$R_{free}$ test set	4488 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	32.9	Xtrriage
Anisotropy	0.187	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 36.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	9436	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	50.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, ZN, VWY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.71	0/4517	0.81	0/6156
1	B	0.77	4/4610 (0.1%)	0.87	2/6283 (0.0%)
All	All	0.74	4/9127 (0.0%)	0.84	2/12439 (0.0%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	23	PRO	N-CA	12.18	1.68	1.47
1	B	238	PRO	N-CA	11.42	1.66	1.47
1	B	136	GLU	CD-OE2	-5.53	1.19	1.25
1	B	237	ALA	C-N	5.09	1.44	1.34

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	238	PRO	CA-N-CD	-6.24	102.76	111.50
1	B	23	PRO	CA-N-CD	-5.96	103.16	111.50

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4417	0	4321	34	1
1	B	4508	0	4425	39	1
2	A	16	0	0	0	0
2	B	16	0	0	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0
4	A	10	0	0	0	0
4	B	10	0	0	0	0
5	A	205	0	0	1	0
5	B	248	0	0	4	0
All	All	9436	0	8746	71	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:23:PRO:N	1:B:23:PRO:CA	1.68	1.46
1:B:23:PRO:N	1:B:23:PRO:C	2.47	0.67
1:A:352:LEU:HD11	1:B:234:PRO:HD3	1.80	0.64
1:B:238:PRO:HB2	1:B:240:LEU:O	1.97	0.63
1:B:6:VAL:HA	1:B:129:ARG:HE	1.64	0.62
1:B:510:VAL:HG21	1:B:541:TYR:CD1	2.34	0.62
1:B:8:CYS:SG	1:B:99:GLY:N	2.73	0.62
1:B:13:SER:O	1:B:44:SER:HA	2.01	0.60
1:A:293:ILE:HG13	1:A:320:LYS:HB3	1.83	0.59
1:B:477:LYS:NZ	1:B:551:GLU:OE2	2.32	0.58
1:B:474[B]:MET:HG2	1:B:590:LEU:HB2	1.85	0.58
1:A:510:VAL:HG21	1:A:541:TYR:CD1	2.40	0.56
1:B:103:VAL:HG12	1:B:103:VAL:O	2.05	0.55
1:B:474[A]:MET:SD	1:B:495:VAL:HG11	2.47	0.54
1:A:12:THR:HG21	1:A:26:CYS:HA	1.88	0.54
1:A:445:PRO:HB3	1:A:468:SER:HB3	1.90	0.54
1:A:277:TYR:HA	1:A:396:TYR:O	2.08	0.53
1:A:368:ALA:O	1:A:393:ALA:HA	2.08	0.53
1:B:31:TYR:CZ	1:B:35:ILE:HG21	2.45	0.52
1:B:195:ILE:HG23	1:B:195:ILE:O	2.10	0.52
1:A:260:ASP:HA	1:A:263:SER:OG	2.09	0.52
1:B:120:TYR:CE2	1:B:409:ARG:HG2	2.44	0.51
1:B:277:TYR:HA	1:B:396:TYR:O	2.10	0.51
1:B:333:ILE:HB	1:B:358:CYS:SG	2.50	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:512:ILE:O	1:A:546:PHE:HA	2.12	0.50
1:A:304:ILE:HG12	1:A:370:ILE:HB	1.93	0.50
1:A:519:ASN:HB3	1:A:530:THR:HG23	1.94	0.49
1:A:60:VAL:HB	5:A:923:HOH:O	2.13	0.49
1:A:281:GLN:HB2	1:A:429:MET:HE2	1.94	0.49
1:A:59:ASP:OD1	1:A:61:THR:OG1	2.25	0.48
1:A:12:THR:CG2	1:A:26:CYS:HA	2.44	0.48
1:B:252:LEU:HB3	1:B:299:TYR:CD1	2.49	0.48
1:A:519:ASN:HB3	1:A:530:THR:CG2	2.43	0.47
1:B:8:CYS:O	1:B:9:ASN:HB2	2.14	0.47
1:B:127:THR:HG23	5:B:970:HOH:O	2.14	0.47
1:B:544:VAL:O	1:B:572:ILE:HA	2.15	0.47
1:B:504:PRO:HB3	1:B:507:ARG:HH21	1.80	0.46
1:A:215:THR:HG22	1:B:193:VAL:HG21	1.97	0.46
1:A:183:THR:O	1:A:225:PHE:HA	2.15	0.46
1:B:124:ASN:OD1	1:B:381:ASN:ND2	2.48	0.46
1:B:183:THR:OG1	1:B:228:THR:OG1	2.22	0.46
1:B:508:LYS:HD3	5:B:905:HOH:O	2.15	0.45
1:A:304:ILE:HA	1:A:370:ILE:O	2.15	0.45
1:A:120:TYR:CE2	1:A:409:ARG:HG2	2.52	0.45
1:A:367:THR:HA	1:A:392:ARG:O	2.17	0.45
1:B:465:LYS:NZ	5:B:808:HOH:O	2.36	0.45
1:A:252:LEU:HB3	1:A:299:TYR:CD2	2.53	0.44
1:B:376:ILE:HD12	1:B:376:ILE:HA	1.91	0.44
1:A:456:VAL:HG23	1:A:457:TYR:CD2	2.53	0.44
1:A:152:ALA:HB2	1:A:167:TRP:CZ3	2.52	0.44
1:B:376:ILE:HG12	1:B:425:VAL:HG11	2.00	0.43
1:A:14:LEU:HB2	1:A:25:LEU:O	2.19	0.43
1:B:121:ILE:HG23	1:B:421:TYR:CE1	2.54	0.43
1:A:296:ALA:O	1:A:300:PRO:HA	2.19	0.43
1:B:13:SER:HB2	1:B:92:LEU:HB2	2.00	0.43
1:B:533:VAL:HG11	1:B:560:ARG:O	2.19	0.43
1:B:86:ASN:HB3	5:B:913:HOH:O	2.18	0.42
1:B:143:GLU:HA	1:B:146:LYS:HD2	2.00	0.42
1:B:451:THR:HG21	1:B:585:LEU:HD23	2.02	0.42
1:A:64:TYR:O	1:A:70:TYR:HA	2.19	0.41
1:A:13:SER:O	1:A:44:SER:HA	2.20	0.41
1:A:472:PHE:HA	1:A:588:THR:O	2.19	0.41
1:A:157:VAL:HG21	1:A:219:LEU:O	2.20	0.41
1:A:269:TYR:OH	1:A:294:GLY:HA3	2.21	0.41
1:A:16:CYS:O	1:A:22:ARG:HA	2.20	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:152:ALA:HB2	1:A:167:TRP:CH2	2.56	0.41
1:B:69:SER:OG	1:B:71:TYR:OH	2.29	0.40
1:B:167:TRP:CZ3	1:B:174:PRO:HD2	2.56	0.40
1:A:182:PHE:HB3	1:A:225:PHE:HB3	2.04	0.40
1:B:377:SER:O	1:B:406:PRO:HA	2.21	0.40
1:B:266:VAL:O	1:B:270:GLN:HG3	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:501:THR:O	1:B:442:ARG:NH1[1_444]	2.15	0.05

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	564/601 (94%)	537 (95%)	24 (4%)	3 (0%)	29 18
1	B	580/601 (96%)	546 (94%)	32 (6%)	2 (0%)	41 31
All	All	1144/1202 (95%)	1083 (95%)	56 (5%)	5 (0%)	34 24

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	351	THR
1	B	9	ASN
1	A	484	VAL
1	A	257	ASN
1	B	161	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	485/523 (93%)	469 (97%)	16 (3%)	38	28
1	B	498/523 (95%)	482 (97%)	16 (3%)	39	29
All	All	983/1046 (94%)	951 (97%)	32 (3%)	38	28

All (32) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	LEU
1	A	12	THR
1	A	35	ILE
1	A	46	ASN
1	A	76	LYS
1	A	81	PHE
1	A	173	ARG
1	A	207	ASP
1	A	209	VAL
1	A	217	TYR
1	A	231	THR
1	A	236	SER
1	A	255	THR
1	A	373	PHE
1	A	392	ARG
1	A	530	THR
1	B	7	LEU
1	B	12	THR
1	B	26	CYS
1	B	76	LYS
1	B	95	ASN
1	B	124	ASN
1	B	160	ASP
1	B	164	HIS
1	B	192	LYS
1	B	215	THR
1	B	220	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	223	ASP
1	B	353	GLU
1	B	373	PHE
1	B	484	VAL
1	B	486	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	86	ASN
1	B	179	ASN
1	B	268	ASN
1	B	516	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 6 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	VWY	B	701	-	16,16,16	0.33	0	22,22,22	0.42	0
4	PO4	B	705	-	4,4,4	0.75	0	6,6,6	0.40	0
4	PO4	A	706	-	4,4,4	1.10	0	6,6,6	0.71	0
4	PO4	A	705	-	4,4,4	0.62	0	6,6,6	0.75	0
2	VWY	A	701	-	16,16,16	0.21	0	22,22,22	0.28	0
4	PO4	B	706	-	4,4,4	0.90	0	6,6,6	0.41	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '2' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	VWY	A	701	-	-	2/12/12/12	0/1/1/1
2	VWY	B	701	-	-	4/12/12/12	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	701	VWY	C4-C3-N1-C2
2	A	701	VWY	O1-C3-N1-C2
2	B	701	VWY	O1-C3-C4-C5
2	B	701	VWY	N1-C3-C4-C5
2	B	701	VWY	O1-C3-C4-C10
2	B	701	VWY	N1-C3-C4-C10

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	572/601 (95%)	0.15	36 (6%) 20 22	24, 48, 106, 130	0
1	B	585/601 (97%)	0.03	26 (4%) 34 37	22, 40, 91, 140	0
All	All	1157/1202 (96%)	0.09	62 (5%) 25 29	22, 44, 97, 140	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	217	TYR	6.8
1	A	229	SER	6.3
1	A	202	LYS	6.0
1	A	169	VAL	5.4
1	B	96	THR	5.2
1	A	179	ASN	5.1
1	B	340	VAL	5.0
1	A	230	HIS	4.7
1	A	228	THR	4.6
1	A	339	ARG	4.6
1	B	45	VAL	4.3
1	B	53	PRO	4.3
1	A	176	LEU	4.2
1	A	215	THR	4.2
1	A	170	GLY	4.0
1	A	149	TYR	4.0
1	A	200	PHE	3.8
1	A	216	THR	3.6
1	A	208	ALA	3.6
1	A	225	PHE	3.5
1	A	195	ILE	3.5
1	B	67	GLY	3.5
1	A	167	TRP	3.4
1	B	193	VAL	3.4

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	91	GLY	3.2
1	A	210	VAL	3.2
1	B	95	ASN	3.2
1	B	218	LYS	3.1
1	A	81	PHE	3.1
1	B	29	CYS	3.0
1	A	256	LEU	2.9
1	A	217	TYR	2.9
1	A	156	GLU	2.8
1	A	219	LEU	2.8
1	A	214	THR	2.8
1	A	227	LEU	2.8
1	B	196	GLY	2.8
1	A	207	ASP	2.7
1	B	66	GLY	2.6
1	B	65	LEU	2.6
1	A	201	GLU	2.6
1	A	182	PHE	2.5
1	A	247	VAL	2.5
1	A	180	TYR	2.5
1	B	71	TYR	2.4
1	B	69	SER	2.4
1	B	8	CYS	2.4
1	B	81	PHE	2.3
1	B	54	GLY	2.3
1	B	157	VAL	2.2
1	B	55	CYS	2.2
1	A	178	ARG	2.2
1	A	177	ASN	2.2
1	B	190	ASN	2.1
1	A	154	VAL	2.1
1	A	337	ARG	2.1
1	B	178	ARG	2.1
1	B	2	VAL	2.1
1	A	153	THR	2.0
1	B	341	GLU	2.0
1	A	257	ASN	2.0
1	B	7	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	VWY	A	701	16/16	0.72	0.32	50,56,60,61	16
2	VWY	B	701	16/16	0.78	0.33	33,37,40,44	16
3	ZN	B	703	1/1	0.90	0.04	74,74,74,74	0
3	ZN	B	704	1/1	0.91	0.08	87,87,87,87	0
4	PO4	B	706	5/5	0.96	0.15	37,40,45,61	0
4	PO4	B	705	5/5	0.97	0.09	39,43,43,48	0
3	ZN	A	704	1/1	0.97	0.05	62,62,62,62	0
4	PO4	A	706	5/5	0.98	0.10	27,36,40,44	0
3	ZN	A	703	1/1	0.98	0.09	43,43,43,43	0
3	ZN	B	702	1/1	0.98	0.07	42,42,42,42	0
4	PO4	A	705	5/5	0.99	0.10	33,38,42,43	0
3	ZN	A	702	1/1	0.99	0.07	47,47,47,47	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around VWY A 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)

**Electron density around VWY B 701:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers [i](#)

There are no such residues in this entry.