



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 3, 2024 – 03:40 am BST

PDB ID : 8RP3
Title : Alpha-Methylacyl-CoA racemase from Mycobacterium tuberculosis (H126A mutant)
Authors : Mojanaga, O.O.; Acharya, K.R.; Lloyd, M.D.
Deposited on : 2024-01-12
Resolution : 2.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.4, CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1544 (2.48-2.44)
Clashscore	141614	1613 (2.48-2.44)
Ramachandran outliers	138981	1598 (2.48-2.44)
Sidechain outliers	138945	1598 (2.48-2.44)
RSRZ outliers	127900	1523 (2.48-2.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	365	 % 74% 19% ...
1	B	365	 % 77% 18% ...

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 10751 atoms, of which 5305 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-methylacyl-CoA racemase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	354	5337	1688	2647	480	506	16	54	4	0
1	B	354	5323	1685	2640	479	503	16	54	2	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	MET	-	initiating methionine	UNP O06543
A	126	ALA	HIS	engineered mutation	UNP O06543
A	361	GLY	-	expression tag	UNP O06543
A	362	SER	-	expression tag	UNP O06543
A	363	GLY	-	expression tag	UNP O06543
A	364	CYS	-	expression tag	UNP O06543
B	0	MET	-	initiating methionine	UNP O06543
B	126	ALA	HIS	engineered mutation	UNP O06543
B	361	GLY	-	expression tag	UNP O06543
B	362	SER	-	expression tag	UNP O06543
B	363	GLY	-	expression tag	UNP O06543
B	364	CYS	-	expression tag	UNP O06543

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
2	A	1	Total	C	H	O	2	0
			10	2	6	2		
2	A	1	Total	C	H	O	2	0
			10	2	6	2		
2	B	1	Total	C	H	O	2	0
			10	2	6	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	28	Total	O	0	0
			28	28		
3	B	33	Total	O	0	0
			33	33		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Alpha-methylacyl-CoA racemase



- Molecule 1: Alpha-methylacyl-CoA racemase



4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	181.11Å 79.15Å 59.10Å 90.00° 91.66° 90.00°	Depositor
Resolution (Å)	90.52 – 2.45 90.52 – 2.45	Depositor EDS
% Data completeness (in resolution range)	100.0 (90.52-2.45) 99.8 (90.52-2.45)	Depositor EDS
R_{merge}	0.37	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 2.45Å)	Xtrriage
Refinement program	REFMAC 5.8.0425, REFMAC 5.8.0425	Depositor
R, R_{free}	0.188 , 0.262 0.198 , 0.265	Depositor DCC
R_{free} test set	1467 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å ²)	40.0	Xtrriage
Anisotropy	0.411	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 5.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	0.218 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10751	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.83	2/2769 (0.1%)	1.54	39/3766 (1.0%)
1	B	0.80	2/2752 (0.1%)	1.43	33/3743 (0.9%)
All	All	0.82	4/5521 (0.1%)	1.49	72/7509 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	7
1	B	0	4
All	All	0	11

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	12	GLU	CD-OE2	7.07	1.33	1.25
1	B	145	GLU	CD-OE2	5.47	1.31	1.25
1	A	321	GLU	CD-OE1	5.16	1.31	1.25
1	A	12	GLU	CD-OE2	5.04	1.31	1.25

All (72) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	MET	CG-SD-CE	18.99	130.58	100.20
1	A	203	ARG	NE-CZ-NH2	-13.18	113.71	120.30
1	A	212	ARG	NE-CZ-NH1	-12.45	114.08	120.30
1	A	284	ARG	NE-CZ-NH1	10.23	125.41	120.30
1	A	267	ARG	NE-CZ-NH1	-9.79	115.41	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	284	ARG	NE-CZ-NH2	-9.59	115.51	120.30
1	B	35	ARG	NE-CZ-NH2	-9.47	115.57	120.30
1	B	120	ARG	NE-CZ-NH1	-9.21	115.70	120.30
1	A	333	ARG	NE-CZ-NH1	-9.04	115.78	120.30
1	A	35	ARG	NE-CZ-NH2	-8.78	115.91	120.30
1	A	203	ARG	NE-CZ-NH1	8.63	124.61	120.30
1	A	62	LYS	CB-CA-C	8.62	127.65	110.40
1	B	12	GLU	OE1-CD-OE2	8.55	133.56	123.30
1	A	321	GLU	OE1-CD-OE2	8.54	133.55	123.30
1	B	54	ARG	CB-CA-C	-8.45	93.50	110.40
1	A	146	ARG	NE-CZ-NH2	8.17	124.38	120.30
1	A	54	ARG	NE-CZ-NH1	-7.97	116.32	120.30
1	B	54	ARG	NE-CZ-NH2	-7.79	116.40	120.30
1	B	333	ARG	NE-CZ-NH1	-7.71	116.44	120.30
1	A	47	ARG	NE-CZ-NH1	7.55	124.08	120.30
1	B	216	MET	CG-SD-CE	-7.39	88.37	100.20
1	B	212	ARG	NE-CZ-NH2	-7.33	116.64	120.30
1	B	48	ASP	CB-CG-OD1	7.12	124.71	118.30
1	A	207	MET	CG-SD-CE	-7.08	88.88	100.20
1	B	35	ARG	NE-CZ-NH1	7.00	123.80	120.30
1	B	123	GLN	N-CA-CB	-6.96	98.08	110.60
1	B	252	LEU	CB-CG-CD1	-6.92	99.23	111.00
1	A	48	ASP	CB-CG-OD1	6.89	124.50	118.30
1	B	120	ARG	NE-CZ-NH2	6.87	123.73	120.30
1	B	265	ARG	NE-CZ-NH2	-6.69	116.96	120.30
1	B	207	MET	CG-SD-CE	-6.36	90.02	100.20
1	B	203	ARG	NE-CZ-NH2	-6.34	117.13	120.30
1	A	48	ASP	CB-CG-OD2	-6.30	112.63	118.30
1	A	71	LEU	CB-CG-CD1	6.29	121.68	111.00
1	A	54	ARG	CB-CA-C	-6.24	97.92	110.40
1	A	270	GLU	CG-CD-OE2	-6.11	106.09	118.30
1	A	241	GLU	CG-CD-OE1	6.07	130.44	118.30
1	A	85	ARG	NE-CZ-NH1	6.04	123.32	120.30
1	B	283[A]	ASP	CB-CG-OD1	5.98	123.69	118.30
1	B	283[B]	ASP	CB-CG-OD1	5.98	123.69	118.30
1	A	295	ASP	CB-CG-OD2	-5.91	112.98	118.30
1	A	12	GLU	OE1-CD-OE2	5.90	130.38	123.30
1	B	285	ASP	CB-CA-C	5.89	122.18	110.40
1	B	254	LEU	CB-CG-CD2	5.74	120.75	111.00
1	A	137	LEU	CB-CG-CD1	5.71	120.72	111.00
1	A	212	ARG	CG-CD-NE	5.69	123.75	111.80
1	A	101	LYS	CB-CG-CD	5.66	126.32	111.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	39	PRO	N-CA-C	5.63	126.74	112.10
1	B	85	ARG	CD-NE-CZ	5.62	131.47	123.60
1	A	258	GLU	CB-CA-C	-5.55	99.30	110.40
1	A	241	GLU	CG-CD-OE2	-5.51	107.28	118.30
1	B	284	ARG	NE-CZ-NH1	5.50	123.05	120.30
1	B	71	LEU	CB-CG-CD1	5.43	120.24	111.00
1	B	145	GLU	CB-CG-CD	5.35	128.65	114.20
1	A	198	MET	CG-SD-CE	-5.33	91.67	100.20
1	A	85	ARG	N-CA-CB	5.29	120.11	110.60
1	B	334	PHE	N-CA-CB	-5.28	101.10	110.60
1	B	145	GLU	OE1-CD-OE2	5.25	129.59	123.30
1	B	47	ARG	NE-CZ-NH2	5.24	122.92	120.30
1	A	50	MET	N-CA-CB	5.22	119.99	110.60
1	A	137	LEU	CB-CG-CD2	-5.21	102.15	111.00
1	A	120	ARG	NE-CZ-NH1	-5.20	117.70	120.30
1	A	356	THR	CA-CB-OG1	-5.19	98.09	109.00
1	B	127	ASP	CB-CG-OD1	-5.15	113.67	118.30
1	A	150	PRO	N-CA-CB	-5.14	96.94	102.60
1	B	54	ARG	CA-CB-CG	5.13	124.68	113.40
1	B	37	ASP	CB-CG-OD1	-5.12	113.69	118.30
1	A	92	LEU	CB-CG-CD2	5.08	119.64	111.00
1	B	47	ARG	CD-NE-CZ	5.07	130.70	123.60
1	A	258	GLU	N-CA-CB	5.04	119.68	110.60
1	B	319	PHE	N-CA-CB	-5.04	101.53	110.60
1	A	223	TYR	CB-CG-CD1	5.02	124.01	121.00

There are no chirality outliers.

All (11) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	105	ARG	Sidechain
1	A	203	ARG	Sidechain
1	A	212	ARG	Sidechain
1	A	233	ARG	Sidechain
1	A	265	ARG	Sidechain
1	A	267	ARG	Sidechain
1	A	55	ARG	Sidechain
1	B	105	ARG	Sidechain
1	B	203	ARG	Sidechain
1	B	212	ARG	Sidechain
1	B	267	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2690	2647	2631	39	0
1	B	2683	2640	2631	33	0
2	A	8	12	12	0	0
2	B	4	6	6	0	0
3	A	28	0	0	2	0
3	B	33	0	0	3	0
All	All	5446	5305	5280	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:THR:HG22	1:A:211:THR:H	1.28	0.98
1:B:326:TRP:O	1:B:327:GLN:NE2	2.28	0.66
1:B:16:ILE:O	1:B:18:PRO:HD2	1.96	0.64
1:B:144:ASP:O	1:B:145:GLU:HG2	1.97	0.64
1:B:206:GLY:HA2	3:B:501:HOH:O	1.98	0.64
1:B:144:ASP:C	1:B:145:GLU:HG2	2.20	0.62
1:A:16:ILE:O	1:A:18:PRO:HD2	2.01	0.59
1:A:209:THR:CG2	1:A:211:THR:HB	2.34	0.58
1:B:345:PRO:HA	3:B:528:HOH:O	2.05	0.56
1:B:80:LEU:HD23	1:B:108:TYR:CE2	2.40	0.56
1:A:209:THR:HG22	1:A:211:THR:N	2.11	0.55
1:A:259:LEU:HD22	1:A:274:LEU:HD13	1.88	0.55
1:B:348:THR:HG22	1:B:349:ILE:N	2.23	0.54
1:A:212:ARG:NH2	1:A:212:ARG:CB	2.71	0.53
1:A:212:ARG:NH2	1:A:212:ARG:HB2	2.25	0.52
1:A:135:GLY:HA2	1:B:302:LEU:O	2.11	0.51
1:B:199:MET:O	1:B:203:ARG:HG3	2.11	0.51
1:A:209:THR:HG22	1:A:211:THR:HB	1.93	0.50
1:A:212:ARG:HB2	1:A:212:ARG:HH21	1.76	0.50
1:B:145:GLU:O	1:B:212:ARG:NH2	2.43	0.50
1:A:138:HIS:O	1:A:212:ARG:HD3	2.12	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:268:TRP:HD1	3:A:520:HOH:O	1.95	0.49
1:A:80:LEU:HD23	1:A:108:TYR:CE2	2.47	0.49
1:A:128:ILE:HG12	1:B:236:ALA:HB1	1.95	0.49
1:A:136:ILE:HD13	1:A:192:SER:HB2	1.94	0.48
1:A:209:THR:HG21	1:A:211:THR:HB	1.96	0.48
1:B:66:GLY:O	1:B:67:LEU:C	2.52	0.48
1:B:113:GLY:HA3	1:B:130:TYR:CE1	2.49	0.48
1:B:347:ALA:O	1:B:348:THR:O	2.32	0.47
1:A:196:ILE:HG12	1:A:199:MET:HB2	1.96	0.47
1:A:113:GLY:HA3	1:A:130:TYR:CE1	2.50	0.46
1:A:344:PRO:HB2	1:A:345:PRO:HD2	1.96	0.46
1:B:320:TYR:CE1	1:B:327:GLN:HB2	2.50	0.45
1:A:217:LEU:O	1:B:151:LEU:HA	2.17	0.45
1:A:107:ILE:HD12	1:A:171:ALA:HB1	1.98	0.45
1:A:224:TYR:HA	1:A:237:VAL:O	2.17	0.45
1:A:66:GLY:O	1:A:67:LEU:C	2.55	0.45
1:A:127:ASP:OD2	1:A:155:GLY:N	2.47	0.45
1:B:53:ASN:ND2	1:B:342:PRO:HB3	2.33	0.44
1:A:150:PRO:HG2	3:A:517:HOH:O	2.17	0.44
1:B:113:GLY:HA3	1:B:130:TYR:CZ	2.52	0.44
1:A:107:ILE:HD12	1:A:171:ALA:CB	2.48	0.44
1:B:115:GLY:HA2	1:B:187:ALA:HB2	2.00	0.44
1:A:98:GLU:O	1:A:101:LYS:HB2	2.18	0.44
1:B:196:ILE:HG12	1:B:199:MET:HB2	1.99	0.44
1:B:334:PHE:CD1	1:B:334:PHE:N	2.86	0.43
1:A:334:PHE:CD1	1:A:334:PHE:N	2.86	0.43
1:B:241:GLU:HA	1:B:242:PRO:HD3	1.87	0.43
1:B:326:TRP:C	1:B:327:GLN:HE21	2.19	0.43
1:B:150:PRO:HG2	3:B:520:HOH:O	2.17	0.43
1:B:49:ALA:C	1:B:51:LEU:N	2.72	0.42
1:A:46:SER:O	1:A:345:PRO:HG3	2.19	0.42
1:A:241:GLU:HA	1:A:242:PRO:HD3	1.85	0.42
1:B:241:GLU:HB2	1:B:244:PHE:CD2	2.54	0.42
1:A:115:GLY:HA2	1:A:187:ALA:HB2	2.02	0.41
1:A:199:MET:O	1:A:203:ARG:HG3	2.21	0.41
1:A:173:TRP:HD1	1:B:4:PRO:O	2.04	0.41
1:B:251:GLY:HA3	1:B:291:PHE:CE1	2.56	0.41
1:A:241:GLU:HB2	1:A:244:PHE:CD2	2.55	0.41
1:A:148:VAL:HA	1:A:149:PRO:HD3	1.89	0.41
1:B:109:ALA:HB1	1:B:164:LEU:HD11	2.03	0.41
1:A:113:GLY:HA3	1:A:130:TYR:CZ	2.56	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:218:ASP:C	1:A:218:ASP:OD1	2.60	0.40
1:A:144:ASP:O	1:A:145:GLU:HG3	2.20	0.40
1:B:212:ARG:HH21	1:B:212:ARG:HD3	1.68	0.40
1:A:255:ASP:HB3	1:A:258:GLU:HG2	2.04	0.40
1:B:323:ASN:O	1:B:324:GLY:C	2.60	0.40
1:B:348:THR:CG2	1:B:349:ILE:N	2.85	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	354/365 (97%)	330 (93%)	22 (6%)	2 (1%)	25	29
1	B	352/365 (96%)	328 (93%)	22 (6%)	2 (1%)	25	29
All	All	706/730 (97%)	658 (93%)	44 (6%)	4 (1%)	25	29

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	348	THR
1	B	95	GLY
1	A	95	GLY
1	A	346	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	274/277 (99%)	262 (96%)	12 (4%)	28	37
1	B	272/277 (98%)	264 (97%)	8 (3%)	42	53
All	All	546/554 (99%)	526 (96%)	20 (4%)	34	45

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	29	LEU
1	A	37	ASP
1	A	54	ARG
1	A	62	LYS
1	A	102	VAL
1	A	133	LEU
1	A	140	ILE
1	A	148	VAL
1	A	149	PRO
1	A	199	MET
1	A	212	ARG
1	A	216	MET
1	B	37	ASP
1	B	56	ILE
1	B	102	VAL
1	B	133	LEU
1	B	176	GLN
1	B	199	MET
1	B	209	THR
1	B	321	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	138	HIS
1	A	176	GLN
1	B	176	GLN
1	B	263	ASN
1	B	327	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	402	-	3,3,3	0.92	0	2,2,2	0.93	0
2	EDO	B	401	-	3,3,3	0.70	0	2,2,2	1.35	0
2	EDO	A	401	-	3,3,3	0.38	0	2,2,2	0.76	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	402	-	-	1/1/1/1	-
2	EDO	B	401	-	-	1/1/1/1	-
2	EDO	A	401	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	EDO	O1-C1-C2-O2
2	B	401	EDO	O1-C1-C2-O2
2	A	402	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/365 (96%)	-0.73	4 (1%) 80 80	26, 44, 81, 123	0
1	B	354/365 (96%)	-0.75	4 (1%) 80 80	27, 44, 81, 106	0
All	All	708/730 (96%)	-0.74	8 (1%) 80 80	26, 44, 81, 123	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	47	ARG	6.3
1	B	47	ARG	4.8
1	B	45	ILE	3.9
1	B	16	ILE	3.9
1	A	45	ILE	3.3
1	A	324	GLY	2.6
1	A	323	ASN	2.2
1	B	346	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	A	401	4/4	0.94	0.27	38,47,52,52	2
2	EDO	A	402	4/4	0.97	0.17	34,49,53,58	2
2	EDO	B	401	4/4	0.97	0.22	34,44,54,54	2

6.5 Other polymers [i](#)

There are no such residues in this entry.