



wwPDB EM Validation Summary Report ⓘ

May 13, 2024 – 09:58 pm BST

PDB ID : 8RPY
EMDB ID : EMD-19426
Title : Escherichia coli 50S subunit in complex with the antimicrobial peptide Api137
Authors : Lauer, S.; Nikolay, R.; Spahn, C.
Deposited on : 2024-01-17
Resolution : 2.64 Å (reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:


EMDB validation analysis : 0.0.1.dev92
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36.2

1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.64 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




























Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826
RNA backbone	4643	859

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	0	56	96% (Green), 4% (Yellow), 0% (Orange), 0% (Red), 0% (Grey)
2	1	50	92% (Green), 8% (Yellow), 0% (Orange), 0% (Red), 0% (Grey)
3	2	46	87% (Green), 13% (Yellow), 0% (Orange), 0% (Red), 0% (Grey)
4	3	64	91% (Green), 9% (Yellow), 0% (Orange), 0% (Red), 0% (Grey)
5	4	38	71% (Green), 26% (Yellow), 3% (Orange), 0% (Red), 0% (Grey)
6	6	66	52% (Green), 14% (Yellow), 35% (Grey), 5% (Red), 0% (Orange)
7	A	2903	79% (Green), 18% (Yellow), 3% (Orange), 0% (Red), 0% (Grey)


Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
8	B	120	 80% 17%
9	C	271	 82% 18%
10	D	209	 86% 14%
11	E	201	 79% 21%
12	F	177	 72% 27%
13	G	176	 92% 8%
14	H	149	 68% 90% 10%
15	J	142	 85% 15%
16	K	122	 90% 10%
17	L	143	 78% 22%
18	M	136	 88% 12%
19	N	120	 75% 25%
20	O	116	 83% 17%
21	P	114	 82% 18%
22	Q	117	 89% 11%
23	R	103	 92% 7%
24	S	110	 81% 18%
25	T	93	 80% 20%
26	U	102	 79% 21%
27	V	94	 68% 32%
28	W	75	 84% 16%
29	X	77	 90% 10%
30	Y	63	 5% 92% 8%
31	Z	58	 84% 16%
32	y	17	 24% 94% 6%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
32	z	17	 <p>A horizontal bar chart representing the quality of chain. The bar is divided into two segments: a red segment on the left labeled '24%' and a green segment on the right labeled '100%'. The green segment extends to the right edge of the chart area.</p>

2 Entry composition [i](#)

There are 32 unique types of molecules in this entry. The entry contains 90576 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Large ribosomal subunit protein bL32.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	0	56	444	269	94	80	1	0	0

- Molecule 2 is a protein called Large ribosomal subunit protein bL33.

Mol	Chain	Residues	Atoms				AltConf	Trace
			Total	C	N	O		
2	1	50	409	263	75	71	0	0

- Molecule 3 is a protein called Large ribosomal subunit protein bL34.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
3	2	46	377	228	90	57	2	0	0

- Molecule 4 is a protein called Large ribosomal subunit protein bL35.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
4	3	64	504	323	105	74	2	0	0

- Molecule 5 is a protein called Large ribosomal subunit protein bL36A.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
5	4	38	302	185	65	48	4	0	0

- Molecule 6 is a protein called Large ribosomal subunit protein bL31.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
6	6	43	333	206	59	62	6	0	0

- Molecule 7 is a RNA chain called 23S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
7	A	2900	62282	27783	11462	20136	2901	1	0

- Molecule 8 is a RNA chain called 5S ribosomal RNA.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	P		
8	B	120	2572	1145	471	836	120	0	0

- Molecule 9 is a protein called Large ribosomal subunit protein uL2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
9	C	271	2082	1288	423	364	7	0	0

- Molecule 10 is a protein called Large ribosomal subunit protein uL3.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
10	D	209	1565	979	288	294	4	0	0

- Molecule 11 is a protein called Large ribosomal subunit protein uL4.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
11	E	201	1552	974	283	290	5	0	0

- Molecule 12 is a protein called Large ribosomal subunit protein uL5.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
12	F	177	1410	899	249	256	6	0	0

- Molecule 13 is a protein called Large ribosomal subunit protein uL6.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
13	G	176	1323	832	243	246	2	0	0

- Molecule 14 is a protein called Large ribosomal subunit protein bL9.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	H	149	Total	C	N	O	S	0	0
			1111	699	197	214	1		

- Molecule 15 is a protein called Large ribosomal subunit protein uL13.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	J	142	Total	C	N	O	S	0	0
			1129	714	212	199	4		

- Molecule 16 is a protein called Large ribosomal subunit protein uL14.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	K	122	Total	C	N	O	S	0	0
			938	587	180	165	6		

- Molecule 17 is a protein called Large ribosomal subunit protein uL15.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	L	143	Total	C	N	O	S	0	0
			1045	649	206	189	1		

- Molecule 18 is a protein called Large ribosomal subunit protein uL16.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	M	136	Total	C	N	O	S	0	0
			1074	686	205	177	6		

- Molecule 19 is a protein called Large ribosomal subunit protein bL17.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	N	120	Total	C	N	O	S	0	0
			960	593	196	166	5		

- Molecule 20 is a protein called Large ribosomal subunit protein uL18.

Mol	Chain	Residues	Atoms				AltConf	Trace
20	O	116	Total	C	N	O	0	0
			892	552	178	162		

- Molecule 21 is a protein called Large ribosomal subunit protein bL19.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
21	P	114	917	574	179	163	1	0	0

- Molecule 22 is a protein called Large ribosomal subunit protein bL20.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
22	Q	117	947	604	192	151		0	0

- Molecule 23 is a protein called Large ribosomal subunit protein bL21.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
23	R	103	816	516	153	145	2	0	0

- Molecule 24 is a protein called Large ribosomal subunit protein uL22.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
24	S	110	868	538	170	157	3	1	0

- Molecule 25 is a protein called Large ribosomal subunit protein uL23.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
25	T	93	738	466	139	131	2	0	0

- Molecule 26 is a protein called Large ribosomal subunit protein uL24.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
26	U	102	779	492	146	141		0	0

- Molecule 27 is a protein called Large ribosomal subunit protein bL25.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
27	V	94	753	479	137	134	3	0	0

- Molecule 28 is a protein called Large ribosomal subunit protein bL27.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	W	75	Total	C	N	O	S	0	0
			575	356	116	102	1		

- Molecule 29 is a protein called Large ribosomal subunit protein bL28.

Mol	Chain	Residues	Atoms					AltConf	Trace
29	X	77	Total	C	N	O	S	0	0
			625	388	129	106	2		

- Molecule 30 is a protein called Large ribosomal subunit protein uL29.

Mol	Chain	Residues	Atoms					AltConf	Trace
30	Y	63	Total	C	N	O	S	0	0
			509	313	99	95	2		

- Molecule 31 is a protein called Large ribosomal subunit protein uL30.

Mol	Chain	Residues	Atoms					AltConf	Trace
31	Z	58	Total	C	N	O	S	0	0
			449	281	87	79	2		


- Molecule 32 is a protein called Apidaecins type 137.

Mol	Chain	Residues	Atoms				AltConf	Trace
32	y	17	Total	C	N	O	0	0
			148	94	33	21		
32	z	17	Total	C	N	O	0	0
			148	94	33	21		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Large ribosomal subunit protein bL32




- Molecule 2: Large ribosomal subunit protein bL33




- Molecule 3: Large ribosomal subunit protein bL34




- Molecule 4: Large ribosomal subunit protein bL35





- Molecule 5: Large ribosomal subunit protein bL36A




- Molecule 6: Large ribosomal subunit protein bL31





• Molecule 7: 23S ribosomal RNA







• Molecule 8: 5S ribosomal RNA




• Molecule 9: Large ribosomal subunit protein uL2




• Molecule 10: Large ribosomal subunit protein uL3




- Molecule 11: Large ribosomal subunit protein uL4




- Molecule 12: Large ribosomal subunit protein uL5




- Molecule 13: Large ribosomal subunit protein uL6




- Molecule 14: Large ribosomal subunit protein bL9



- Molecule 15: Large ribosomal subunit protein uL13





- Molecule 16: Large ribosomal subunit protein uL14

Chain K:  90% 10%




- Molecule 17: Large ribosomal subunit protein uL15

Chain L:  78% 22%





- Molecule 18: Large ribosomal subunit protein uL16

Chain M:  88% 12%





- Molecule 19: Large ribosomal subunit protein bL17

Chain N:  75% 25%




- Molecule 20: Large ribosomal subunit protein uL18

Chain O:  83% 17%




- Molecule 21: Large ribosomal subunit protein bL19

Chain P:  82% 18%





- Molecule 22: Large ribosomal subunit protein bL20

Chain Q:  89% 11%





- Molecule 23: Large ribosomal subunit protein bL21

Chain R:  92% 7%




- Molecule 24: Large ribosomal subunit protein uL22

Chain S:  81% 18%




- Molecule 25: Large ribosomal subunit protein uL23

Chain T:  80% 20%



- Molecule 26: Large ribosomal subunit protein uL24

Chain U:  79% 21%




- Molecule 27: Large ribosomal subunit protein bL25

Chain V:  68% 32%




- Molecule 28: Large ribosomal subunit protein bL27

Chain W:  84% 16%




- Molecule 29: Large ribosomal subunit protein bL28



Chain X:  90% 10%





- Molecule 30: Large ribosomal subunit protein uL29




- Molecule 31: Large ribosomal subunit protein uL30



- Molecule 32: Apidaecins type 137



- Molecule 32: Apidaecins type 137



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	424506	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	24.6	Depositor
Minimum defocus (nm)	500	Depositor
Maximum defocus (nm)	2000	Depositor
Magnification	Not provided	
Image detector	GATAN K3 BIOQUANTUM (6k x 4k)	Depositor
Maximum map value	1.538	Depositor
Minimum map value	-0.573	Depositor
Average map value	-0.002	Depositor
Map value standard deviation	0.067	Depositor
Recommended contour level	0.124	Depositor
Map size (\AA)	353.99997, 353.99997, 353.99997	wwPDB
Map dimensions	300, 300, 300	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (\AA)	1.18, 1.18, 1.18	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	0	0.25	0/450	0.54	0/599
2	1	0.26	0/416	0.49	0/554
3	2	0.26	0/380	0.64	0/498
4	3	0.24	0/513	0.56	0/676
5	4	0.27	0/303	0.58	0/397
6	6	0.26	0/339	0.49	0/454
7	A	0.30	1/69756 (0.0%)	0.66	4/108822 (0.0%)
8	B	0.31	1/2876 (0.0%)	0.65	0/4483
9	C	0.27	0/2121	0.56	0/2852
10	D	0.27	0/1586	0.52	0/2134
11	E	0.26	0/1571	0.52	0/2113
12	F	0.25	0/1434	0.53	0/1926
13	G	0.26	0/1343	0.51	0/1816
14	H	0.24	0/1122	0.48	0/1515
15	J	0.26	0/1152	0.49	0/1551
16	K	0.26	0/947	0.55	0/1268
17	L	0.27	0/1054	0.58	0/1403
18	M	0.26	0/1093	0.55	0/1460
19	N	0.26	0/973	0.58	0/1301
20	O	0.25	0/902	0.53	0/1209
21	P	0.27	0/929	0.56	0/1242
22	Q	0.28	0/960	0.53	0/1278
23	R	0.27	0/829	0.53	0/1107
24	S	0.25	0/875	0.54	0/1170
25	T	0.26	0/744	0.53	0/994
26	U	0.27	0/787	0.52	0/1051
27	V	0.28	0/766	0.54	0/1025
28	W	0.28	0/582	0.55	0/769
29	X	0.26	0/635	0.58	0/848
30	Y	0.25	0/510	0.52	0/677
31	Z	0.31	0/453	0.55	0/605
32	y	0.48	0/155	0.66	0/212
32	z	0.24	0/155	0.59	0/212
All	All	0.29	2/98711 (0.0%)	0.63	4/148221 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
8	B	1	U	OP3-P	-10.61	1.48	1.61
7	A	1	G	OP3-P	-10.60	1.48	1.61

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
7	A	1913	A	OP1-P-O3'	-10.79	81.45	105.20
7	A	1913	A	OP2-P-O3'	-10.25	82.66	105.20
7	A	1914	C	OP1-P-OP2	7.09	130.24	119.60
7	A	2321	U	C2-N1-C1'	5.46	124.25	117.70

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	0	444	0	461	1	0
2	1	409	0	440	4	0
3	2	377	0	418	5	0
4	3	504	0	574	7	0
5	4	302	0	343	10	0
6	6	333	0	332	8	0
7	A	62282	0	31324	239	0
8	B	2572	0	1302	13	0
9	C	2082	0	2157	31	0
10	D	1565	0	1616	28	0
11	E	1552	0	1619	28	0
12	F	1410	0	1447	36	0
13	G	1323	0	1374	10	0
14	H	1111	0	1148	10	0
15	J	1129	0	1162	14	0
16	K	938	0	1012	12	0
17	L	1045	0	1117	35	0
18	M	1074	0	1157	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
19	N	960	0	1000	20	0
20	O	892	0	923	16	0
21	P	917	0	965	18	0
22	Q	947	0	1022	11	0
23	R	816	0	839	7	0
24	S	868	0	934	14	0
25	T	738	0	807	11	0
26	U	779	0	834	15	0
27	V	753	0	780	22	0
28	W	575	0	592	8	0
29	X	625	0	655	5	0
30	Y	509	0	543	3	0
31	Z	449	0	491	6	0
32	y	148	0	152	0	0
32	z	148	0	152	0	0
All	All	90576	0	59692	543	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 543 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
27:V:62:THR:OG1	27:V:69:GLU:OE2	1.83	0.97
31:Z:26:LEU:O	31:Z:37:ARG:NH1	2.02	0.92
7:A:2326:C:O2'	7:A:2327:A:OP1	1.89	0.91
14:H:113:SER:O	14:H:116:ARG:NH1	2.03	0.91
30:Y:34:SER:OG	30:Y:36:GLN:OE1	1.87	0.90

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	0	54/56 (96%)	53 (98%)	1 (2%)	0	100	100
2	1	48/50 (96%)	47 (98%)	1 (2%)	0	100	100
3	2	44/46 (96%)	44 (100%)	0	0	100	100
4	3	62/64 (97%)	59 (95%)	3 (5%)	0	100	100
5	4	36/38 (95%)	35 (97%)	1 (3%)	0	100	100
6	6	41/66 (62%)	39 (95%)	2 (5%)	0	100	100
9	C	269/271 (99%)	257 (96%)	12 (4%)	0	100	100
10	D	207/209 (99%)	198 (96%)	9 (4%)	0	100	100
11	E	199/201 (99%)	191 (96%)	7 (4%)	1 (0%)	29	43
12	F	175/177 (99%)	168 (96%)	7 (4%)	0	100	100
13	G	174/176 (99%)	169 (97%)	5 (3%)	0	100	100
14	H	147/149 (99%)	141 (96%)	6 (4%)	0	100	100
15	J	140/142 (99%)	135 (96%)	5 (4%)	0	100	100
16	K	120/122 (98%)	117 (98%)	3 (2%)	0	100	100
17	L	141/143 (99%)	127 (90%)	14 (10%)	0	100	100
18	M	134/136 (98%)	129 (96%)	5 (4%)	0	100	100
19	N	118/120 (98%)	110 (93%)	8 (7%)	0	100	100
20	O	114/116 (98%)	112 (98%)	2 (2%)	0	100	100
21	P	112/114 (98%)	108 (96%)	4 (4%)	0	100	100
22	Q	115/117 (98%)	112 (97%)	3 (3%)	0	100	100
23	R	101/103 (98%)	95 (94%)	5 (5%)	1 (1%)	15	22
24	S	109/110 (99%)	105 (96%)	4 (4%)	0	100	100
25	T	91/93 (98%)	85 (93%)	6 (7%)	0	100	100
26	U	100/102 (98%)	89 (89%)	11 (11%)	0	100	100
27	V	92/94 (98%)	86 (94%)	6 (6%)	0	100	100
28	W	73/75 (97%)	70 (96%)	3 (4%)	0	100	100
29	X	75/77 (97%)	74 (99%)	1 (1%)	0	100	100
30	Y	61/63 (97%)	59 (97%)	2 (3%)	0	100	100
31	Z	56/58 (97%)	54 (96%)	2 (4%)	0	100	100
32	y	15/17 (88%)	11 (73%)	4 (27%)	0	100	100
32	z	15/17 (88%)	14 (93%)	1 (7%)	0	100	100
All	All	3238/3322 (98%)	3093 (96%)	143 (4%)	2 (0%)	54	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
23	R	54	VAL
11	E	83	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	0	47/47 (100%)	47 (100%)	0	100	100
2	1	45/45 (100%)	45 (100%)	0	100	100
3	2	38/38 (100%)	38 (100%)	0	100	100
4	3	51/51 (100%)	51 (100%)	0	100	100
5	4	34/34 (100%)	33 (97%)	1 (3%)	42	60
6	6	40/59 (68%)	40 (100%)	0	100	100
9	C	216/216 (100%)	216 (100%)	0	100	100
10	D	164/164 (100%)	164 (100%)	0	100	100
11	E	165/165 (100%)	164 (99%)	1 (1%)	86	93
12	F	148/148 (100%)	147 (99%)	1 (1%)	84	91
13	G	137/137 (100%)	137 (100%)	0	100	100
14	H	114/114 (100%)	114 (100%)	0	100	100
15	J	116/116 (100%)	116 (100%)	0	100	100
16	K	103/103 (100%)	103 (100%)	0	100	100
17	L	102/102 (100%)	102 (100%)	0	100	100
18	M	109/109 (100%)	109 (100%)	0	100	100
19	N	100/100 (100%)	99 (99%)	1 (1%)	76	86
20	O	86/86 (100%)	86 (100%)	0	100	100
21	P	99/99 (100%)	99 (100%)	0	100	100
22	Q	89/89 (100%)	89 (100%)	0	100	100
23	R	84/84 (100%)	84 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
24	S	94/93 (101%)	93 (99%)	1 (1%)	73	85
25	T	80/80 (100%)	80 (100%)	0	100	100
26	U	83/83 (100%)	83 (100%)	0	100	100
27	V	78/78 (100%)	78 (100%)	0	100	100
28	W	57/57 (100%)	57 (100%)	0	100	100
29	X	67/67 (100%)	67 (100%)	0	100	100
30	Y	55/55 (100%)	54 (98%)	1 (2%)	59	75
31	Z	48/48 (100%)	48 (100%)	0	100	100
32	y	17/17 (100%)	16 (94%)	1 (6%)	19	30
32	z	17/17 (100%)	17 (100%)	0	100	100
All	All	2683/2701 (99%)	2676 (100%)	7 (0%)	92	96

5 of 7 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
19	N	2	ARG
24	S	110	ARG
32	y	17	ARG
30	Y	7	ARG
12	F	29	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
10	D	173	GLN
13	G	103	ASN
17	L	99	ASN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
7	A	2897/2903 (99%)	339 (11%)	14 (0%)
8	B	119/120 (99%)	11 (9%)	3 (2%)
All	All	3016/3023 (99%)	350 (11%)	17 (0%)

5 of 350 RNA backbone outliers are listed below:

Mol	Chain	Res	Type
7	A	10	A
7	A	34	U
7	A	35	G
7	A	46	G
7	A	51	G

5 of 17 RNA pucker outliers are listed below:

Mol	Chain	Res	Type
8	B	44	G
8	B	66	A
7	A	784	G
7	A	1020	A
7	A	1022	G

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-19426. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 150




Y Index: 150




Z Index: 150


6.2.2 Raw map



X Index: 150



Y Index: 150




Z Index: 150


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 150




Y Index: 159




Z Index: 165


6.3.2 Raw map



X Index: 150



Y Index: 159




Z Index: 165


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.124. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_19426_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 884 nm³; this corresponds to an approximate mass of 798 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum i




*Reported resolution corresponds to spatial frequency of 0.379 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.379 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.64	-	-
Author-provided FSC curve	2.64	2.93	2.70
Unmasked-calculated*	3.12	3.86	3.16

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.12 differs from the reported value 2.64 by more than 10 %

9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-19426 and PDB model 8RPY. Per-residue inclusion information can be found in section 3 on page 10.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 0.124 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)


This section was not generated.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.124).



































9.4 Atom inclusion [i](#)



At the recommended contour level, 96% of all backbone atoms, 93% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (0.124) and Q-score for the entire model and for each chain.

Chain	Atom inclusion
All	 0.9310
0	 0.9320
1	 0.9180
2	 0.9440
3	 0.9450
4	 0.9310
6	 0.7490
A	 0.9500
B	 0.9810
C	 0.9470
D	 0.9260
E	 0.9080
F	 0.8020
G	 0.8710
H	 0.3200
J	 0.9410
K	 0.8630
L	 0.8930
M	 0.9200
N	 0.9410
O	 0.9040
P	 0.8840
Q	 0.9570
R	 0.8960
S	 0.9100
T	 0.9040
U	 0.9270
V	 0.9010
W	 0.9320
X	 0.9400
Y	 0.8890
Z	 0.9150
y	 0.5610
z	 0.5470

