



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 5, 2023 – 04:13 AM EDT

PDB ID : 3RQF
Title : Cerebral cavernous malformation 3 (CCM3) in complex with paxillin LD2
Authors : Li, X.; Zhang, R.; Boggon, T.J.
Deposited on : 2011-04-28
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.35
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	214	 5% (poor fit), 77% (0-1 outliers), 14% (2 outliers), 8% (3+ outliers), 8% (not modelled)
1	B	214	 11% (poor fit), 82% (0-1 outliers), 8% (2 outliers), 8% (3+ outliers), 8% (not modelled)
1	C	214	 10% (poor fit), 77% (0-1 outliers), 15% (2 outliers), 8% (3+ outliers), 8% (not modelled)
1	D	214	 8% (poor fit), 80% (0-1 outliers), 8% (2 outliers), 10% (3+ outliers), 8% (not modelled)
2	E	13	 85% (poor fit), 85% (0-1 outliers), 15% (2 outliers), 8% (3+ outliers), 8% (not modelled)

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 6474 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Programmed cell death protein 10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	197	Total 1611	C 1030	N 273	O 302	S 6	0	0	0
1	B	197	Total 1605	C 1024	N 271	O 304	S 6	0	0	0
1	C	197	Total 1598	C 1017	N 271	O 301	S 9	0	0	0
1	D	192	Total 1565	C 998	N 267	O 295	S 5	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q9BUL8
A	0	HIS	-	expression tag	UNP Q9BUL8
B	-1	GLY	-	expression tag	UNP Q9BUL8
B	0	HIS	-	expression tag	UNP Q9BUL8
C	-1	GLY	-	expression tag	UNP Q9BUL8
C	0	HIS	-	expression tag	UNP Q9BUL8
D	-1	GLY	-	expression tag	UNP Q9BUL8
D	0	HIS	-	expression tag	UNP Q9BUL8

- Molecule 2 is a protein called Paxillin LD2 peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	E	11	Total 91	C 57	N 15	O 19	0	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total 1	O 1	0	0



Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total O 2 2	0	0
3	D	1	Total O 1 1	0	0



- Molecule 2: Paxillin LD2 peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.86Å 115.47Å 124.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	42.52 – 2.70 42.52 – 2.70	Depositor EDS
% Data completeness (in resolution range)	99.7 (42.52-2.70) 99.7 (42.52-2.70)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.22 (at 2.69Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.237 , 0.291 0.233 , 0.284	Depositor DCC
R_{free} test set	1292 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	67.3	Xtrriage
Anisotropy	0.106	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 56.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6474	wwPDB-VP
Average B, all atoms (Å ²)	85.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/1636	0.47	0/2200
1	B	0.34	0/1628	0.47	0/2189
1	C	0.34	0/1619	0.49	0/2170
1	D	0.34	0/1587	0.47	0/2132
2	E	0.32	0/90	0.46	0/120
All	All	0.34	0/6560	0.47	0/8811

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1611	0	1655	19	0
1	B	1605	0	1646	10	0
1	C	1598	0	1643	20	0
1	D	1565	0	1607	9	0
2	E	91	0	94	0	0
3	A	1	0	0	0	0
3	B	2	0	0	0	0
3	D	1	0	0	0	0
All	All	6474	0	6645	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 4.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:197:LEU:O	1:D:201:THR:HG22	1.93	0.69
1:C:21:PRO:HG2	1:C:60:GLN:HB2	1.77	0.66
1:D:21:PRO:HG2	1:D:60:GLN:HE21	1.62	0.64
1:A:85:ALA:HB2	1:A:115:SER:HB3	1.79	0.64
1:B:21:PRO:HG2	1:B:60:GLN:HB2	1.82	0.61
1:A:157:ARG:HA	1:A:160:LEU:HD23	1.85	0.57
1:C:10:ASN:O	1:C:14:THR:HG23	2.04	0.57
1:D:59:THR:O	1:D:63:ILE:HG12	2.04	0.57
1:A:131:ILE:HG12	1:A:178:LEU:HD21	1.86	0.57
1:C:109:ALA:O	1:C:113:ILE:HG12	2.04	0.57
1:C:27:TYR:OH	1:C:49:ILE:HD11	2.05	0.56
1:C:1:MET:HG3	1:C:34:GLU:HG3	1.89	0.53
1:A:32:GLU:HA	1:C:8:MET:HG2	1.90	0.53
1:A:197:LEU:O	1:A:201:THR:HG22	2.09	0.52
1:D:82:ARG:NH2	1:D:119:ASP:OD1	2.42	0.52
1:A:37:ASN:HD22	1:A:40:ALA:H	1.59	0.51
1:B:131:ILE:HD12	1:B:178:LEU:HG	1.93	0.51
1:B:34:GLU:OE2	1:B:45:ARG:NH2	2.41	0.51
1:C:51:ALA:HB1	1:C:58:LEU:CD2	2.42	0.49
1:C:63:ILE:HG21	1:D:73:GLU:HG3	1.94	0.49
1:C:106:LYS:HD3	1:C:140:GLU:HB3	1.94	0.49
1:A:141:LEU:O	1:A:144:THR:HG22	2.13	0.48
1:C:14:THR:CG2	1:C:53:LYS:HA	2.44	0.48
1:C:33:LEU:HD11	1:D:58:LEU:HD11	1.94	0.47
1:D:78:GLU:O	1:D:82:ARG:HG3	2.15	0.47
1:A:141:LEU:HD23	1:A:167:PHE:CZ	2.50	0.47
1:C:180:THR:HG22	1:C:184:ASP:OD2	2.15	0.46
1:A:176:ASP:HA	1:A:179:LYS:HD2	1.98	0.46
1:A:85:ALA:HB2	1:A:115:SER:CB	2.45	0.46
1:B:127:PHE:O	1:B:131:ILE:HG12	2.15	0.45
1:C:20:MET:N	1:C:21:PRO:CD	2.79	0.45
1:A:85:ALA:O	1:A:86:ASP:C	2.56	0.44
1:A:37:ASN:ND2	1:A:40:ALA:H	2.16	0.43
1:B:100:PHE:HE1	1:B:208:PHE:HB3	1.83	0.43
1:A:184:ASP:HB3	1:A:186:LYS:HG3	1.99	0.43
1:D:157:ARG:HG3	1:D:159:ALA:H	1.84	0.43
1:A:74:VAL:HG22	1:A:188:ILE:HG23	2.00	0.43
1:B:51:ALA:HB1	1:B:58:LEU:CD2	2.49	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:14:THR:HG21	1:C:53:LYS:HA	2.00	0.43
1:C:51:ALA:HB1	1:C:58:LEU:HD23	1.99	0.42
1:A:17:MET:C	1:A:19:SER:H	2.22	0.42
1:B:89:GLU:O	1:B:92:MET:HG2	2.19	0.42
1:B:20:MET:HE2	1:B:20:MET:HB2	1.92	0.42
1:A:141:LEU:HA	1:A:144:THR:HG22	2.01	0.42
1:C:3:MET:CG	1:C:7:GLU:HB3	2.49	0.42
1:A:65:LYS:HB3	1:B:29:VAL:HG22	2.01	0.41
1:C:1:MET:HE1	1:C:45:ARG:HB2	2.03	0.41
1:C:45:ARG:O	1:C:49:ILE:HG12	2.20	0.41
1:A:39:SER:HA	1:A:42:GLN:HE21	1.85	0.41
1:A:134:ILE:HG21	1:A:174:PHE:HE2	1.85	0.41
1:B:123:ASP:OD1	1:B:126:ARG:HB2	2.21	0.41
1:C:1:MET:HE3	1:C:41:ALA:O	2.22	0.40
1:C:97:GLU:HB2	1:C:100:PHE:HD2	1.86	0.40
1:D:116:LYS:O	1:D:120:GLU:HG2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	195/214 (91%)	185 (95%)	9 (5%)	1 (0%)	29	54
1	B	193/214 (90%)	190 (98%)	3 (2%)	0	100	100
1	C	191/214 (89%)	188 (98%)	3 (2%)	0	100	100
1	D	188/214 (88%)	180 (96%)	7 (4%)	1 (0%)	29	54
2	E	9/13 (69%)	9 (100%)	0	0	100	100
All	All	776/869 (89%)	752 (97%)	22 (3%)	2 (0%)	41	66

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	87	ASP
1	D	157	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/194 (92%)	172 (96%)	7 (4%)	32	61
1	B	179/194 (92%)	172 (96%)	7 (4%)	32	61
1	C	178/194 (92%)	171 (96%)	7 (4%)	32	61
1	D	174/194 (90%)	168 (97%)	6 (3%)	37	66
2	E	11/13 (85%)	11 (100%)	0	100	100
All	All	721/789 (91%)	694 (96%)	27 (4%)	34	63

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	17	MET
1	A	71	SER
1	A	83	MET
1	A	124	ARG
1	A	173	SER
1	A	188	ILE
1	A	201	THR
1	B	20	MET
1	B	38	LEU
1	B	88	VAL
1	B	92	MET
1	B	97	GLU
1	B	113	ILE
1	B	178	LEU
1	C	31	ASN
1	C	55	ASN
1	C	60	GLN
1	C	151	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	161	GLU
1	C	178	LEU
1	C	207	THR
1	D	37	ASN
1	D	87	ASP
1	D	112	GLN
1	D	113	ILE
1	D	201	THR
1	D	207	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (19) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	60	GLN
1	A	147	ASN
1	B	55	ASN
1	B	60	GLN
1	B	75	ASN
1	B	163	GLN
1	C	55	ASN
1	C	60	GLN
1	C	146	ASN
1	C	163	GLN
1	C	206	GLN
1	D	37	ASN
1	D	42	GLN
1	D	60	GLN
1	D	112	GLN
1	D	155	GLN
1	D	189	ASN
1	D	195	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	197/214 (92%)	0.41	10 (5%) 28 26	39, 71, 107, 133	0
1	B	197/214 (92%)	0.63	23 (11%) 4 3	40, 81, 156, 201	0
1	C	197/214 (92%)	0.62	22 (11%) 5 4	42, 75, 155, 193	0
1	D	192/214 (89%)	0.56	17 (8%) 9 7	42, 79, 159, 207	0
2	E	11/13 (84%)	4.59	11 (100%) 0 0	141, 142, 144, 144	0
All	All	794/869 (91%)	0.61	83 (10%) 6 4	39, 77, 150, 207	0

All (83) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	143	SER	12.3
1	C	149	PHE	9.7
2	E	145	LEU	7.0
1	B	158	ARG	6.9
1	D	155	GLN	6.3
1	B	208	PHE	5.8
2	E	144	GLU	5.1
1	C	145	VAL	5.0
1	A	91	TYR	4.8
1	B	99	GLU	4.6
1	C	148	VAL	4.5
2	E	151	GLU	4.5
1	C	207	THR	4.4
1	B	11	GLU	4.3
1	D	10	ASN	4.2
1	D	154	TYR	4.2
1	A	92	MET	4.2
1	B	210	THR	4.1
1	C	100	PHE	4.0
2	E	147	ARG	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	13	GLU	3.9
2	E	148	LEU	3.8
1	A	16	SER	3.8
1	D	87	ASP	3.7
1	C	206	GLN	3.7
1	C	203	LEU	3.7
2	E	146	ASP	3.6
1	B	96	PRO	3.5
1	C	147	ASN	3.5
1	B	124	ARG	3.5
1	C	143	ASP	3.4
1	C	209	LYS	3.3
1	C	142	LEU	3.3
1	B	128	LEU	3.2
1	B	100	PHE	3.2
1	D	108	ARG	3.2
1	B	204	ILE	3.2
1	D	11	GLU	3.1
1	D	206	GLN	3.1
2	E	150	LEU	3.0
1	A	89	GLU	3.0
1	C	205	LEU	3.0
1	D	186	LYS	2.9
1	C	97	GLU	2.9
2	E	149	LEU	2.9
1	C	141	LEU	2.8
1	A	208	PHE	2.8
1	A	210	THR	2.8
1	C	151	LYS	2.8
1	B	168	VAL	2.7
1	C	138	ILE	2.7
1	B	164	LYS	2.7
1	B	12	ALA	2.7
1	B	92	MET	2.7
1	C	35	ARG	2.7
1	B	103	LEU	2.6
1	C	124	ARG	2.6
1	C	87	ASP	2.6
1	B	150	LYS	2.5
1	D	201	THR	2.5
1	B	122	ASN	2.5
1	B	85	ALA	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	124	ARG	2.5
2	E	152	LEU	2.5
1	B	98	PRO	2.4
1	A	17	MET	2.4
1	B	102	ASP	2.3
1	A	212	ALA	2.3
1	D	12	ALA	2.3
1	C	150	LYS	2.3
1	C	103	LEU	2.2
1	A	35	ARG	2.2
1	D	207	THR	2.2
1	B	125	VAL	2.2
1	B	149	PHE	2.2
1	C	152	TYR	2.1
1	A	36	VAL	2.1
1	D	128	LEU	2.1
1	D	129	GLN	2.1
1	D	151	LYS	2.1
1	D	152	TYR	2.1
1	B	121	ILE	2.1
2	E	153	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.