



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 22, 2020 – 11:58 AM BST

PDB ID : 4RQO  
Title : Crystal structure of L-Serine Dehydratase from Legionella pneumophila  
Authors : Thoden, J.B.; Holden, H.M.; Grant, G.A.  
Deposited on : 2014-11-04  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.13.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.13.1


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)
RSRZ outliers	127900	1356 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	476	<div style="display: flex; align-items: center;"> <div style="width: 9%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 83%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 7%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">9%      83%      9%      • 7%</p>
1	B	476	<div style="display: flex; align-items: center;"> <div style="width: 4%; height: 10px; background-color: red; margin-right: 5px;"></div> <div style="width: 83%; height: 10px; background-color: green; margin-right: 5px;"></div> <div style="width: 9%; height: 10px; background-color: yellow; margin-right: 5px;"></div> <div style="width: 6%; height: 10px; background-color: grey;"></div> </div> <p style="font-size: small; margin-top: 5px;">4%      83%      9%      • 6%</p>

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 6976 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called L-serine dehydratase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	N	O	S	Se			
1	A	444	3359	2120	574	639	8	18	0	0	0
1	B	448	3395	2142	580	646	8	19	0	0	0

There are 36 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MSE	-	EXPRESSION TAG	UNP R9SE63
A	-16	GLY	-	EXPRESSION TAG	UNP R9SE63
A	-15	SER	-	EXPRESSION TAG	UNP R9SE63
A	-14	SER	-	EXPRESSION TAG	UNP R9SE63
A	-13	HIS	-	EXPRESSION TAG	UNP R9SE63
A	-12	HIS	-	EXPRESSION TAG	UNP R9SE63
A	-11	HIS	-	EXPRESSION TAG	UNP R9SE63
A	-10	HIS	-	EXPRESSION TAG	UNP R9SE63
A	-9	HIS	-	EXPRESSION TAG	UNP R9SE63
A	-8	HIS	-	EXPRESSION TAG	UNP R9SE63
A	-7	GLU	-	EXPRESSION TAG	UNP R9SE63
A	-6	ASN	-	EXPRESSION TAG	UNP R9SE63
A	-5	LEU	-	EXPRESSION TAG	UNP R9SE63
A	-4	TYR	-	EXPRESSION TAG	UNP R9SE63
A	-3	PHE	-	EXPRESSION TAG	UNP R9SE63
A	-2	GLN	-	EXPRESSION TAG	UNP R9SE63
A	-1	GLY	-	EXPRESSION TAG	UNP R9SE63
A	0	SER	-	EXPRESSION TAG	UNP R9SE63
B	-17	MSE	-	EXPRESSION TAG	UNP R9SE63
B	-16	GLY	-	EXPRESSION TAG	UNP R9SE63
B	-15	SER	-	EXPRESSION TAG	UNP R9SE63
B	-14	SER	-	EXPRESSION TAG	UNP R9SE63
B	-13	HIS	-	EXPRESSION TAG	UNP R9SE63
B	-12	HIS	-	EXPRESSION TAG	UNP R9SE63
B	-11	HIS	-	EXPRESSION TAG	UNP R9SE63

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-10	HIS	-	EXPRESSION TAG	UNP R9SE63
B	-9	HIS	-	EXPRESSION TAG	UNP R9SE63
B	-8	HIS	-	EXPRESSION TAG	UNP R9SE63
B	-7	GLU	-	EXPRESSION TAG	UNP R9SE63
B	-6	ASN	-	EXPRESSION TAG	UNP R9SE63
B	-5	LEU	-	EXPRESSION TAG	UNP R9SE63
B	-4	TYR	-	EXPRESSION TAG	UNP R9SE63
B	-3	PHE	-	EXPRESSION TAG	UNP R9SE63
B	-2	GLN	-	EXPRESSION TAG	UNP R9SE63
B	-1	GLY	-	EXPRESSION TAG	UNP R9SE63
B	0	SER	-	EXPRESSION TAG	UNP R9SE63

- Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe<sub>4</sub>S<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Fe S 8 4 4	0	0
2	B	1	Total Fe S 8 4 4	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	2	Total Cl 2 2	0	0


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	95	Total 95	O 95	0	0
4	B	109	Total 109	O 109	0	0


### 3 Residue-property plots i

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: L-serine dehydratase



- Molecule 1: L-serine dehydratase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.39Å 81.39Å 267.52Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.45 – 2.25 29.43 – 2.25	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.45-2.25) 99.7 (29.43-2.25)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	0.10	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.62 (at 2.24Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.198 , 0.258 0.210 , 0.262	Depositor DCC
$R_{free}$ test set	2538 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.7	Xtrriage
Anisotropy	0.081	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 31.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.033 for -h,-k,l	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	6976	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SF4, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.70	0/3397	0.79	2/4558 (0.0%)
1	B	0.81	0/3433	0.90	6/4605 (0.1%)
All	All	0.76	0/6830	0.85	8/9163 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	240	ARG	NE-CZ-NH1	8.75	124.67	120.30
1	B	240	ARG	NE-CZ-NH2	-8.21	116.19	120.30
1	B	399	ARG	NE-CZ-NH1	6.29	123.44	120.30
1	A	79	VAL	CB-CA-C	-5.99	100.01	111.40
1	A	128	MSE	CB-CA-C	-5.34	99.73	110.40
1	B	218	MSE	CG-SE-CE	5.32	110.61	98.90
1	B	247	ARG	NE-CZ-NH2	-5.30	117.65	120.30
1	B	88	MSE	CG-SE-CE	5.02	109.95	98.90

There are no chirality outliers.

All (1) planarity outliers are listed below:



Mol	Chain	Res	Type	Group
1	A	167	ASN	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3359	0	3377	23	0
1	B	3395	0	3419	24	0
2	A	8	0	0	0	0
2	B	8	0	0	0	0
3	B	2	0	0	0	0
4	A	95	0	0	1	0
4	B	109	0	0	0	0
All	All	6976	0	6796	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:180:LEU:HD22	1:B:366:ILE:HG22	1.51	0.88
1:B:13:ILE:HD13	1:B:84:MSE:HE2	1.63	0.79
1:B:13:ILE:HD13	1:B:84:MSE:CE	2.11	0.79
1:A:290:THR:HG21	1:A:458:CYS:SG	2.26	0.75
1:A:24:MSE:HE1	1:A:79:VAL:HG21	1.70	0.73
1:B:180:LEU:HD22	1:B:366:ILE:CG2	2.21	0.68
1:A:53:SER:OG	1:A:126:ASN:ND2	2.27	0.67
1:B:214:ILE:CG2	1:B:218:MSE:HE3	2.25	0.66
1:A:290:THR:CG2	1:A:458:CYS:SG	2.87	0.62
1:A:448:MSE:HE3	4:A:671:HOH:O	2.03	0.58
1:A:214:ILE:HG23	1:A:286:VAL:HG11	1.86	0.58
1:A:24:MSE:HE1	1:A:79:VAL:CG2	2.34	0.58
1:B:290:THR:HG21	1:B:458:CYS:SG	2.44	0.58
1:B:290:THR:HG23	1:B:292:GLY:H	1.70	0.57
1:A:447:SER:O	1:A:448:MSE:HE2	2.05	0.55
1:A:180:LEU:CD2	1:A:366:ILE:HG22	2.38	0.54

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:240:ARG:HD3	1:B:277:GLU:HB3	1.90	0.53
1:B:218:MSE:HE1	1:B:320:PHE:HE1	1.74	0.53
1:B:308:HIS:HE1	1:B:362:LEU:HD21	1.74	0.52
1:A:194:MSE:HG3	1:A:321:LEU:O	2.10	0.52
1:B:371:ASN:ND2	1:B:422:GLN:HB2	2.25	0.52
1:A:290:THR:HG23	1:A:292:GLY:H	1.75	0.52
1:A:240:ARG:HD3	1:A:277:GLU:HB3	1.91	0.51
1:A:301:LEU:HD13	1:A:319:TYR:CD2	2.48	0.47
1:B:180:LEU:CD2	1:B:366:ILE:CG2	2.93	0.47
1:B:53:SER:OG	1:B:126:ASN:ND2	2.48	0.46
1:A:95:ASN:O	1:A:96:LEU:HD23	2.17	0.45
1:A:78:THR:O	1:A:79:VAL:C	2.53	0.45
1:A:39:LEU:HD13	1:A:140:LEU:HB3	1.99	0.45
1:A:9:PHE:HB3	1:A:214:ILE:HD12	1.98	0.44
1:B:290:THR:HG23	1:B:292:GLY:N	2.31	0.44
1:A:71:LEU:HD23	1:A:71:LEU:HA	1.90	0.43
1:A:309:ASP:O	1:A:310:ARG:C	2.56	0.43
1:B:297:ILE:N	1:B:298:PRO:HD2	2.35	0.42
1:B:436:GLY:O	1:B:439:MSE:HB2	2.18	0.42
1:A:148:ILE:HD12	1:A:336:ILE:HG12	2.01	0.41
1:A:311:MSE:HE3	1:A:315:ASP:HB3	2.03	0.41
1:A:330:TYR:CE1	1:A:377:MSE:HE3	2.56	0.41
1:B:129:ARG:HD3	1:B:142:GLU:HB2	2.02	0.41
1:A:378:GLU:OE1	1:B:398:GLU:OE2	2.38	0.41
1:B:377:MSE:O	1:B:378:GLU:C	2.59	0.41
1:B:35:GLU:OE1	1:B:102:LYS:NZ	2.46	0.40
1:B:308:HIS:HE1	1:B:362:LEU:CD2	2.34	0.40
1:B:352:SER:HA	1:B:376:ALA:HB3	2.04	0.40
1:B:290:THR:CG2	1:B:458:CYS:SG	3.08	0.40
1:B:451:LEU:HD12	1:B:451:LEU:HA	1.95	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	438/476 (92%)	416 (95%)	20 (5%)	2 (0%)	29	29
1	B	442/476 (93%)	424 (96%)	17 (4%)	1 (0%)	47	55
All	All	880/952 (92%)	840 (96%)	37 (4%)	3 (0%)	41	46

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	79	VAL
1	A	141	ILE
1	B	338	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/368 (98%)	344 (96%)	15 (4%)	30	34
1	B	364/368 (99%)	345 (95%)	19 (5%)	23	24
All	All	723/736 (98%)	689 (95%)	34 (5%)	26	29

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	GLU
1	A	79	VAL
1	A	98	ASN
1	A	104	GLU
1	A	109	GLU
1	A	167	ASN
1	A	212	LEU
1	A	284	ARG
1	A	290	THR
1	A	310	ARG

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	326	ILE
1	A	398	GLU
1	A	422	GLN
1	A	427	LYS
1	A	441	SER
1	B	36	GLN
1	B	98	ASN
1	B	156	GLU
1	B	157	GLU
1	B	160	ASP
1	B	178	GLU
1	B	227	LYS
1	B	253	LYS
1	B	257	SER
1	B	258	VAL
1	B	261	GLN
1	B	284	ARG
1	B	290	THR
1	B	326	ILE
1	B	398	GLU
1	B	422	GLN
1	B	427	LYS
1	B	434	GLN
1	B	451	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	95	ASN
1	A	98	ASN
1	A	126	ASN
1	A	224	ASN
1	A	267	HIS
1	A	422	GLN
1	B	98	ASN
1	B	126	ASN
1	B	186	HIS
1	B	208	HIS
1	B	224	ASN
1	B	306	GLN
1	B	308	HIS
1	B	422	GLN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	434	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SF4	B	501	1	0,12,12	0.00	-	-		
2	SF4	A	500	1	0,12,12	0.00	-	-		

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	B	501	1	-	-	0/6/5/5
2	SF4	A	500	1	-	-	0/6/5/5

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	426/476 (89%)	0.48	43 (10%) <b>7</b> <b>7</b>	26, 45, 72, 98	0
1	B	429/476 (90%)	0.18	21 (4%) <b>29</b> <b>32</b>	25, 38, 58, 103	0
All	All	855/952 (89%)	0.33	64 (7%) <b>14</b> <b>15</b>	25, 41, 67, 103	0

All (64) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	309	ASP	8.0
1	A	38	ASN	6.2
1	A	136	ASN	4.6
1	B	446	THR	4.4
1	A	446	THR	4.3
1	B	445	GLU	4.3
1	A	262	SER	4.2
1	A	138	ASN	4.1
1	A	160	ASP	3.8
1	A	249	LEU	3.7
1	A	296	ILE	3.5
1	A	333	GLY	3.4
1	B	308	HIS	3.4
1	A	445	GLU	3.4
1	A	250	ILE	3.4
1	B	137	ALA	3.3
1	B	447	SER	3.1
1	A	261	GLN	3.1
1	B	297	ILE	3.1
1	A	161	LYS	3.0
1	A	184	LYS	3.0
1	A	447	SER	2.9
1	A	309	ASP	2.9
1	A	335	SER	2.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	A	212	LEU	2.8
1	B	310	ARG	2.6
1	A	119	GLU	2.6
1	B	296	ILE	2.6
1	B	405	VAL	2.5
1	A	443	TYR	2.5
1	A	134	ASP	2.5
1	A	187	HIS	2.5
1	B	307	ALA	2.5
1	B	160	ASP	2.5
1	B	2	ASN	2.4
1	A	123	LYS	2.4
1	A	347	VAL	2.4
1	A	297	ILE	2.4
1	A	149	GLY	2.3
1	A	135	GLY	2.3
1	A	159	PHE	2.3
1	A	209	LYS	2.3
1	B	200	THR	2.3
1	A	336	ILE	2.3
1	A	244	ASP	2.2
1	A	167	ASN	2.2
1	B	136	ASN	2.2
1	A	158	ASP	2.2
1	A	295	GLY	2.2
1	B	135	GLY	2.2
1	A	32	GLN	2.2
1	B	167	ASN	2.2
1	A	201	TRP	2.2
1	A	36	GLN	2.2
1	A	67	ILE	2.1
1	B	401	ALA	2.1
1	A	349	VAL	2.1
1	A	404	ALA	2.0
1	A	310	ARG	2.0
1	A	43	THR	2.0
1	B	356	ALA	2.0
1	B	406	LYS	2.0
1	B	358	LEU	2.0
1	A	355	ALA	2.0



## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	CL	B	502	1/1	0.85	0.22	69,69,69,69	0
2	SF4	A	500	8/8	0.95	0.11	30,33,34,36	0
3	CL	B	503	1/1	0.95	0.28	60,60,60,60	0
2	SF4	B	501	8/8	0.96	0.09	28,31,34,35	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.