



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 15, 2020 – 06:32 am BST

PDB ID : 4RQY
Title : RE-REFINED STRUCTURE OF 1TE0 - STRUCTURAL ANALYSIS of
DEGS, A STRESS SENSOR OF THE BACTERIAL PERIPLASM
Authors : Sauer, R.T.; Grant, R.A.
Deposited on : 2014-11-05
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	319	 10% 94% 6%
1	B	319	 15% 89% 9%

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 8912 atoms, of which 4407 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protease degS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	299	4461	1385	2246	394	431	5	0	0	0
1	B	291	4306	1341	2161	384	415	5	0	0	0

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	78	Total 78	O 78	0	0
3	B	57	Total 57	O 57	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protease degS



- Molecule 1: Protease degS



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 3	Depositor
Cell constants a, b, c, α , β , γ	166.28Å 166.28Å 166.28Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.52 – 2.20 28.52 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.1 (28.52-2.20) 99.1 (28.52-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.18 (at 2.20Å)	Xtrriage
Refinement program	PHENIX dev_1760	Depositor
R, R_{free}	0.182 , 0.220 0.186 , 0.225	Depositor DCC
R_{free} test set	1927 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	47.5	Xtrriage
Anisotropy	0.000	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 68.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.031 for -l,-k,-h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8912	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.40	0/2239	0.56	0/3043
1	B	0.37	0/2169	0.54	0/2949
All	All	0.39	0/4408	0.55	0/5992

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2215	2246	2246	0	0
1	B	2145	2161	2173	4	0
2	A	10	0	0	0	0
3	A	78	0	0	0	0
3	B	57	0	0	1	0
All	All	4505	4407	4419	4	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

All (4) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:THR:OG1	3:B:449:HOH:O	2.10	0.68
1:B:179:ILE:HD12	1:B:239:GLN:HB2	1.94	0.49
1:B:338:ASP:O	1:B:339:ASP:CB	2.65	0.44
1:B:181:LEU:HD21	1:B:247:LYS:HE3	2.01	0.42

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	289/319 (91%)	287 (99%)	2 (1%)	0	100	100
1	B	283/319 (89%)	278 (98%)	5 (2%)	0	100	100
All	All	572/638 (90%)	565 (99%)	7 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/258 (93%)	239 (100%)	0	100	100
1	B	231/258 (90%)	231 (100%)	0	100	100
All	All	470/516 (91%)	470 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	A	402	-	4,4,4	0.92	0	6,6,6	0.51	0
2	PO4	A	401	-	4,4,4	0.83	0	6,6,6	0.50	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	299/319 (93%)	0.41	33 (11%) 5 4	33, 56, 109, 141	0
1	B	291/319 (91%)	0.65	47 (16%) 1 1	37, 63, 145, 176	0
All	All	590/638 (92%)	0.53	80 (13%) 3 2	33, 59, 129, 176	0

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	220	PHE	6.6
1	A	162	TYR	6.2
1	A	275	GLY	5.8
1	A	69	ASN	5.6
1	B	313	ILE	5.6
1	A	228	THR	5.6
1	B	339	ASP	5.5
1	A	276	ILE	5.4
1	A	220	PHE	5.4
1	A	165	GLY	5.2
1	A	266	ILE	5.1
1	B	335	VAL	5.1
1	B	352	PRO	5.0
1	A	70	SER	5.0
1	A	353	ALA	5.0
1	B	221	ASP	5.0
1	B	183	PRO	5.0
1	A	74	LEU	4.9
1	B	184	THR	4.9
1	A	73	GLN	4.8
1	B	296	ALA	4.7
1	B	340	LYS	4.7
1	B	185	GLY	4.5
1	B	289	PRO	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	278	GLN	4.5
1	B	300	VAL	4.5
1	B	266	ILE	4.4
1	B	280	GLN	4.3
1	B	337	ARG	4.3
1	B	264	ARG	4.2
1	B	297	GLY	4.1
1	A	186	ARG	4.0
1	B	342	LEU	4.0
1	B	299	GLN	3.9
1	B	229	PRO	3.9
1	B	37	PHE	3.8
1	B	219	SER	3.6
1	A	135	THR	3.6
1	B	303	LEU	3.5
1	B	165	GLY	3.5
1	B	291	GLY	3.5
1	B	344	LEU	3.5
1	B	40	THR	3.4
1	B	338	ASP	3.3
1	A	68	THR	3.3
1	B	301	ASN	3.3
1	A	221	ASP	3.3
1	B	159	GLY	3.2
1	A	166	GLN	3.1
1	B	305	ILE	3.0
1	A	279	LEU	3.0
1	B	341	GLN	3.0
1	B	316	LEU	2.9
1	A	67	ASN	2.8
1	B	213	MET	2.7
1	A	203	GLY	2.6
1	A	264	ARG	2.6
1	B	336	MET	2.6
1	A	167	THR	2.5
1	B	290	ASP	2.5
1	A	41	ASP	2.5
1	A	39	SER	2.4
1	A	136	GLY	2.4
1	B	38	ASP	2.4
1	B	136	GLY	2.4
1	B	298	ILE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	345	GLN	2.3
1	A	204	ALA	2.2
1	B	71	HIS	2.2
1	B	283	VAL	2.2
1	B	258	TYR	2.1
1	B	146	ARG	2.1
1	A	161	PRO	2.1
1	A	163	ASN	2.1
1	A	66	LEU	2.1
1	B	205	LEU	2.1
1	A	160	ASN	2.0
1	B	157	ALA	2.0
1	A	205	LEU	2.0
1	A	157	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PO4	A	402	5/5	0.94	0.15	66,69,82,91	5
2	PO4	A	401	5/5	0.97	0.11	51,74,80,88	0

6.5 Other polymers [i](#)

There are no such residues in this entry.