



wwPDB X-ray Structure Validation Summary Report ⓘ

Sep 14, 2023 – 01:40 PM EDT

PDB ID : 4RRP
Title : Crystal Structure of the Fab complexed with antigen Asf1p, Northeast Structural Genomics Consortium (NESG) Target PdR16
Authors : Kuzin, A.; Lew, S.; Seetharaman, J.; Mao, L.; Xiao, R.; Oconnell, P.T.; Maglaqui, M.; Bailey, L.; Everett, J.K.; Acton, T.B.; Montelione, G.T.; Hunt, J.F.; Tong, L.; Chaperone-Enabled Studies of Epigenetic Regulation Enzymes (CEBS); Northeast Structural Genomics Consortium (NESG)
Deposited on : 2014-11-06
Resolution : 2.79 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.79 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	216	83% 14% ..
1	B	216	85% 14%
1	C	216	% 73% 22% . .
1	D	216	85% 14% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	E	216	 % 79% 20% .
1	F	216	 7% 68% 28% ..
2	G	238	 77% 13% . 8%
2	H	238	 79% 13% . 7%
2	I	238	 3% 74% 16% . 8%
2	J	238	 85% 7% . 7%
2	K	238	 3% 66% 21% . 10%
2	L	238	 13% 67% 21% . 10%
3	M	160	 % 80% 15% ..
3	N	160	 3% 76% 18% . 5%
3	O	160	 % 86% 9% .
3	P	160	 % 83% 13% ..
3	Q	160	 3% 72% 22% ..
3	R	160	 6% 71% 18% . 11%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 27161 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Fab antibody, light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	213	Total 1655	C 1035	N 282	O 333	S 5	0	2	0
1	B	216	Total 1656	C 1035	N 278	O 336	S 7	0	0	0
1	C	212	Total 1627	C 1019	N 274	O 329	S 5	0	0	0
1	D	214	Total 1650	C 1032	N 279	O 333	S 6	0	1	0
1	E	214	Total 1642	C 1027	N 276	O 334	S 5	0	0	0
1	F	214	Total 1642	C 1027	N 276	O 333	S 6	0	0	0

- Molecule 2 is a protein called Fab antibody, heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	G	219	Total 1633	C 1031	N 272	O 323	S 7	0	1	0
2	H	221	Total 1651	C 1041	N 276	O 328	S 6	0	1	0
2	I	220	Total 1634	C 1032	N 271	O 325	S 6	0	0	0
2	J	222	Total 1646	C 1038	N 273	O 328	S 7	0	0	0
2	K	214	Total 1598	C 1012	N 264	O 316	S 6	0	0	0
2	L	214	Total 1599	C 1012	N 264	O 316	S 7	0	0	0

- Molecule 3 is a protein called Antigen Asf1p.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	M	153	Total	C	N	O	S	0	0	0
			1222	784	198	238	2			
3	N	152	Total	C	N	O	S	0	1	0
			1225	788	199	236	2			
3	O	153	Total	C	N	O	S	0	0	0
			1218	781	196	239	2			
3	P	156	Total	C	N	O	S	0	1	0
			1252	802	206	242	2			
3	Q	153	Total	C	N	O	S	0	1	0
			1234	794	201	237	2			
3	R	143	Total	C	N	O	S	0	0	0
			1147	737	187	221	2			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
M	0	GLY	-	expression tag	UNP E7KE71
M	1	SER	-	expression tag	UNP E7KE71
N	0	GLY	-	expression tag	UNP E7KE71
N	1	SER	-	expression tag	UNP E7KE71
O	0	GLY	-	expression tag	UNP E7KE71
O	1	SER	-	expression tag	UNP E7KE71
P	0	GLY	-	expression tag	UNP E7KE71
P	1	SER	-	expression tag	UNP E7KE71
Q	0	GLY	-	expression tag	UNP E7KE71
Q	1	SER	-	expression tag	UNP E7KE71
R	0	GLY	-	expression tag	UNP E7KE71
R	1	SER	-	expression tag	UNP E7KE71

- Molecule 4 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	21	Total	O	0	0
			21	21		
5	G	22	Total	O	0	0
			22	22		
5	B	20	Total	O	0	0
			20	20		
5	H	27	Total	O	0	0
			27	27		
5	C	15	Total	O	0	0
			15	15		
5	I	14	Total	O	0	0
			14	14		
5	D	16	Total	O	0	0
			16	16		
5	J	21	Total	O	0	0
			21	21		
5	K	3	Total	O	0	0
			3	3		
5	F	3	Total	O	0	0
			3	3		
5	L	2	Total	O	0	0
			2	2		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	M	10	Total O 10 10	0	0
5	N	15	Total O 15 15	0	0
5	O	16	Total O 16 16	0	0
5	P	13	Total O 13 13	0	0
5	Q	3	Total O 3 3	0	0
5	R	2	Total O 2 2	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Fab antibody, light chain




- Molecule 1: Fab antibody, light chain




- Molecule 1: Fab antibody, light chain




- Molecule 1: Fab antibody, light chain




- Molecule 1: Fab antibody, light chain

Chain E: 





- Molecule 1: Fab antibody, light chain

Chain F: 




- Molecule 2: Fab antibody, heavy chain

Chain G: 



- Molecule 2: Fab antibody, heavy chain

Chain H: 




- Molecule 2: Fab antibody, heavy chain

Chain I: 








• Molecule 2: Fab antibody, heavy chain





• Molecule 2: Fab antibody, heavy chain




• Molecule 2: Fab antibody, heavy chain




• Molecule 3: Antigen Asf1p





• Molecule 3: Antigen Asf1p









• Molecule 3: Antigen Asf1p




• Molecule 3: Antigen Asf1p



• Molecule 3: Antigen Asf1p



• Molecule 3: Antigen Asf1p



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	92.72Å 181.31Å 259.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.69 – 2.79 49.69 – 2.79	Depositor EDS
% Data completeness (in resolution range)	99.4 (49.69-2.79) 99.4 (49.69-2.79)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.94 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.183 , 0.253 0.186 , 0.255	Depositor DCC
R_{free} test set	1999 reflections (1.84%)	wwPDB-VP
Wilson B-factor (Å ²)	38.7	Xtrriage
Anisotropy	0.376	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 41.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	27161	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.04% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/1694	0.60	0/2299
1	B	0.52	0/1692	0.60	1/2297 (0.0%)
1	C	0.49	0/1663	0.59	0/2259
1	D	0.52	0/1689	0.61	0/2293
1	E	0.41	0/1678	0.53	0/2279
1	F	0.37	0/1678	0.55	0/2279
2	G	0.53	0/1675	0.63	0/2283
2	H	0.51	0/1691	0.60	1/2305 (0.0%)
2	I	0.51	0/1674	0.67	2/2283 (0.1%)
2	J	0.53	0/1686	0.64	0/2299
2	K	0.38	0/1637	0.51	0/2233
2	L	0.38	0/1637	0.54	0/2231
3	M	0.50	0/1250	0.60	0/1704
3	N	0.47	0/1256	0.61	0/1712
3	O	0.51	0/1245	0.60	0/1697
3	P	0.52	0/1284	0.59	0/1750
3	Q	0.40	0/1266	0.55	0/1727
3	R	0.40	0/1173	0.55	0/1597
All	All	0.47	0/27568	0.59	4/37527 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	I	138	LEU	CA-CB-CG	5.95	128.97	115.30
2	I	66	ARG	NE-CZ-NH2	-5.20	117.70	120.30
1	B	66	ARG	NE-CZ-NH1	-5.16	117.72	120.30
2	H	178	LEU	CA-CB-CG	5.07	126.97	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	-1	MET	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1655	0	1619	18	0
1	B	1656	0	1613	17	0
1	C	1627	0	1588	25	0
1	D	1650	0	1612	12	0
1	E	1642	0	1599	17	0
1	F	1642	0	1599	37	0
2	G	1633	0	1601	16	0
2	H	1651	0	1619	15	0
2	I	1634	0	1602	20	0
2	J	1646	0	1612	8	0
2	K	1598	0	1565	36	0
2	L	1599	0	1564	27	0
3	M	1222	0	1197	12	0
3	N	1225	0	1210	13	0
3	O	1218	0	1195	5	0
3	P	1252	0	1237	11	0
3	Q	1234	0	1221	16	0
3	R	1147	0	1124	12	0
4	A	7	0	10	3	0
5	A	21	0	0	0	0
5	B	20	0	0	0	0
5	C	15	0	0	0	0
5	D	16	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	F	3	0	0	0	0
5	G	22	0	0	1	0
5	H	27	0	0	0	0
5	I	14	0	0	0	0
5	J	21	0	0	0	0
5	K	3	0	0	0	0
5	L	2	0	0	0	0
5	M	10	0	0	0	0
5	N	15	0	0	0	0
5	O	16	0	0	0	0
5	P	13	0	0	0	0
5	Q	3	0	0	0	0
5	R	2	0	0	0	0
All	All	27161	0	26387	301	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 301 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:189:HIS:O	1:C:211:ARG:NH1	2.12	0.81
3:Q:35:LYS:O	3:Q:36:HIS:ND1	2.17	0.76
1:F:214:CYS:HG	2:L:127:SER:HG	1.34	0.75
1:C:120:PRO:HD3	1:C:132:VAL:HG13	1.70	0.73
1:F:12:SER:HB2	1:F:107:LYS:HD3	1.70	0.73

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	213/216 (99%)	201 (94%)	10 (5%)	2 (1%)	17	46
1	B	214/216 (99%)	201 (94%)	12 (6%)	1 (0%)	29	61
1	C	210/216 (97%)	197 (94%)	12 (6%)	1 (0%)	29	61
1	D	213/216 (99%)	198 (93%)	15 (7%)	0	100	100
1	E	212/216 (98%)	189 (89%)	22 (10%)	1 (0%)	29	61
1	F	212/216 (98%)	182 (86%)	27 (13%)	3 (1%)	11	34
2	G	216/238 (91%)	212 (98%)	3 (1%)	1 (0%)	29	61
2	H	220/238 (92%)	213 (97%)	7 (3%)	0	100	100
2	I	218/238 (92%)	203 (93%)	14 (6%)	1 (0%)	29	61
2	J	220/238 (92%)	212 (96%)	8 (4%)	0	100	100
2	K	210/238 (88%)	189 (90%)	18 (9%)	3 (1%)	11	34
2	L	208/238 (87%)	184 (88%)	18 (9%)	6 (3%)	4	15
3	M	149/160 (93%)	142 (95%)	3 (2%)	4 (3%)	5	17
3	N	149/160 (93%)	139 (93%)	8 (5%)	2 (1%)	12	36
3	O	149/160 (93%)	142 (95%)	5 (3%)	2 (1%)	12	36
3	P	155/160 (97%)	146 (94%)	8 (5%)	1 (1%)	25	56
3	Q	152/160 (95%)	137 (90%)	14 (9%)	1 (1%)	22	53
3	R	137/160 (86%)	121 (88%)	12 (9%)	4 (3%)	4	15
All	All	3457/3684 (94%)	3208 (93%)	216 (6%)	33 (1%)	15	44

5 of 33 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	K	172	SER
1	F	138	ASN
2	L	207	VAL
3	O	130	VAL
3	R	3	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	190/191 (100%)	180 (95%)	10 (5%)	22	54
1	B	191/191 (100%)	183 (96%)	8 (4%)	30	63
1	C	187/191 (98%)	168 (90%)	19 (10%)	7	22
1	D	190/191 (100%)	181 (95%)	9 (5%)	26	59
1	E	189/191 (99%)	175 (93%)	14 (7%)	13	37
1	F	189/191 (99%)	171 (90%)	18 (10%)	8	25
2	G	183/201 (91%)	168 (92%)	15 (8%)	11	33
2	H	185/201 (92%)	172 (93%)	13 (7%)	15	40
2	I	183/201 (91%)	171 (93%)	12 (7%)	16	44
2	J	185/201 (92%)	176 (95%)	9 (5%)	25	57
2	K	179/201 (89%)	164 (92%)	15 (8%)	11	31
2	L	180/201 (90%)	167 (93%)	13 (7%)	14	38
3	M	140/146 (96%)	135 (96%)	5 (4%)	35	69
3	N	141/146 (97%)	131 (93%)	10 (7%)	14	39
3	O	140/146 (96%)	135 (96%)	5 (4%)	35	69
3	P	144/146 (99%)	136 (94%)	8 (6%)	21	51
3	Q	142/146 (97%)	130 (92%)	12 (8%)	10	31
3	R	132/146 (90%)	123 (93%)	9 (7%)	16	42
All	All	3070/3228 (95%)	2866 (93%)	204 (7%)	16	44

5 of 204 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	K	68	THR
2	L	58	SER
3	R	80	SER
2	K	110	THR
1	F	89	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 6 such sidechains are listed below:

Mol	Chain	Res	Type
2	I	164	HIS
2	I	199	ASN
1	F	89	GLN
2	H	28	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	137	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	PEG	A	701	-	6,6,6	0.50	0	5,5,5	0.87	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PEG	A	701	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	701	PEG	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	213/216 (98%)	-0.23	0 100 100	13, 26, 42, 61	0
1	B	216/216 (100%)	-0.31	1 (0%) 91 88	15, 33, 53, 76	0
1	C	212/216 (98%)	0.03	3 (1%) 75 70	12, 41, 71, 80	0
1	D	214/216 (99%)	-0.24	1 (0%) 91 88	14, 25, 43, 74	0
1	E	214/216 (99%)	0.02	2 (0%) 84 80	32, 52, 68, 83	0
1	F	214/216 (99%)	0.67	15 (7%) 16 9	50, 71, 86, 104	0
2	G	219/238 (92%)	-0.34	1 (0%) 91 88	14, 23, 47, 74	0
2	H	221/238 (92%)	-0.22	1 (0%) 91 88	15, 26, 59, 86	0
2	I	220/238 (92%)	0.01	8 (3%) 42 32	12, 28, 75, 89	0
2	J	222/238 (93%)	-0.45	0 100 100	11, 25, 45, 70	0
2	K	214/238 (89%)	0.22	6 (2%) 53 43	26, 63, 87, 106	0
2	L	214/238 (89%)	0.79	30 (14%) 2 1	31, 68, 96, 104	0
3	M	153/160 (95%)	-0.09	1 (0%) 87 84	16, 29, 64, 81	0
3	N	152/160 (95%)	0.05	5 (3%) 46 36	17, 34, 79, 88	1 (0%)
3	O	153/160 (95%)	-0.16	1 (0%) 87 84	15, 27, 68, 79	0
3	P	156/160 (97%)	-0.26	1 (0%) 89 86	16, 25, 54, 76	0
3	Q	153/160 (95%)	0.27	5 (3%) 46 36	28, 54, 86, 95	0
3	R	143/160 (89%)	0.52	9 (6%) 20 12	32, 64, 96, 106	0
All	All	3503/3684 (95%)	0.01	90 (2%) 56 46	11, 36, 82, 106	1 (0%)

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	L	112	SER	5.7
2	L	119	PRO	4.7
2	L	149	PRO	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	161	SER	4.2
1	F	106	ILE	4.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	PEG	A	701	7/7	0.95	0.31	31,37,43,43	0

6.5 Other polymers [i](#)

There are no such residues in this entry.