



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 31, 2021 – 07:16 PM EDT

PDB ID : 1RSV
Title : azide complex of the diferrous E238A mutant R2 subunit of ribonucleotide reductase
Authors : Assarsson, M.; Andersson, M.E.; Hogbom, M.; Persson, B.O.; Sahlin, M.; Barra, A.L.; Sjoberg, B.M.; Nordlund, P.; Graslund, A.
Deposited on : 2003-12-10
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	 3% 56% 29% 6% 9%
1	B	375	 3% 48% 33% 10% 9%

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 5919 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Ribonucleoside-diphosphate reductase 1 beta chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	340	2783	1782	464	524	13	0	0	0
1	B	341	2789	1785	465	526	13	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	122	PHE	TYR	engineered mutation	UNP P69924
A	238	ALA	GLU	engineered mutation	UNP P69924
B	122	PHE	TYR	engineered mutation	UNP P69924
B	238	ALA	GLU	engineered mutation	UNP P69924

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Fe	0	0
			2	2		
2	B	2	Total	Fe	0	0
			2	2		

- Molecule 3 is MERCURY (II) ION (three-letter code: HG) (formula: Hg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	6	Total	Hg	0	0
			6	6		
3	B	6	Total	Hg	0	0
			6	6		

- Molecule 4 is AZIDE ION (three-letter code: AZI) (formula: N₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total N 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	161	Total O 161 161	0	0
5	B	167	Total O 167 167	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ribonucleoside-diphosphate reductase 1 beta chain



- Molecule 1: Ribonucleoside-diphosphate reductase 1 beta chain



ASN
VAL
GLN
VAL
VAL
ALA
PRO
GLN
GLU
VAL
GLU
VAL
SER
SER
TYR
LEU
VAL
VAL
GLY
GLN
ILE
ASP
SER
GLU
VAL
VAL
ASP
THR
ASP
ASP
LEU
SER
ASN
ASN
PHE
GLN
LEU

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.89Å 84.50Å 113.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	17.00 – 2.20 16.74 – 2.19	Depositor EDS
% Data completeness (in resolution range)	(Not available) (17.00-2.20) 97.2 (16.74-2.19)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 2.18Å)	Xtrriage
Refinement program	TNT	Depositor
R, R_{free}	0.182 , 0.256 0.174 , 0.171	Depositor DCC
R_{free} test set	1798 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	17.4	Xtrriage
Anisotropy	0.489	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 79.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5919	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, AZI, HG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.04	16/2847 (0.6%)	1.11	9/3862 (0.2%)
1	B	1.04	15/2853 (0.5%)	1.11	9/3870 (0.2%)
All	All	1.04	31/5700 (0.5%)	1.11	18/7732 (0.2%)

The worst 5 of 31 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	214	CYS	CB-SG	8.64	1.97	1.82
1	B	222	GLU	CD-OE2	7.24	1.33	1.25
1	A	41	GLU	CD-OE2	6.99	1.33	1.25
1	B	271	GLU	CD-OE2	6.96	1.33	1.25
1	B	188	GLU	CD-OE2	6.94	1.33	1.25

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	214	CYS	CA-CB-SG	10.29	132.53	114.00
1	B	214	CYS	CB-CA-C	7.28	124.96	110.40
1	B	89	ARG	NE-CZ-NH2	-7.07	116.77	120.30
1	A	132	ASP	CB-CG-OD2	-6.60	112.36	118.30
1	B	54	ASP	CB-CG-OD2	-6.57	112.39	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2783	0	2732	121	0
1	B	2789	0	2735	208	0
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	6	0	0	4	0
3	B	6	0	0	1	0
4	A	3	0	0	1	0
5	A	161	0	0	17	0
5	B	167	0	0	28	0
All	All	5919	0	5467	311	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

The worst 5 of 311 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:2004:HG:HG	5:B:2175:HOH:O	1.15	1.18
3:A:2009:HG:HG	5:A:2165:HOH:O	1.34	1.05
3:A:2012:HG:HG	5:A:2151:HOH:O	1.32	1.03
1:A:205:ALA:HB1	1:A:315:ARG:HD3	1.36	1.02
3:A:2007:HG:HG	5:A:2127:HOH:O	1.37	1.00

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	338/375 (90%)	324 (96%)	14 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	339/375 (90%)	323 (95%)	15 (4%)	1 (0%)	41	46
All	All	677/750 (90%)	647 (96%)	29 (4%)	1 (0%)	51	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	340	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	306/339 (90%)	282 (92%)	24 (8%)	12	13
1	B	307/339 (91%)	266 (87%)	41 (13%)	4	3
All	All	613/678 (90%)	548 (89%)	65 (11%)	6	6

5 of 65 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	292	ARG
1	B	296	MET
1	B	9	LYS
1	A	340	VAL
1	B	297	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	76	ASN
1	B	128	ASN
1	B	247	HIS
1	B	87	GLN
1	B	131	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 17 ligands modelled in this entry, 16 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	AZI	A	901	2	0,2,2	-	-	0,1,1	-	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	901	AZI	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	340/375 (90%)	-0.51	3 (0%) 84 83	14, 25, 51, 73	0
1	B	341/375 (90%)	-0.25	13 (3%) 40 38	12, 28, 58, 84	0
All	All	681/750 (90%)	-0.38	16 (2%) 60 58	12, 27, 56, 84	0

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	340	VAL	6.5
1	B	297	ILE	6.3
1	B	289	TYR	5.0
1	B	291	PHE	4.1
1	B	341	SER	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	AZI	A	901	3/3	0.91	0.23	22,22,30,39	3
3	HG	A	2011	1/1	0.93	0.17	40,40,40,40	1
3	HG	B	2002	1/1	0.97	0.06	39,39,39,39	1
3	HG	B	2003	1/1	0.97	0.04	49,49,49,49	1
3	HG	B	2006	1/1	0.97	0.10	35,35,35,35	1
3	HG	A	2008	1/1	0.97	0.06	37,37,37,37	1
3	HG	B	2013	1/1	0.98	0.05	26,26,26,26	1
2	FE	B	1001	1/1	0.98	0.04	41,41,41,41	1
3	HG	B	2010	1/1	0.99	0.03	42,42,42,42	1
2	FE	B	1002	1/1	0.99	0.06	24,24,24,24	0
2	FE	A	1003	1/1	0.99	0.04	22,22,22,22	1
2	FE	A	1004	1/1	1.00	0.03	17,17,17,17	0
3	HG	B	2004	1/1	1.00	0.03	23,23,23,23	1
3	HG	A	2009	1/1	1.00	0.03	24,24,24,24	1
3	HG	A	2001	1/1	1.00	0.02	24,24,24,24	1
3	HG	A	2012	1/1	1.00	0.04	31,31,31,31	1
3	HG	A	2007	1/1	1.00	0.02	24,24,24,24	1

6.5 Other polymers [i](#)

There are no such residues in this entry.