



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 11:44 PM EDT

PDB ID : 7RTK
Title : Structure of the (NIAU)2 complex with N-terminal mutation of ISCU2 Y35D at 2.5 Å resolution
Authors : Boniecki, M.T.; Cygler, M.
Deposited on : 2021-08-13
Resolution : 2.50 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : ?? (??), CSD ??CSD?? (????)
Xtriage (Phenix) : 1.13
EDS : 2.35.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	406	
2	B	91	
3	C	77	
4	D	143	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
13	EDT	B	108	-	-	-	X
6	EDO	A	527	-	-	-	X
6	EDO	A	531	-	-	-	X
8	PG4	A	516	-	-	-	X

2 Entry composition [i](#)

There are 17 unique types of molecules in this entry. The entry contains 5798 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine desulfurase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	395	3071	1935	535	580	21	0	5	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	52	MET	-	initiating methionine	UNP Q9Y697
A	53	GLY	-	expression tag	UNP Q9Y697
A	54	SER	-	expression tag	UNP Q9Y697
A	55	SER	-	expression tag	UNP Q9Y697

- Molecule 2 is a protein called LYR motif-containing protein 4.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	84	713	449	140	123	1	0	5	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	11	ALA	SER	variant	UNP Q9HD34

- Molecule 3 is a protein called Acyl carrier protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	75	528	331	84	112	1	0	0	0


- Molecule 4 is a protein called Iron-sulfur cluster assembly enzyme ISCU, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	D	124	919	580	156	175	8	0	3	0

There are 11 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
D	33	MET	-	initiating methionine	UNP Q9H1K1
D	34	ALA	-	expression tag	UNP Q9H1K1
D	35	ASP	TYR	conflict	UNP Q9H1K1
D	168	LEU	-	expression tag	UNP Q9H1K1
D	169	GLU	-	expression tag	UNP Q9H1K1
D	170	HIS	-	expression tag	UNP Q9H1K1
D	171	HIS	-	expression tag	UNP Q9H1K1
D	172	HIS	-	expression tag	UNP Q9H1K1
D	173	HIS	-	expression tag	UNP Q9H1K1
D	174	HIS	-	expression tag	UNP Q9H1K1
D	175	HIS	-	expression tag	UNP Q9H1K1

- Molecule 5 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: $C_8H_{10}NO_6P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
5	A	1	15	8	1	5	1	0	0

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0
6	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	A	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	B	1	Total 4	C 2	O 2	0	0
6	C	1	Total 4	C 2	O 2	0	0
6	D	1	Total 4	C 2	O 2	0	0
6	D	1	Total 4	C 2	O 2	0	0
6	D	1	Total 4	C 2	O 2	0	0

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	A	1	Total	C	O	0	0
			6	3	3		
7	D	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 9 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: $C_4H_{10}O_3$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	A	1	Total C O 7 4 3	0	0
9	A	1	Total C O 7 4 3	0	0
9	A	1	Total C O 7 4 3	0	0
9	A	1	Total C O 7 4 3	0	0
9	A	1	Total C O 7 4 3	0	0
9	B	1	Total C O 7 4 3	0	0
9	B	1	Total C O 7 4 3	0	0
9	B	1	Total C O 7 4 3	0	0

- Molecule 10 is 2,5,8,11,14,17-HEXAOXANONADECAN-19-OL (three-letter code: P15) (formula: $C_{13}H_{28}O_7$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
10	A	1	Total	C	O	0	0
			20	13	7		

- Molecule 11 is 2,3-DIHYDROXY-1,4-DITHIOBUTANE (three-letter code: DTT) (formula: $C_4H_{10}O_2S_2$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
11	A	1	Total	C	O	S	0	0
			8	4	2	2		

- Molecule 12 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
12	A	1	Total C O 10 6 4	0	0
12	A	1	Total C O 10 6 4	0	0
12	A	1	Total C O 10 6 4	0	0

- Molecule 13 is {[-(BIS-CARBOXYMETHYL-AMINO)-ETHYL]-CARBOXYMETHYL-AMINO}-ACETIC ACID (three-letter code: EDT) (formula: C₁₀H₁₆N₂O₈).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
13	B	1	20	10	2	8	0	0

- Molecule 14 is S-[2-({N-[(2R)-2-hydroxy-3,3-dimethyl-4-(phosphonoxy)butanoyl]-beta-alanyl}amino)ethyl] dodecanethioate (three-letter code: 8Q1) (formula: C₂₃H₄₅N₂O₈PS).




Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
14	C	1	34	23	2	7	1	1	0	0

- Molecule 15 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
15	C	1	12	6	1	4	1	0	0

- Molecule 16 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: C₁₀H₂₂O₆).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
16	D	1	16	10	6	0	0


- Molecule 17 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
17	A	163	Total	O	0	0
			163	163		
17	B	30	Total	O	0	0
			30	30		
17	C	4	Total	O	0	0
			4	4		
17	D	12	Total	O	0	0
			12	12		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cysteine desulfurase, mitochondrial




- Molecule 2: LYR motif-containing protein 4



- Molecule 3: Acyl carrier protein



- Molecule 4: Iron-sulfur cluster assembly enzyme ISCU, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	86.35Å 86.35Å 246.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.00 – 2.50 49.00 – 2.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (49.00-2.50) 100.0 (49.00-2.50)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.43 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.19.1_4122	Depositor
R, R_{free}	0.169 , 0.214 0.167 , 0.212	Depositor DCC
R_{free} test set	1663 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	44.0	Xtrriage
Anisotropy	0.191	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 55.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	5798	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PGE, EDT, PLP, 8Q1, P15, GOL, PG4, EDO, MES, 1PE, PEG, DTT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.31	2/3142 (0.1%)	0.53	2/4251 (0.0%)
2	B	0.25	0/737	0.54	0/989
3	C	0.24	0/532	0.44	0/726
4	D	0.26	0/941	0.46	0/1273
All	All	0.29	2/5352 (0.0%)	0.51	2/7239 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	105[A]	ARG	C-O	5.57	1.33	1.23
1	A	105[B]	ARG	C-O	5.57	1.33	1.23

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	105[A]	ARG	CA-C-O	5.19	131.00	120.10
1	A	105[B]	ARG	CA-C-O	5.19	131.00	120.10

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3071	0	3061	30	0
2	B	713	0	738	7	0
3	C	528	0	467	6	0
4	D	919	0	931	12	0
5	A	15	0	7	2	0
6	A	84	0	126	9	0
6	B	16	0	24	0	0
6	C	4	0	6	0	0
6	D	12	0	18	0	0
7	A	12	0	16	3	0
7	D	6	0	8	0	0
8	A	13	0	18	0	0
9	A	35	0	50	1	0
9	B	21	0	30	1	0
10	A	20	0	28	0	0
11	A	8	0	10	1	0
12	A	30	0	42	1	0
13	B	20	0	12	1	0
14	C	34	0	0	0	0
15	C	12	0	12	0	0
16	D	16	0	22	3	0
17	A	163	0	0	1	0
17	B	30	0	0	1	0
17	C	4	0	0	0	0
17	D	12	0	0	0	0
All	All	5798	0	5626	54	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (54) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:29:ARG:HH21	13:B:108:EDT:H061	1.35	0.90
1:A:105[B]:ARG:NH1	17:A:601:HOH:O	2.23	0.71
1:A:245:ASN:HD21	11:A:523:DTT:H2	1.55	0.71
1:A:175:TYR:HD1	6:A:502:EDO:H22	1.56	0.70
1:A:208:GLU:OE2	1:A:389:SER:OG	2.12	0.67
4:D:112:LYS:HG2	16:D:202:1PE:H222	1.78	0.64
4:D:95:CYS:SG	4:D:135:LYS:NZ	2.60	0.64
1:A:237:VAL:HG21	1:A:254:ILE:HG12	1.80	0.63
3:C:11:ILE:HD13	3:C:65:VAL:HG22	1.79	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:114:VAL:HG13	4:D:150:LEU:HD22	1.81	0.63
4:D:112:LYS:HE3	16:D:202:1PE:H252	1.80	0.62
6:A:506:EDO:H11	4:D:44:GLU:HG2	1.83	0.60
1:A:287:GLN:HG3	6:A:532:EDO:H11	1.82	0.60
2:B:41:ARG:NH2	3:C:35:ASP:OD2	2.35	0.60
2:B:12:LEU:HD11	2:B:60:LYS:HG3	1.82	0.60
1:A:297:PRO:HD3	6:A:512:EDO:H21	1.87	0.57
2:B:34[B]:ARG:NE	17:B:202:HOH:O	2.38	0.56
1:A:321[B]:ARG:NH2	7:A:504:GOL:O2	2.37	0.56
1:A:366:LEU:HD23	1:A:406:ILE:HG13	1.86	0.56
1:A:109:GLN:HE22	12:A:528:PGE:H5	1.71	0.55
1:A:377:SER:HB2	1:A:405:SER:O	2.09	0.53
1:A:137:GLY:HA3	1:A:278:VAL:HG13	1.91	0.53
1:A:150:ILE:HB	1:A:200:VAL:HG22	1.89	0.53
1:A:139:ALA:HB2	1:A:149:LEU:HD21	1.91	0.52
1:A:201:SER:HA	1:A:230:HIS:O	2.10	0.52
4:D:133:PRO:HA	4:D:136:LEU:HG	1.91	0.51
1:A:161:ASP:OD1	1:A:164:ARG:NH2	2.41	0.51
1:A:367:LEU:HD11	1:A:377:SER:HB3	1.93	0.51
1:A:432[B]:ARG:HH12	6:A:505:EDO:H11	1.76	0.51
1:A:153:GLN:HG2	9:A:521:PEG:H21	1.93	0.50
7:A:504:GOL:H12	2:B:34[A]:ARG:HH21	1.76	0.50
4:D:112:LYS:HZ2	16:D:202:1PE:H131	1.76	0.50
3:C:47:GLU:HG2	3:C:52:THR:O	2.11	0.49
1:A:338:PRO:HD3	6:A:507:EDO:H11	1.95	0.48
3:C:54:ILE:HG12	3:C:71:TYR:CE2	2.49	0.48
1:A:317:TYR:HB2	6:A:513:EDO:H11	1.95	0.48
1:A:181:SER:HB2	1:A:345:ASP:HB2	1.95	0.48
6:A:506:EDO:H12	4:D:43:TYR:CE2	2.49	0.48
1:A:142:TYR:CE2	1:A:228:TYR:HE2	2.34	0.46
3:C:59:ALA:HA	3:C:62:ILE:HD12	1.97	0.45
4:D:132:PRO:HD2	4:D:135:LYS:HB2	1.99	0.44
1:A:127:ALA:HB3	5:A:501:PLP:O4P	2.17	0.44
1:A:289:ARG:HB2	1:A:291[A]:MET:HE2	2.01	0.43
3:C:3:ILE:O	3:C:6:ARG:HG3	2.18	0.43
2:B:27[B]:ASN:OD1	2:B:28:TYR:N	2.52	0.43
2:B:38:ASP:HB3	9:B:103:PEG:H21	2.00	0.43
4:D:56:SER:HB3	4:D:59:VAL:HB	2.02	0.42
6:A:506:EDO:H12	4:D:43:TYR:HE2	1.84	0.42
1:A:349:HIS:NE2	1:A:353:CYS:O	2.52	0.42
1:A:440:TRP:O	1:A:444:GLN:HG2	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:D:105:ALA:HB1	4:D:146:ILE:HD11	2.03	0.41
1:A:232:ASP:OD2	5:A:501:PLP:N1	2.54	0.41
1:A:189:LEU:O	1:A:193:ILE:HG13	2.21	0.40
1:A:321[B]:ARG:NH2	7:A:504:GOL:HO2	2.19	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	396/406 (98%)	383 (97%)	13 (3%)	0	100	100
2	B	87/91 (96%)	86 (99%)	1 (1%)	0	100	100
3	C	73/77 (95%)	69 (94%)	4 (6%)	0	100	100
4	D	125/143 (87%)	125 (100%)	0	0	100	100
All	All	681/717 (95%)	663 (97%)	18 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	330/346 (95%)	324 (98%)	6 (2%)	59	81
2	B	73/80 (91%)	72 (99%)	1 (1%)	67	86

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	C	48/66 (73%)	48 (100%)	0	100	100
4	D	98/118 (83%)	94 (96%)	4 (4%)	30	55
All	All	549/610 (90%)	538 (98%)	11 (2%)	60	79

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	105[A]	ARG
1	A	105[B]	ARG
1	A	172	GLN
1	A	377	SER
1	A	379	SER
1	A	381	CYS
2	B	5	SER
4	D	51	SER
4	D	139	SER
4	D	140[A]	MET
4	D	140[B]	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	245	ASN
4	D	78	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

51 ligands are modelled in this entry.

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	395/406 (97%)	-0.33	6 (1%) 73 75	25, 37, 70, 115	3 (0%)
2	B	84/91 (92%)	-0.40	2 (2%) 59 62	29, 41, 67, 100	0
3	C	75/77 (97%)	1.28	18 (24%) 0 0	46, 78, 111, 124	0
4	D	124/143 (86%)	-0.01	8 (6%) 18 19	35, 57, 83, 131	0
All	All	678/717 (94%)	-0.10	34 (5%) 28 30	25, 43, 86, 131	3 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	17	VAL	6.4
1	A	381	CYS	5.7
1	A	378	GLY	5.3
1	A	380	ALA	4.9
3	C	2	THR	4.8
3	C	72	ILE	4.0
3	C	22	VAL	4.0
2	B	3	ALA	3.8
3	C	3	ILE	3.8
1	A	455	THR	3.7
3	C	23	THR	3.6
3	C	73	ASN	3.4
1	A	379	SER	3.3
3	C	7	VAL	3.0
4	D	155	LEU	3.0
3	C	6	ARG	2.9
3	C	76	GLN	2.9
3	C	68	ALA	2.8
3	C	11	ILE	2.8
3	C	25	ASN	2.7
2	B	2	ALA	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
4	D	154	LYS	2.6
4	D	158	GLU	2.5
4	D	118	LEU	2.5
3	C	20	GLU	2.4
4	D	157	GLN	2.4
3	C	74	GLY	2.3
3	C	24	ASN	2.3
4	D	153	TYR	2.3
1	A	172	GLN	2.3
4	D	150	LEU	2.3
3	C	50	PHE	2.3
4	D	83	GLY	2.1
3	C	21	GLU	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
6	EDO	D	205	4/4	0.42	0.39	73,77,82,95	0
6	EDO	A	531	4/4	0.61	0.43	50,67,73,73	0
13	EDT	B	108	20/20	0.62	0.43	44,61,76,79	20
6	EDO	A	526	4/4	0.66	0.34	72,74,80,85	0
6	EDO	A	532	4/4	0.66	0.35	73,76,79,82	0
6	EDO	A	513	4/4	0.67	0.25	68,69,74,80	0
8	PG4	A	516	13/13	0.70	0.43	41,63,77,80	13
6	EDO	A	527	4/4	0.70	0.56	41,56,58,59	4
6	EDO	B	101	4/4	0.71	0.34	56,59,77,82	0
16	1PE	D	202	16/16	0.71	0.28	68,87,96,101	0


Continued on next page...


Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
6	EDO	A	510	4/4	0.72	0.35	38,40,43,49	4
15	MES	C	302	12/12	0.72	0.30	63,85,103,105	12
12	PGE	A	529	10/10	0.72	0.25	75,82,86,88	0
6	EDO	A	530	4/4	0.73	0.28	55,67,84,90	0
9	PEG	A	533	7/7	0.74	0.39	63,72,95,96	0
10	P15	A	520	20/20	0.74	0.38	34,71,84,85	20
6	EDO	A	519	4/4	0.74	0.28	52,54,58,65	4
9	PEG	A	522	7/7	0.75	0.24	64,71,81,95	0
9	PEG	A	521	7/7	0.76	0.35	62,66,77,83	0
6	EDO	D	203	4/4	0.76	0.40	74,74,80,85	0
9	PEG	B	103	7/7	0.77	0.36	42,57,64,67	7
6	EDO	A	525	4/4	0.77	0.27	66,67,71,76	0
6	EDO	D	201	4/4	0.77	0.30	68,74,82,85	0
9	PEG	B	105	7/7	0.79	0.27	51,57,61,64	7
11	DTT	A	523	8/8	0.80	0.38	53,73,86,93	8
6	EDO	A	505	4/4	0.81	0.28	59,63,63,68	0
9	PEG	A	534	7/7	0.81	0.39	72,79,85,88	0
6	EDO	A	502	4/4	0.82	0.37	57,61,66,75	0
6	EDO	C	303	4/4	0.82	0.39	56,59,64,64	4
6	EDO	A	535	4/4	0.82	0.22	57,60,63,67	0
6	EDO	A	511	4/4	0.83	0.24	47,55,57,64	4
6	EDO	A	514	4/4	0.83	0.36	42,52,60,69	4
6	EDO	B	106	4/4	0.84	0.39	73,77,78,94	0
7	GOL	D	204	6/6	0.85	0.25	56,62,68,71	6
12	PGE	A	524	10/10	0.85	0.25	43,52,65,68	10
6	EDO	A	507	4/4	0.86	0.23	57,59,60,71	0
12	PGE	A	528	10/10	0.86	0.25	29,48,57,59	10
6	EDO	B	104	4/4	0.87	0.21	69,70,74,76	0
9	PEG	A	517	7/7	0.87	0.25	53,62,73,85	0
6	EDO	B	107	4/4	0.88	0.35	54,66,66,77	0
6	EDO	A	515	4/4	0.89	0.28	34,43,47,47	4
6	EDO	A	509	4/4	0.90	0.42	72,74,76,77	0
9	PEG	B	102	7/7	0.90	0.19	34,56,60,65	0
7	GOL	A	504	6/6	0.90	0.23	47,73,80,82	0
7	GOL	A	503	6/6	0.91	0.18	25,39,47,47	6
6	EDO	A	506	4/4	0.92	0.36	32,37,38,44	4
6	EDO	A	508	4/4	0.92	0.37	42,46,53,56	4
14	8Q1	C	301	34/35	0.95	0.13	35,44,58,59	0
6	EDO	A	518	4/4	0.97	0.50	43,49,52,66	4
6	EDO	A	512	4/4	0.98	0.20	28,33,35,45	4
5	PLP	A	501	15/16	0.99	0.15	29,33,41,42	0

The following is a graphical depiction of the model fit to experimental electron density of all

instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.