



Full wwPDB EM Validation Report ⓘ

Nov 23, 2022 – 03:10 PM EST

PDB ID : 7RTT
EMDB ID : EMD-24688
Title : Cryo-EM structure of a TTYH2 cis-dimer
Authors : Li, B.; Brohawn, S.G.; Hoel, C.M.
Deposited on : 2021-08-15
Resolution : 3.50 Å (reported)

This is a Full wwPDB EM Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev43
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : 1.9.9
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.31.3


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	
1	B	540	
2	C	3	
2	D	3	
2	E	3	
2	F	3	

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 6268 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein tweety homolog 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
			Total	C	N	O	S		
1	A	396	3058	1971	503	559	25	0	0
1	B	395	3053	1968	502	558	25	0	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	533	SER	-	expression tag	UNP Q3TH73
A	534	ASN	-	expression tag	UNP Q3TH73
A	535	SER	-	expression tag	UNP Q3TH73
A	536	LEU	-	expression tag	UNP Q3TH73
A	537	GLU	-	expression tag	UNP Q3TH73
A	538	VAL	-	expression tag	UNP Q3TH73
A	539	LEU	-	expression tag	UNP Q3TH73
A	540	PHE	-	expression tag	UNP Q3TH73
A	541	GLN	-	expression tag	UNP Q3TH73
B	533	SER	-	expression tag	UNP Q3TH73
B	534	ASN	-	expression tag	UNP Q3TH73
B	535	SER	-	expression tag	UNP Q3TH73
B	536	LEU	-	expression tag	UNP Q3TH73
B	537	GLU	-	expression tag	UNP Q3TH73
B	538	VAL	-	expression tag	UNP Q3TH73
B	539	LEU	-	expression tag	UNP Q3TH73
B	540	PHE	-	expression tag	UNP Q3TH73
B	541	GLN	-	expression tag	UNP Q3TH73

- Molecule 2 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				AltConf	Trace
2	C	3	Total	C	N	O	0	0
			39	22	2	15		
2	D	3	Total	C	N	O	0	0
			39	22	2	15		
2	E	3	Total	C	N	O	0	0
			39	22	2	15		
2	F	3	Total	C	N	O	0	0
			38	22	2	14		


- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		AltConf
3	A	1	Total	Ca	0
			1	1	
3	B	1	Total	Ca	0
			1	1	


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Protein tweety homolog 2



- Molecule 1: Protein tweety homolog 2



ASP ILE LEU PHE ASP GLY ASP ASP PRO PHE ASN ASN GLN GLN ALA ARG ARG ILE ALA ALA HIS ASN PRO THR THR ARG GLY GLN LEU HIS SER SER PHE CYS ARG TYR SER SER GLY LEU GLY ASP GLN CYS LEU LEU HIS GLN PRO PRO SER GLN THR ILE SER ASN ASN ALA PRO VAL VAL SER LEU PHE THR TYR MET ASN GLN ALA

ILE LEU PHE ASP GLY ASP ASN ARG TYR ASN VAL PRO LEU ILE ARG GLY HIS SER PRO THR THR TYR SER SER MET ARG PRO THR TYR MET SER VAL ALA ASP GLU HIS ARG LEU TYR PHE PRO SER ILE SER ASN SER LEU VAL VAL SER LEU PHE GLN

- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




MAG1
MAG2
BMA3

- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




MAG1
MAG2
BMA3

- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose





MAG1
MAG2
BMA3

- Molecule 2: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




MAG1
MAG2
BMA3

4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	96342	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TECNAI ARCTICA	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	1.0038	Depositor
Minimum defocus (nm)	Not provided	
Maximum defocus (nm)	Not provided	
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	58.931	Depositor
Minimum map value	-37.143	Depositor
Average map value	0.000	Depositor
Map value standard deviation	1.000	Depositor
Recommended contour level	7.0	Depositor
Map size (Å)	291.072, 291.072, 291.072	wwPDB
Map dimensions	256, 256, 256	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.137, 1.137, 1.137	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.24	0/3120	0.46	0/4251
1	B	0.25	0/3115	0.45	0/4244
All	All	0.25	0/6235	0.45	0/8495

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3058	0	3076	28	0
1	B	3053	0	3071	32	0
2	C	39	0	34	0	0
2	D	39	0	34	1	0
2	E	39	0	34	0	0
2	F	38	0	31	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
All	All	6268	0	6280	60	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (60) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:305:ILE:HG13	1:B:306:SER:H	1.55	0.71
1:A:156:ARG:HH12	1:A:344:LEU:HB2	1.61	0.65
1:A:156:ARG:NH1	1:A:343:ASP:OD1	2.30	0.65
1:B:291:SER:OG	1:B:295:ARG:NH2	2.32	0.63
1:B:156:ARG:HH21	1:B:344:LEU:HB2	1.66	0.59
1:B:56:LEU:HD11	1:B:254:ILE:HG22	1.86	0.58
1:A:242:LEU:O	1:A:246:LEU:HG	2.04	0.57
1:A:125:LEU:HD22	1:A:198:LEU:HD11	1.86	0.57
1:A:89:VAL:O	1:A:412:ARG:NH2	2.37	0.56
1:B:111:ASN:OD1	1:B:212:ARG:NH1	2.39	0.55
1:B:236:ARG:HD2	1:B:239:LYS:HG3	1.89	0.54
1:A:67:LEU:HD11	1:A:411:PRO:HG3	1.91	0.53
1:B:170:THR:OG1	1:B:325:GLN:OE1	2.26	0.53
1:A:241:LEU:O	1:A:245:MET:HG2	2.09	0.53
1:B:305:ILE:HG13	1:B:306:SER:N	2.24	0.51
1:A:18:HIS:ND1	1:A:272:ASP:OD1	2.44	0.51
1:A:30:VAL:HG21	1:A:44:SER:HB2	1.94	0.50
1:A:114:THR:HB	1:A:388:GLY:HA3	1.94	0.49
1:A:266:ALA:O	1:A:270:THR:HG22	2.13	0.49
1:A:103:ALA:O	1:A:395:PHE:HD1	1.95	0.49
1:A:282:LEU:HD11	1:A:295:ARG:HH11	1.77	0.49
1:A:260:LEU:HB2	1:A:396:SER:OG	2.13	0.49
1:A:349:LEU:HB2	2:D:1:NAG:H82	1.96	0.48
1:B:121:LEU:HD12	1:B:381:ILE:HG12	1.96	0.47
1:B:197:GLN:O	1:B:200:LYS:HG2	2.14	0.47
1:B:166:ASP:N	1:B:166:ASP:OD1	2.47	0.47
1:B:209:GLU:OE1	1:B:212:ARG:NH2	2.48	0.46
1:B:277:PRO:HG3	1:B:378:LEU:HD11	1.97	0.46
1:A:245:MET:HA	1:A:248:CYS:SG	2.56	0.46
1:A:148:VAL:O	1:A:151:GLU:HG3	2.16	0.46
1:B:273:PHE:CG	1:B:273:PHE:O	2.69	0.46
1:A:56:LEU:HD11	1:A:254:ILE:HG22	1.99	0.45
1:B:24:GLY:HA3	1:B:28:GLN:HE21	1.82	0.45
1:B:57:GLY:O	1:B:61:ILE:HG12	2.17	0.45
1:B:147:LYS:HD3	1:B:147:LYS:HA	1.78	0.44
1:B:251:LEU:HA	1:B:254:ILE:HG12	1.99	0.44
1:B:260:LEU:HB2	1:B:396:SER:OG	2.18	0.44
1:B:375:LEU:HD23	1:B:375:LEU:HA	1.85	0.44
1:A:322:MET:SD	1:A:351:LEU:HD11	2.58	0.43
1:B:172:LYS:O	1:B:176:GLN:HG2	2.17	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:13:TRP:CD1	1:B:14:VAL:HG23	2.54	0.43
1:B:114:THR:HB	1:B:388:GLY:HA3	2.01	0.43
1:B:156:ARG:HE	1:B:344:LEU:HB2	1.82	0.43
1:A:197:GLN:O	1:A:200:LYS:HG2	2.18	0.43
1:B:316:GLN:OE1	1:B:358:LEU:HD23	2.18	0.43
1:A:42:GLN:O	1:A:46:LEU:HG	2.18	0.42
1:A:13:TRP:CD1	1:A:14:VAL:HG23	2.53	0.42
1:B:34:PHE:CG	1:B:273:PHE:HB2	2.54	0.42
1:A:305:ILE:O	1:A:306:SER:OG	2.35	0.42
1:A:121:LEU:HD22	1:A:205:THR:HG21	2.02	0.41
1:B:224:LEU:HD23	1:B:224:LEU:HA	1.85	0.41
1:A:12:TRP:HA	1:A:15:VAL:HG12	2.03	0.41
1:B:31:ASP:OD1	1:B:31:ASP:N	2.51	0.41
1:B:145:ARG:NH1	1:B:353:ASN:HB3	2.35	0.41
1:A:224:LEU:HD23	1:A:224:LEU:HA	1.85	0.41
1:A:236:ARG:O	1:A:236:ARG:HG3	2.20	0.41
1:B:148:VAL:O	1:B:151:GLU:HG3	2.21	0.41
1:B:114:THR:HG23	1:B:212:ARG:NH2	2.36	0.40
1:B:156:ARG:NH2	1:B:344:LEU:HB2	2.35	0.40
1:A:167:TYR:O	1:A:171:LEU:HD23	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	392/540 (73%)	377 (96%)	15 (4%)	0	100	100
1	B	391/540 (72%)	378 (97%)	13 (3%)	0	100	100
All	All	783/1080 (72%)	755 (96%)	28 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	335/465 (72%)	335 (100%)	0	100	100
1	B	335/465 (72%)	335 (100%)	0	100	100
All	All	670/930 (72%)	670 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

12 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	C	1	1,2	14,14,15	0.35	0	17,19,21	0.40	0
2	NAG	C	2	2	14,14,15	0.21	0	17,19,21	0.42	0
2	BMA	C	3	2	11,11,12	0.58	0	15,15,17	0.72	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	D	1	1,2	14,14,15	0.25	0	17,19,21	0.47	0
2	NAG	D	2	2	14,14,15	0.41	0	17,19,21	0.35	0
2	BMA	D	3	2	11,11,12	0.60	0	15,15,17	0.74	0
2	NAG	E	1	1,2	14,14,15	0.34	0	17,19,21	0.57	0
2	NAG	E	2	2	14,14,15	0.24	0	17,19,21	0.37	0
2	BMA	E	3	2	11,11,12	0.57	0	15,15,17	0.78	0
2	NAG	F	1	1,2	14,14,15	0.23	0	17,19,21	0.54	0
2	NAG	F	2	2	14,14,15	0.41	0	17,19,21	0.34	0
2	BMA	F	3	2	10,10,12	0.63	0	14,14,17	0.59	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	C	1	1,2	-	0/6/23/26	0/1/1/1
2	NAG	C	2	2	-	2/6/23/26	0/1/1/1
2	BMA	C	3	2	-	0/2/19/22	0/1/1/1
2	NAG	D	1	1,2	-	2/6/23/26	0/1/1/1
2	NAG	D	2	2	-	2/6/23/26	0/1/1/1
2	BMA	D	3	2	-	0/2/19/22	0/1/1/1
2	NAG	E	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	E	2	2	-	2/6/23/26	0/1/1/1
2	BMA	E	3	2	-	1/2/19/22	0/1/1/1
2	NAG	F	1	1,2	-	1/6/23/26	0/1/1/1
2	NAG	F	2	2	-	2/6/23/26	0/1/1/1
2	BMA	F	3	2	-	-	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (13) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	C	2	NAG	O5-C5-C6-O6
2	D	1	NAG	O5-C5-C6-O6
2	F	2	NAG	O5-C5-C6-O6
2	D	1	NAG	C4-C5-C6-O6
2	F	2	NAG	C4-C5-C6-O6

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
2	C	2	NAG	C4-C5-C6-O6
2	E	2	NAG	C4-C5-C6-O6
2	D	2	NAG	O5-C5-C6-O6
2	E	2	NAG	O5-C5-C6-O6
2	D	2	NAG	C4-C5-C6-O6
2	E	3	BMA	O5-C5-C6-O6
2	E	1	NAG	C3-C2-N2-C7
2	F	1	NAG	C3-C2-N2-C7



There are no ring outliers.


1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.







5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-24688. These allow visual inspection of the internal detail of the map and identification of artifacts.

No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map




The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 129



Y Index: 134




Z Index: 173


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal surface views [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the 3D surface view of the map at the recommended contour level 7.0. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.5 Mask visualisation

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 46 nm³; this corresponds to an approximate mass of 42 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum [i](#)



*Reported resolution corresponds to spatial frequency of 0.286\AA^{-1}


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.

9 Map-model fit [i](#)


This section contains information regarding the fit between EMDB map EMD-24688 and PDB model 7RTT. Per-residue inclusion information can be found in section 3 on page 5.

9.1 Map-model overlay [i](#)




The images above show the 3D surface view of the map at the recommended contour level 7.0 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [\(i\)](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [\(i\)](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (7.0).















9.4 Atom inclusion [i](#)



At the recommended contour level, 71% of all backbone atoms, 68% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary [i](#)

The table lists the average atom inclusion at the recommended contour level (7.0) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.6790	 0.4330
A	 0.6848	 0.4360
B	 0.6870	 0.4360
C	 0.4103	 0.3840
D	 0.4359	 0.3260
E	 0.3846	 0.2090
F	 0.4211	 0.3560

