



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 17, 2024 – 07:20 AM EST

PDB ID : 3RUF
Title : Alternative analogs as viable substrates of UDP-hexose 4-epimerases
Authors : Bhatt, V.S.; Guan, W.; Wang, P.G.
Deposited on : 2011-05-05
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.36
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	8085 (2.00-2.00)
Ramachandran outliers	138981	9054 (2.00-2.00)
Sidechain outliers	138945	9053 (2.00-2.00)
RSRZ outliers	127900	7900 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	351	 1% 89% 6% . .
1	B	351	 2% 89% 6% . .
1	S	351	 3% 92% . .
1	b	351	 1% 90% 5% .

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 11427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called WbgU.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2681	1712	459	502	8	0	0	0
1	B	336	2681	1712	459	502	8	0	0	0
1	S	336	2681	1712	459	502	8	0	0	0
1	b	336	2674	1706	459	501	8	0	0	0

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-8	HIS	-	expression tag	UNP Q7BJX9
A	-7	HIS	-	expression tag	UNP Q7BJX9
A	-6	HIS	-	expression tag	UNP Q7BJX9
A	-5	HIS	-	expression tag	UNP Q7BJX9
A	-4	HIS	-	expression tag	UNP Q7BJX9
A	-3	HIS	-	expression tag	UNP Q7BJX9
B	-8	HIS	-	expression tag	UNP Q7BJX9
B	-7	HIS	-	expression tag	UNP Q7BJX9
B	-6	HIS	-	expression tag	UNP Q7BJX9
B	-5	HIS	-	expression tag	UNP Q7BJX9
B	-4	HIS	-	expression tag	UNP Q7BJX9
B	-3	HIS	-	expression tag	UNP Q7BJX9
S	-8	HIS	-	expression tag	UNP Q7BJX9
S	-7	HIS	-	expression tag	UNP Q7BJX9
S	-6	HIS	-	expression tag	UNP Q7BJX9
S	-5	HIS	-	expression tag	UNP Q7BJX9
S	-4	HIS	-	expression tag	UNP Q7BJX9
S	-3	HIS	-	expression tag	UNP Q7BJX9
b	-8	HIS	-	expression tag	UNP Q7BJX9
b	-7	HIS	-	expression tag	UNP Q7BJX9
b	-6	HIS	-	expression tag	UNP Q7BJX9

Continued on next page...



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
3	A	1	25	9	2	12	2	0	0
3	B	1	25	9	2	12	2	0	0
3	S	1	25	9	2	12	2	0	0
3	b	1	25	9	2	12	2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	111	Total	O	0	0
			111	111		
5	B	115	Total	O	0	0
			115	115		
5	S	101	Total	O	0	0
			101	101		
5	b	102	Total	O	0	0
			102	102		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: WbgU




- Molecule 1: WbgU



- Molecule 1: WbgU



- Molecule 1: WbgU



4 Data and refinement statistics i

Property	Value	Source
Space group	P 32	Depositor
Cell constants a, b, c, α , β , γ	77.98Å 77.98Å 223.87Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	37.00 – 2.00 37.32 – 1.48	Depositor EDS
% Data completeness (in resolution range)	99.1 (37.00-2.00) 58.4 (37.32-1.48)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.00 (at 1.48Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.204 , 0.248 0.206 , 0.247	Depositor DCC
R_{free} test set	7537 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	36.4	Xtrriage
Anisotropy	0.025	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 32.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.021 for -h,-k,l 0.437 for h,-h-k,-l 0.027 for -k,-h,-l	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	11427	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAD, UDP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.10	9/2739 (0.3%)	0.92	2/3717 (0.1%)
1	B	1.17	11/2739 (0.4%)	0.96	8/3717 (0.2%)
1	S	1.03	0/2739	0.91	5/3717 (0.1%)
1	b	1.05	3/2731 (0.1%)	0.92	7/3706 (0.2%)
All	All	1.09	23/10948 (0.2%)	0.93	22/14857 (0.1%)

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	318	LYS	CD-CE	12.09	1.81	1.51
1	A	7	GLU	CB-CG	8.94	1.69	1.52
1	B	7	GLU	CB-CG	8.07	1.67	1.52
1	B	318	LYS	CB-CG	7.54	1.72	1.52
1	B	318	LYS	CE-NZ	7.39	1.67	1.49
1	A	186	PHE	CE1-CZ	6.32	1.49	1.37
1	B	1	TYR	CD1-CE1	6.19	1.48	1.39
1	A	67	GLU	CG-CD	5.92	1.60	1.51
1	A	49	PHE	CE2-CZ	5.72	1.48	1.37
1	A	5	TYR	CD2-CE2	5.70	1.47	1.39
1	b	43	VAL	CB-CG1	5.67	1.64	1.52
1	B	116	ALA	CA-CB	5.67	1.64	1.52
1	A	25	ALA	CA-CB	5.53	1.64	1.52
1	B	25	ALA	CA-CB	5.51	1.64	1.52
1	B	67	GLU	CG-CD	5.38	1.60	1.51
1	b	172	VAL	CB-CG2	5.37	1.64	1.52
1	A	176	TYR	CD2-CE2	5.31	1.47	1.39
1	A	5	TYR	CD1-CE1	5.23	1.47	1.39
1	B	318	LYS	CG-CD	5.15	1.70	1.52
1	B	137	PHE	CD1-CE1	5.14	1.49	1.39
1	A	137	PHE	CE2-CZ	5.09	1.47	1.37
1	b	186	PHE	CE2-CZ	5.04	1.47	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	186	PHE	CE1-CZ	5.00	1.46	1.37

All (22) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	258	ASP	CB-CG-OD2	7.34	124.91	118.30
1	B	299	ARG	NE-CZ-NH1	7.24	123.92	120.30
1	B	239	ILE	CB-CA-C	-6.57	98.45	111.60
1	S	240	ASP	CB-CG-OD1	6.31	123.98	118.30
1	B	260	ILE	CG1-CB-CG2	-6.29	97.57	111.40
1	b	240	ASP	CB-CG-OD1	6.19	123.87	118.30
1	S	148	ASP	CB-CG-OD1	5.93	123.63	118.30
1	A	239	ILE	CB-CA-C	-5.82	99.97	111.60
1	B	239	ILE	CG1-CB-CG2	5.67	123.87	111.40
1	S	154	LYS	CD-CE-NZ	-5.63	98.75	111.70
1	b	96	LEU	CB-CG-CD2	5.56	120.46	111.00
1	A	235	ASP	CB-CG-OD1	5.39	123.16	118.30
1	B	299	ARG	NE-CZ-NH2	-5.34	117.63	120.30
1	B	318	LYS	CG-CD-CE	5.32	127.84	111.90
1	b	195	ASN	N-CA-C	5.28	125.25	111.00
1	b	304	ARG	NE-CZ-NH1	5.26	122.93	120.30
1	b	148	ASP	CB-CG-OD1	5.23	123.01	118.30
1	B	318	LYS	CB-CA-C	5.22	120.83	110.40
1	S	304	ARG	NE-CZ-NH1	5.21	122.91	120.30
1	b	304	ARG	NE-CZ-NH2	-5.03	117.78	120.30
1	S	179	VAL	CG1-CB-CG2	-5.02	102.87	110.90
1	b	239	ILE	CB-CA-C	-5.01	101.57	111.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

Due to software issues we are unable to calculate clashes - this section is therefore empty.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	332/351 (95%)	321 (97%)	10 (3%)	1 (0%)	41	37
1	B	332/351 (95%)	323 (97%)	7 (2%)	2 (1%)	25	19
1	S	332/351 (95%)	321 (97%)	9 (3%)	2 (1%)	25	19
1	b	332/351 (95%)	322 (97%)	8 (2%)	2 (1%)	25	19
All	All	1328/1404 (95%)	1287 (97%)	34 (3%)	7 (0%)	29	23

All (7) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	b	286	HIS
1	S	3	SER
1	B	286	HIS
1	b	103	SER
1	A	103	SER
1	B	103	SER
1	S	103	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	290/304 (95%)	275 (95%)	15 (5%)	23	19
1	B	290/304 (95%)	276 (95%)	14 (5%)	25	22
1	S	290/304 (95%)	283 (98%)	7 (2%)	49	51
1	b	289/304 (95%)	279 (96%)	10 (4%)	36	35
All	All	1159/1216 (95%)	1113 (96%)	46 (4%)	31	29

All (46) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	46	LEU
1	A	67	GLU
1	A	109	VAL
1	A	152	LEU
1	A	157	GLU
1	A	183	THR
1	A	239	ILE
1	A	253	LYS
1	A	268	ARG
1	A	273	GLU
1	A	299	ARG
1	A	315	ASP
1	A	318	LYS
1	A	320	ARG
1	A	330	ARG
1	B	67	GLU
1	B	109	VAL
1	B	117	THR
1	B	152	LEU
1	B	157	GLU
1	B	176	TYR
1	B	239	ILE
1	B	279	TYR
1	B	283	ASN
1	B	299	ARG
1	B	315	ASP
1	B	324	LYS
1	B	326	ARG
1	B	338	ARG
1	S	71	ARG
1	S	157	GLU
1	S	179	VAL
1	S	183	THR
1	S	239	ILE
1	S	279	TYR
1	S	338	ARG
1	b	3	SER
1	b	15	SER
1	b	46	LEU
1	b	71	ARG
1	b	130	LYS
1	b	157	GLU
1	b	239	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	b	318	LYS
1	b	338	ARG
1	b	341	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (18) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	11	GLN
1	A	41	GLN
1	A	149	HIS
1	A	158	ASN
1	A	173	ASN
1	A	178	GLN
1	A	195	ASN
1	B	41	GLN
1	B	149	HIS
1	B	195	ASN
1	S	149	HIS
1	S	158	ASN
1	S	173	ASN
1	S	195	ASN
1	S	283	ASN
1	b	149	HIS
1	b	173	ASN
1	b	195	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry

9 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAD	A	343	-	42,48,48	1.18	3 (7%)	50,73,73	1.85	11 (22%)
3	UDP	A	344	-	24,26,26	1.19	2 (8%)	37,40,40	1.83	11 (29%)
3	UDP	B	344	-	24,26,26	1.16	1 (4%)	37,40,40	1.73	8 (21%)
4	SO4	A	345	-	4,4,4	1.45	0	6,6,6	0.62	0
3	UDP	b	343	-	24,26,26	0.96	1 (4%)	37,40,40	1.55	8 (21%)
2	NAD	b	344	-	42,48,48	1.21	4 (9%)	50,73,73	1.64	11 (22%)
2	NAD	B	343	-	42,48,48	1.08	3 (7%)	50,73,73	1.80	10 (20%)
2	NAD	S	343	-	42,48,48	1.07	1 (2%)	50,73,73	1.63	7 (14%)
3	UDP	S	344	-	24,26,26	1.13	3 (12%)	37,40,40	1.51	8 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAD	A	343	-	-	6/26/62/62	0/5/5/5
3	UDP	A	344	-	-	1/16/32/32	0/2/2/2
3	UDP	B	344	-	-	2/16/32/32	0/2/2/2
3	UDP	b	343	-	-	1/16/32/32	0/2/2/2
2	NAD	b	344	-	-	6/26/62/62	0/5/5/5
2	NAD	B	343	-	-	6/26/62/62	0/5/5/5
2	NAD	S	343	-	-	6/26/62/62	0/5/5/5
3	UDP	S	344	-	-	1/16/32/32	0/2/2/2

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	343	NAD	O4B-C1B	4.27	1.47	1.41
3	A	344	UDP	C2-N1	3.79	1.44	1.38
2	b	344	NAD	C7N-N7N	3.08	1.38	1.33
3	B	344	UDP	C2-N1	3.06	1.43	1.38
2	S	343	NAD	C7N-N7N	2.84	1.38	1.33
2	B	343	NAD	C2N-N1N	2.63	1.38	1.35
2	B	343	NAD	O4B-C1B	2.59	1.44	1.41
3	S	344	UDP	C6-C5	2.52	1.40	1.35
3	b	343	UDP	C6-C5	2.42	1.40	1.35
3	S	344	UDP	O2-C2	2.40	1.27	1.23
3	A	344	UDP	C6-C5	2.32	1.40	1.35
2	b	344	NAD	C2A-N3A	2.31	1.35	1.32
2	B	343	NAD	C2A-N3A	2.27	1.35	1.32
2	A	343	NAD	C4A-N3A	2.24	1.38	1.35
2	b	344	NAD	C2N-N1N	2.21	1.37	1.35
2	A	343	NAD	O3D-C3D	2.14	1.48	1.43
2	b	344	NAD	C3B-C4B	2.06	1.58	1.53
3	S	344	UDP	PB-O3B	2.06	1.62	1.54

All (74) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	343	NAD	N3A-C2A-N1A	-7.05	117.67	128.68
2	S	343	NAD	N3A-C2A-N1A	-6.48	118.55	128.68
2	B	343	NAD	N3A-C2A-N1A	-6.11	119.13	128.68
2	b	344	NAD	N3A-C2A-N1A	-5.02	120.83	128.68
2	A	343	NAD	O7N-C7N-C3N	-4.90	113.77	119.63
2	b	344	NAD	O7N-C7N-C3N	-4.37	114.40	119.63
3	A	344	UDP	O3B-PB-O3A	-4.37	89.98	104.64
2	B	343	NAD	O2N-PN-O1N	4.22	133.10	112.24
2	A	343	NAD	O2N-PN-O1N	4.16	132.79	112.24
2	B	343	NAD	O7N-C7N-C3N	-4.00	114.84	119.63
3	b	343	UDP	C4-N3-C2	-3.80	121.57	126.58
3	A	344	UDP	N3-C2-N1	3.73	119.84	114.89
3	B	344	UDP	O3B-PB-O3A	-3.69	92.27	104.64
3	A	344	UDP	C4-N3-C2	-3.64	121.78	126.58
3	A	344	UDP	C5-C4-N3	3.58	120.20	114.84
3	b	343	UDP	C5-C4-N3	3.56	120.16	114.84
3	B	344	UDP	O4-C4-C5	-3.53	118.96	125.16
3	B	344	UDP	C4-N3-C2	-3.37	122.14	126.58
3	S	344	UDP	C5-C4-N3	3.27	119.74	114.84
3	S	344	UDP	O2B-PB-O1B	3.25	123.40	110.68
2	b	344	NAD	C2N-C3N-C4N	3.25	121.94	118.26

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	343	NAD	C2N-C3N-C4N	3.20	121.88	118.26
2	B	343	NAD	O4B-C1B-C2B	-3.19	102.27	106.93
3	B	344	UDP	N3-C2-N1	3.17	119.10	114.89
2	S	343	NAD	O2N-PN-O1N	3.13	127.69	112.24
2	A	343	NAD	C3N-C2N-N1N	-3.12	117.38	120.43
3	B	344	UDP	C1'-N1-C2	3.08	123.14	117.57
3	S	344	UDP	C4-N3-C2	-3.04	122.57	126.58
2	b	344	NAD	O7N-C7N-N7N	3.04	126.90	122.58
2	B	343	NAD	C3N-C7N-N7N	2.99	121.34	117.75
3	b	343	UDP	N3-C2-N1	2.94	118.79	114.89
2	A	343	NAD	C2A-N1A-C6A	2.94	123.78	118.75
2	S	343	NAD	O7N-C7N-C3N	-2.93	116.13	119.63
3	A	344	UDP	O4-C4-C5	-2.88	120.09	125.16
3	A	344	UDP	O2B-PB-O1B	2.84	121.81	110.68
2	S	343	NAD	O3D-C3D-C4D	-2.81	102.91	111.05
2	B	343	NAD	O4D-C1D-C2D	-2.79	102.86	106.93
3	S	344	UDP	O3B-PB-O3A	-2.78	95.31	104.64
2	A	343	NAD	C3N-C7N-N7N	2.77	121.07	117.75
3	B	344	UDP	C5-C4-N3	2.76	118.97	114.84
2	S	343	NAD	O2B-C2B-C3B	2.66	120.43	111.82
3	b	343	UDP	O2-C2-N1	-2.65	119.26	122.79
2	A	343	NAD	PN-O3-PA	-2.63	123.81	132.83
3	A	344	UDP	O2B-PB-O3A	2.60	113.34	104.64
2	b	344	NAD	C4A-C5A-N7A	-2.56	106.73	109.40
3	S	344	UDP	O2A-PA-O1A	2.53	124.74	112.24
3	A	344	UDP	C6-N1-C2	-2.52	117.76	120.99
2	b	344	NAD	C2A-N1A-C6A	2.50	123.02	118.75
3	B	344	UDP	O2B-PB-O3A	2.49	113.00	104.64
2	b	344	NAD	O2B-C2B-C3B	2.49	119.88	111.82
3	b	343	UDP	O4-C4-C5	-2.46	120.83	125.16
2	b	344	NAD	C3N-C2N-N1N	-2.44	118.04	120.43
2	B	343	NAD	PN-O3-PA	-2.43	124.48	132.83
3	b	343	UDP	O2B-PB-O1B	2.43	120.20	110.68
2	A	343	NAD	O4D-C1D-C2D	-2.43	103.37	106.93
3	B	344	UDP	O2B-PB-O1B	2.38	120.02	110.68
3	b	343	UDP	O3'-C3'-C2'	-2.34	104.25	111.82
3	S	344	UDP	N3-C2-N1	2.33	117.99	114.89
2	B	343	NAD	O2N-PN-O5D	-2.32	96.96	107.75
3	S	344	UDP	O4-C4-C5	-2.32	121.08	125.16
3	b	343	UDP	O3B-PB-O3A	-2.31	96.89	104.64
2	b	344	NAD	C5A-C6A-N6A	2.24	123.76	120.35
2	b	344	NAD	O3D-C3D-C4D	-2.23	104.60	111.05

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	S	343	NAD	C2A-N1A-C6A	2.21	122.54	118.75
2	S	343	NAD	C3D-C2D-C1D	-2.20	97.67	100.98
2	A	343	NAD	C2N-C3N-C4N	2.16	120.71	118.26
2	B	343	NAD	C3N-C2N-N1N	-2.14	118.34	120.43
2	A	343	NAD	O2N-PN-O5D	-2.12	97.90	107.75
2	A	343	NAD	O4B-C1B-C2B	-2.10	103.86	106.93
3	A	344	UDP	C1'-N1-C2	2.07	121.31	117.57
3	A	344	UDP	O3B-PB-O2B	-2.05	99.80	107.64
3	A	344	UDP	O2'-C2'-C1'	2.01	116.76	110.02
3	S	344	UDP	O3'-C3'-C2'	-2.01	105.31	111.82
2	b	344	NAD	C5N-C4N-C3N	-2.01	117.96	120.34

There are no chirality outliers.

All (29) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	343	NAD	C5D-O5D-PN-O1N
2	B	343	NAD	PN-O3-PA-O5B
2	B	343	NAD	C5D-O5D-PN-O1N
2	S	343	NAD	C5D-O5D-PN-O1N
2	b	344	NAD	C5D-O5D-PN-O1N
3	B	344	UDP	PB-O3A-PA-O5'
2	A	343	NAD	PN-O3-PA-O5B
2	b	344	NAD	PN-O3-PA-O5B
3	A	344	UDP	PB-O3A-PA-O5'
3	S	344	UDP	PB-O3A-PA-O5'
3	b	343	UDP	PB-O3A-PA-O5'
2	A	343	NAD	PA-O3-PN-O1N
2	A	343	NAD	C5D-O5D-PN-O2N
2	B	343	NAD	C5D-O5D-PN-O2N
2	S	343	NAD	C5D-O5D-PN-O2N
2	b	344	NAD	C5D-O5D-PN-O2N
2	B	343	NAD	PA-O3-PN-O1N
2	b	344	NAD	O4B-C4B-C5B-O5B
2	B	343	NAD	O4B-C4B-C5B-O5B
2	A	343	NAD	O4B-C4B-C5B-O5B
3	B	344	UDP	PA-O3A-PB-O3B
2	A	343	NAD	C5D-O5D-PN-O3
2	B	343	NAD	C5D-O5D-PN-O3
2	S	343	NAD	C5D-O5D-PN-O3
2	b	344	NAD	C5D-O5D-PN-O3
2	S	343	NAD	O4B-C4B-C5B-O5B

Continued on next page...


Continued from previous page...



Mol	Chain	Res	Type	Atoms
2	S	343	NAD	PA-O3-PN-O1N
2	S	343	NAD	PA-O3-PN-O2N
2	b	344	NAD	PA-O3-PN-O2N



There are no ring outliers.


No monomer is involved in short contacts.



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	336/351 (95%)	-0.27	4 (1%) 79 78	27, 38, 66, 82	0
1	B	336/351 (95%)	-0.25	6 (1%) 68 66	28, 39, 68, 83	0
1	S	336/351 (95%)	-0.15	9 (2%) 54 53	29, 41, 67, 92	0
1	b	336/351 (95%)	-0.12	5 (1%) 73 72	28, 41, 66, 85	0
All	All	1344/1404 (95%)	-0.20	24 (1%) 68 66	27, 40, 67, 92	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	b	2	MET	8.9
1	S	2	MET	6.0
1	B	286	HIS	4.9
1	B	301	GLY	4.6
1	b	1	TYR	4.6
1	A	301	GLY	4.4
1	S	1	TYR	4.4
1	B	225	TYR	4.0
1	b	286	HIS	3.8
1	b	300	SER	3.7
1	S	300	SER	3.7
1	B	279	TYR	3.6
1	A	282	LEU	3.6
1	S	340	LEU	3.1
1	A	293	ILE	2.9
1	B	219	LEU	2.8
1	A	286	HIS	2.6
1	B	285	ILE	2.5
1	S	10	GLN	2.5
1	S	297	GLU	2.5
1	S	338	ARG	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	S	286	HIS	2.2
1	b	337	VAL	2.0
1	S	279	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	UDP	A	344	25/25	0.91	0.12	47,50,53,54	0
3	UDP	b	343	25/25	0.92	0.09	38,49,52,52	0
2	NAD	b	344	44/44	0.94	0.10	30,36,41,45	0
2	NAD	A	343	44/44	0.95	0.10	23,31,38,41	0
2	NAD	B	343	44/44	0.95	0.10	27,32,39,40	0
3	UDP	B	344	25/25	0.95	0.09	45,51,53,54	0
3	UDP	S	344	25/25	0.95	0.09	42,48,49,52	0
2	NAD	S	343	44/44	0.95	0.11	29,35,40,44	0
4	SO4	A	345	5/5	0.99	0.11	30,30,30,30	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.


Electron density around UDP A 344:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around UDP b 343:**




$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)








Electron density around NAD B 343:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around UDP B 344:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around UDP S 344:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAD S 343:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.